

In order to see the correspondence of the fields with IIB matrix model, we express the bosonic 32×32 matrices in terms of the **10-dimensional indices**.

$(\mu, \nu, \dots = 0, 1, \dots, 9, \sharp = 10)$.

$$W = m_{\sharp}, A_{\mu} = m_{\mu}, B_{\mu} = m_{\mu\sharp}, C_{\mu_1\mu_2} = m_{\mu_1\mu_2}, \\ H_{\mu_1\cdots\mu_4} = m_{\mu_1\cdots\mu_4\sharp}, Z_{\mu_1\cdots\mu_5} = m_{\mu_1\cdots\mu_5}.$$

Then, the action is decomposed as

$$S = 96\mu Tr \left(-W^2 - A_{\mu}A^{\mu} + B_{\mu}B^{\mu} + \frac{1}{2}C_{\mu_1\mu_2}C^{\mu_1\mu_2} \right. \\ \left. - \frac{1}{4!}H_{\mu_1\cdots\mu_4}H^{\mu_1\cdots\mu_4} - \frac{1}{5!}Z_{\mu_1\cdots\mu_5}Z^{\mu_1\cdots\mu_5} + \frac{i}{16}\bar{\psi}\psi \right) \\ + 32i Tr \left(3C_{\mu_1\mu_2}[B^{\mu_1}, B^{\mu_2}] + C_{\mu_1\mu_2}[C^{\mu_2\mu_3}, C^{\mu_3\mu_1}] \right) \\ + [\text{cubic interactions involving } (W, A, H, Z, \psi)].$$

- The rank-1 and rank-5 fields (in 11 dimensions) have a positive mass, while the rank-2 fields are tachyonic.

$$\underbrace{\Gamma_A \Gamma^A}_{\text{no sum}} = \underbrace{\Gamma_{A_1\cdots A_5} \Gamma^{A_1\cdots A_5}}_{\text{no sum}} = +\mathbf{1}_{32 \times 32}, \\ \underbrace{\Gamma_{A_1 A_2} \Gamma^{A_1 A_2}}_{\text{no sum}} = -\mathbf{1}_{32 \times 32}.$$

- The rank-1 and rank-5 fields has a stable trivial commutative classical solution:

$$W = A_{\mu} = H_{\mu_1\cdots\mu_4} = Z_{\mu_1\cdots\mu_5} = 0.$$

- For the rank-2 tachyonic fields $B_{\mu}, C_{\mu_1\mu_2}$, the trivial solution $B_{\mu} = C_{\mu_1\mu_2} = 0$ is unstable. \Rightarrow They may incorporate an interesting stable non-commutative solution!

4. Resolution of the equations of motion

From now on, we set the fermions and the positive-mass-square bosonic fields to zero:

$$S = 96\mu Tr \left(B_{\mu}B^{\mu} + \frac{1}{2}C_{\mu_1\mu_2}C^{\mu_1\mu_2} \right) \\ + 32i Tr \left(3C_{\mu_1\mu_2}[B^{\mu_1}, B^{\mu_2}] + C_{\mu_1\mu_2}[C^{\mu_2\mu_3}, C^{\mu_3\mu_1}] \right).$$

The equations of motion:

$$B_{\mu} = -i\mu^{-1}[B^{\nu}, C_{\mu\nu}], \\ C_{\mu_1\mu_2} = -i\mu^{-1}([B_{\mu_1}, B_{\mu_2}] + [C_{\mu_1}{}^{\rho}, C_{\mu_2\rho}]).$$

We integrate out the rank-2 fields (in 10 dimensions) $C_{\mu_1\mu_2}$ by solving the latter equation of motions iteratively.

$$C_{\mu_1\mu_2} = -i\mu^{-1}[B_{\mu_1}, B_{\mu_2}] + i\mu^{-3}[[B_{\mu_1}, B_{\rho}], [B_{\mu_2}, B^{\rho}]] \\ - 2i\mu^{-5}[[B_{\mu_1}, B_{\rho}], [[B_{\mu_2}, B_{\chi}], [B^{\rho}, B^{\chi}]]] + \mathcal{O}(\mu^{-7}). (*)$$

Then, the action reduces to

$$S = Tr \left(96\mu B_{\mu}B^{\mu} + 48\mu^{-1}[B_{\mu_1}, B_{\mu_2}][B^{\mu_1}, B^{\mu_2}] \right) \\ + [\text{higher-order commutators of the order } \mathcal{O}(\mu^{-2k+1}) \text{ with } k = 2, 3, \dots].$$

We consider the classical solution of the equation of motion $B_{\mu} = -i\mu^{-1}[B^{\nu}, C_{\mu\nu}]$ with $C_{\mu_1\mu_2}$ substituted for (*).

Fuzzy-sphere classical solution

1. $SO(3) \times SO(3) \times SO(3)$ fuzzy spheres

This describes a space formed by the Cartesian product of three fuzzy spheres.

$$[B_i, B_j] = i\mu r \epsilon_{ijk} B_k, \quad B_1^2 + B_2^2 + B_3^2 = \mu^2 r^2 \frac{N^2 - 1}{4}, \quad (i, j, k = 1, 2, 3) \\ [B_l, B_m] = i\mu r \epsilon_{lmn} B_n, \quad B_4^2 + B_5^2 + B_6^2 = \mu^2 r^2 \frac{N^2 - 1}{4}, \quad (l, m, n = 4, 5, 6) \\ [B_p, B_q] = i\mu r \epsilon_{pqr} B_r, \quad B_7^2 + B_8^2 + B_9^2 = \mu^2 r^2 \frac{N^2 - 1}{4}, \quad (p, q, r = 7, 8, 9) \\ B_0 = 0, [B_{\mu}, B_{\nu}] = 0, \text{ (otherwise).}$$

(We consider the Cartesian product of three spheres instead of a single $SO(3)$ fuzzy sphere

$$[B_i, B_j] = i\mu r \epsilon_{ijk} B_k \text{ (for } i, j, k = 1, 2, 3), \\ B_{\mu} = 0 \text{ (for } \mu = 0, 4, 5, \dots, 9),$$

because the solution $B_4 = \dots = B_9 = 0$ is trivially unstable.)

2. $SO(9)$ fuzzy sphere

Generally, the $SO(2k+1)$ fuzzy sphere (S^{2k} fuzzy sphere) is constructed by the n -fold symmetric tensor product of $(2k+1)$ -dimensional gamma matrices:

$$B_p = \frac{\mu r}{2} [(\Gamma_p^{(2k)} \otimes \mathbf{1} \otimes \dots \otimes \mathbf{1}) + \dots + (\mathbf{1} \otimes \dots \otimes \mathbf{1} \otimes \Gamma_p^{(2k)})]_{\text{sym}}. \\ B_p B_p = \frac{\mu^2 r^2}{4} n(n+2k) \mathbf{1}_{N_k \times N_k}.$$

We start with the ansatz for the rank-2 fields C_{pq} for the $SO(2k+1)$ fuzzy spheres:

$$C_{pq} = -i\mu^{-1} f(r) B_{pq}.$$

Then, by solving the equations of motion for B_{μ} and $C_{\mu\nu}$ simultaneously, we obtain a constraint on the radius parameter r as

$$r = \frac{1}{2k}.$$

Comparison of the classical energy

- Trivial commutative solution $B_0 = \dots = B_9 = 0$:

$$E_{B_{\mu}=0} = -S_{B_{\mu}=0} = 0.$$

- $SO(3) \times SO(3) \times SO(3)$ fuzzy spheres ($N_1 = n+1$):

$$E_{SO(3)^3} = -S_{SO(3)^3} = -\frac{16\mu}{r_{SO(3)^3}} Tr(B_{\mu}B^{\mu}) \\ = -12\mu^3 N_1(N_1-1)(N_1+1) \\ \sim -\mathcal{O}(\mu^3 n^3) = -\mathcal{O}(\mu^3 N_1^3).$$

- $SO(9)$ fuzzy sphere:

$$E_{SO(9)} = -S_{SO(9)} = -\frac{5}{8}\mu^3 n(n+8)N_4 \\ \sim -\mathcal{O}(\mu^3 n^{12}) = -\mathcal{O}(\mu^3 N_4^{\frac{6}{5}}),$$

where the size of the matrices B_p is

$$N_4 = \frac{(n+1)(n+2)(n+3)^2(n+4)^2(n+5)^2(n+6)(n+7)}{302400} \sim \mathcal{O}(n^{10}).$$

When the size of the matrices are the same...

$(N_1 = N_4)$	E
0	Commutative Solution
$-N^{65}$	$SO(9)$ fuzzy sphere
$-N^3$	$SO(3) \times SO(3) \times SO(3)$ fuzzy spheres