Monte Carlo Studies of the GWW Phase Transition in Large-N Gauge Theories

(arXiv:0710.5873)

Takehiro Azuma [Tata Institite of Fundamental Research (TIFR)]

From Strings to LHC II, Dec. 20th 2007, 15:15 ∼ 15:45

Collaboration with Pallab Basu and Spenta R. Wadia

Contents

1 Introduction **2**

2 The model **4**

3 Gross-Witten-Wadia phase transition **5**

4 SO(6) R-symmetry breaking **9**

5 Conclusion **14**

1 Introduction

AdS/CFT correspondence: J. M. Maldacena, hep-th/9711200

duality between type IIB superstring on $AdS_5 \times S^5$ and $\mathcal{N} = 4$ SYM gauge theory.



- **Nonperturbative studies of superstring.**
- **Quantum description of the blackhole.**

Thermodynamic aspects of quantum gravity in AdS spacetime.

S. W. Hawking and D. N. Page, Commun. Math. Phys. 87, 577 (1983).

AdS spacetime allows two Schwarzschild blackhole solutions.

- **Small black hole (SBH): Horizon radius smaller than AdS.**
  Negative specific heat. Unstable.
- **Big black hole (BBH): horizon radius comparable to AdS. Positive specific heat.**

First-order phase transition between the AdS space and the BBH.

Hawking-Page transition corresponds to a large-N deconfinement transition in the gauge theory at strong coupling. E. Witten, hep-th/9803131.

Studies of blackhole physics on the SYM side.

2 The model

Zero-mode action of the bosonic sector of $\mathcal{N}=4$ SYM on S^3 at finite temperature.

$$Z = \int dM_\mu dA e^{-S}, \quad \text{where}$$
$$S = N \int_0^\beta dt \left\{ \mathrm{tr} \sum_{\mu=1}^{D} (D_t M_\mu(t))^2 - \frac{\lambda}{2} \mathrm{tr} \sum_{\mu,\nu=1}^{D} [M_\mu(t), M_\nu(t)]^2 + m^2 \mathrm{tr} \sum_{\mu=1}^{D} M_\mu^2(t) \right\}.$$

- $D_t M_\mu(t) = \partial_t M_\mu(t) - i[A, M_\mu(t)]$: covariant derivative
  (A = zero mode of the time component of the gauge field on S^3)
- $M_\mu(t)$: SO(6) scalar fields ($\mu, \nu, \cdots = 1, 2, \cdots, D$, here $D = 6$)
- U(N) gauge symmetry : $M_\mu(t) \to g(t) M_\mu(t) g^\dagger(t)$, $A \to g(t) A g^\dagger(t) + i g(t) \frac{dg(t)^\dagger}{dt}$
- $\beta = 1/T$: inverse temperature
  Periodic boundary condition : $A(t+\beta) = A(t)$, $M_\mu(t+\beta) = M_\mu(t)$.
- Static and diagonal gauge: $A = \frac{1}{\beta}\mathrm{diag}(\alpha_1, \alpha_2, \cdots, \alpha_N)$, where $\{\alpha_a\} \in [-\pi, \pi]$.
  Gauge fixing term : $S_{\text{g.f.}} = -\sum_{a,b=1, a\neq b}^{N} \log \sin |(\alpha_a - \alpha_b)/2|$.

3 Gross-Witten-Wadia phase transition

Effective action of the SYM theory on S^3 at finite temperature

→ Described by Polyakov line U.



Phase structure of the YM theory and blackhole states in supergravity.

L. Alvarez-Gaume, C. Gomez, H. Liu and S.R. Wadia hep-th/0502227

Gross-Witten-Wadia (GWW) third-order phase transition of the partition function

$$Z_g = \int dU \exp(\frac{Ng}{2}(\mathrm{tr}\, U + \mathrm{tr}\, U^\dagger)),$$

$$\langle |u_1| \rangle = \begin{cases} \frac{g}{2}, & (g < 1) \\ 1 - \frac{1}{2g} & (g > 1). \end{cases}$$

$$\langle |u_2| \rangle = \begin{cases} 0, & (g < 1) \\ 1 - \frac{2}{g} + \frac{1}{g^2} & (g > 1). \end{cases}$$

MC simulation for $N = 128$, $u_n = \frac{1}{N}\mathrm{tr}\, U^n$,

- $g < 1$: no gap on unit circle.
- $g > 1$: a system has a gap.

Saddle point of the gauge field for our model:

$$S' = S + N\mu \int_0^\beta dt(\mathrm{tr}\, U + \mathrm{tr}\, U^\dagger).$$

$D = 2$ **case**

High-temperature $(\beta = 0.2)$ **case** $(\lambda = m = 1.0,\ N = 48)$.



No GWW-type phase transition.

Low-temperature ($\beta = 2.0$) case ($\lambda = m = 1.0,\ N = 48$).





Critical point is $\mu_c \simeq 0.22$.

- $\langle |u_1| \rangle_{|\mu=0} = 0,\ \langle |u_1| \rangle_{|\mu=\infty} = 1.$
- At the critical point $\mu = \mu_c$, $\langle |u_1| \rangle$ and $\frac{d\langle |u_1| \rangle}{d\mu}$ are continuous.

$$\langle |u_1| \rangle = \begin{cases} q_1 \frac{\mu}{\mu_c} + r_1 (\frac{\mu}{\mu_c})^2, & (\mu < \mu_c),\ \ r_1 = \frac{1}{2}(1 - \frac{3}{2}q_1 - \frac{1}{2}q_2), \\ 1 - q_2 (\frac{\mu}{\mu_c})^{-1} - r_2 (\frac{\mu}{\mu_c})^{-2}, & (\mu > \mu_c),\ \ r_2 = \frac{1}{2}(1 - \frac{1}{2}q_1 - \frac{3}{2}q_2), \end{cases}$$

Parameters are fitted as

$$(q_1, q_2) = (0.503 \pm 0.011, 0.538 \pm 0.004) \Rightarrow (r_1, r_2) = (-0.0121, -0.0293).$$

r_1, r_2's contribution is small. **GWW-type third order phase transition.**

$$\langle |u_2| \rangle = 1 - \frac{2\mu_c}{\mu} + \frac{\mu_c^2}{\mu^2}, \quad (\mu > \mu_c).$$

$D = 6$ case

Low-temperature ($\beta = 2.0$) case ($\lambda = m = 1.0$, $N = 16$).



Critical point is $\mu_c \simeq 0.20$.

$$(q_1, q_2) = (0.538 \pm 0.020, 0.542 \pm 0.007) \Rightarrow (r_1, r_2) = (-0.039, -0.041).$$

The result is insensitive to the dimensionality.

4 SO(6) R-symmetry breaking

Supergravity-side analysis of the 10-dim small black hole (SBH) in $AdS_5 \times S^5$..

L. Alvarez-Gaume, P. Basu, M. Marino and S.R. Wadia, hep-th/0605041

Metric of 10-dim SBH in $AdS_5 \times S^5$ → not symmetric under SO(6) rotation of S^5.

$$\Updownarrow$$

Corresponding saddle point → transform under SO(6) R-symmetry group.

Supergravity analysis : SO(6) R-symmetry is not spontaneously broken.

How about in the gauge theory side?

⇒ Monte Carlo simulation of the gauge theory.

Order parameter for the SO(6) R-symmetry breaking

Eigenvalue $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_6$ of integrated "momentum of inertia"

$$I_{\mu\nu} = \frac{1}{N} \int_0^\beta dt \mathrm{tr}\, M_\mu(t) M_\nu(t).$$

(in analogy to the IKKT model's case) hep-th/9811220, 0003208, 0005147, 0104260, 0108041, 0108070, 0402194

Large-N extrapolation of the VEV $\langle \lambda_\mu \rangle$.

Large-N limit : Eigenvalues $\langle \lambda_\mu \rangle$ converge to the same value.

SO(6) R-symmetry is not broken.



Figure 1: $(\beta, \lambda) = (0.1, 0.1)$ (upper left), $(\beta, \lambda) = (1.0, 0.1)$ (upper right), $(\beta, \lambda) = (0.1, 1.0)$ (lower left), $(\beta, \lambda) = (1.0, 1.0)$ (lower right), for $N = 16, 20, 24, 28, 32$, $m = 1.0$.

Specific vacuum configurations of the gauge field



(1)uniform (2)clumped

- **Uniform distrobution :** $\alpha_a = \frac{\pi}{N}(2a - N)$ $(a = 1, 2, \cdots, N)$

  $\rightarrow \langle \mathrm{tr}\, U^n \rangle = 0$ for any $n \neq 0 \rightarrow$ Corresponds to $\mathrm{AdS}_5 \times \mathrm{S}^5$ geometry.

  E. Witten hep-th/9803131

- **Clumped distribution : blackhole state**

U: zero mode of the Polyakov line on S^3:

$$u_n = \frac{1}{N}\mathrm{tr}\, U^n = \frac{1}{N}\sum_{a=1}^{N} e^{i\alpha_a}, \quad U = P\exp\left(i\int_0^\beta dt A(t)\right).$$

SO(6) R-symmetry is not broken in these cases, too.



Figure 2: Uniform (left) and clumped (right) distribution for $\boldsymbol{\beta = 0.1}$, $\boldsymbol{\lambda = 1.0}$, $\boldsymbol{m = 1.0}$.

5 Conclusion

Zero mode effective action of the $\mathcal{N} = 4$ SYM theory on S^3.

- **Gross-Witten-Wadia (GWW) type third-order phase transition of the matrix model.**
- **SO(6) R-symmetry of the Yang-Mills theory is unbroken.**

Further development

- **Effect of the fermion**
- **Extension to higher-dimensional system, such as $S^1 \times S^1 \times S^2$.**