

Monte Carlo studies of the 4d supersymmetric reduced model with the Chern-Simons term

Takehiro Azuma

High-energy Accelerator Research Organization (KEK)

JPS Meeting 2004 at Kouchi University, Sep. 28th 2004 15:30-15:45

collaborated with K.N. Anagnostopoulos, K. Nagao and J. Nishimura

Contents

1	Introduction	2
2	Supersymmetric YMCS model	3
3	Result of the numerical studies	4
4	Conclusion	10

1 Introduction

Motivation of the fuzzy-sphere studies:

- Relation between the **non-commutative field theory** and the **superstring theory**.
- Novel regularization scheme alternative to the lattice regularization.
- Prototype of the **curved-space background** in the large- N reduced model.

Matrix models on the homogeneous space

[hep-th/0101102](#), [0103192](#), [0204256](#), [0207115](#), [0209057](#), [0301055](#), [0303120](#), [0307007](#), [0309264](#), [0312241](#), [0403242](#), [0405096](#), [0405277](#)

2 Supersymmetric YMCS model

4d supersymmetric Yang-Mills-Chern-Simons (YMCS) model:

$$S = N \text{tr} \left(-\frac{1}{4} \sum_{\mu, \nu=1}^4 [A_\mu, A_\nu]^2 + \frac{2}{3} i \alpha \sum_{\mu, \nu, \rho=1}^3 \epsilon_{\mu\nu\rho} A_\mu A_\nu A_\rho + \bar{\psi}_\alpha \sum_{\mu=1}^4 (\Gamma_\mu)_{\alpha\beta} [A_\mu, \psi_\beta] \right),$$

A_μ (ψ_α): $N \times N$ traceless hermitian (complex) matrices.

$$\Gamma_1 = i\sigma_1 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \quad \Gamma_2 = i\sigma_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \Gamma_3 = i\sigma_3 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad \Gamma_4 = \sigma_4 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Classical equation of motion:

$$[A_\nu, [A_\nu, A_\mu]] + i\alpha \epsilon_{\mu\nu\rho} [A_\nu, A_\rho] = 0.$$

Fuzzy S^2 classical solution:

$$A_\mu^{(S^2)} = \begin{cases} \alpha L_\mu^{(N)}, & (\text{for } \mu = 1, 2, 3), \\ 0, & (\text{for } \mu = 4), \end{cases}$$

Numerical studies via the hybrid Monte Carlo (HMC) algorithm up to $N = 16$.



3 Result of the numerical studies



Fuzzy-sphere initial condition of the Monte Carlo simulation:

$$A_\mu^{(0)} = A_\mu^{(S^2)}.$$

The observables to study ($\tilde{\alpha} = \alpha\sqrt{N}$):

$$\begin{aligned} \frac{1}{N^2} \langle S \rangle &\simeq -\frac{\tilde{\alpha}^4}{24} \underbrace{-\frac{1}{2}}_{\text{one-loop}}, \\ \langle \frac{1}{N} \text{tr}(F_{\mu\nu}^2) \rangle &\simeq \frac{\tilde{\alpha}^4}{2} \underbrace{+6}_{\text{one-loop}}. \\ \frac{1}{\sqrt{N}} \langle M \rangle &\simeq -\frac{\tilde{\alpha}^3}{6} \underbrace{+0}_{\text{one-loop}}, \\ R &= \langle \sqrt{\frac{1}{N} \sum_{\mu=1}^4 \text{tr} A_\mu^2} \rangle. \end{aligned}$$





Transition point α_{tr} :

$$\alpha_{\text{tr}} \simeq \begin{cases} 1.1 & (N = 4), \\ 0.5 & (N = 8) \Rightarrow \alpha_{\text{tr}} \simeq O(\frac{1}{N}), \\ 0.3 & (N = 16) \end{cases}$$

Behavior of the eigenvalue distribution function $f(x)$ of the Casimir operator

$$Q = A_1^2 + A_2^2 + A_3^2 + A_4^2.$$

- $\alpha < \alpha_{\text{tr}}$

Spikes in the history of $\frac{1}{N} \text{tr } A_\mu^2$.

Power-law tail behavior $f(x) \simeq x^{-2}$.

W. Krauth and M. Staudacher, Phys. Lett. B 453, 253 (1999), [hep-th/9902113].

- $\alpha > \alpha_{\text{tr}}$

No spikes in the history of $\frac{1}{N} \text{tr } A_\mu^2$.

Peaks of $f(x)$ around the classical radius $Q = \frac{N^2-1}{4} \alpha^2$.

Absence of the power-law behavior.




Figure 1: The history of $\frac{1}{N} \text{tr } A_\mu^2$ and the eigenvalue distribution $f(x)$ for $N = 4$.

Argument from one-loop effective action

One-loop perturbation around the fuzzy sphere solution


$$A_\mu = \begin{cases} \beta L_\mu^{(N)}, & (\text{for } \mu = 1, 2, 3), \\ 0, & (\text{for } \mu = 4), \end{cases}$$

One-loop effective action at large N :

$$W = N^2 \left(\frac{\tilde{\beta}^4}{8} - \frac{\tilde{\alpha}\tilde{\beta}^3}{6} \right) - N^2 \log N.$$

Minimum at $\tilde{\beta} = \tilde{\alpha}$.

The fuzzy sphere is always stable for fixed $\tilde{\alpha}(> 0)$ at large N .



4 Conclusion

Four-dimensional supersymmetric Yang-Mills-Chern-Simons (YMCS) model via the hybrid Monte Carlo (HMC) algorithm.

- The transition point $\alpha_{\text{tr}} \simeq O(\frac{1}{N})$.
 - * $\alpha < \alpha_{\text{tr}}$: Power-law behavior of eigenvalue distribution function $f(x) \simeq x^{-2}$.
 - * $\alpha > \alpha_{\text{tr}}$: Stability of the fuzzy sphere.
Absence of the power-law behavior.

Future direction

- Simulation for larger N using more efficient codes.