Perturbative dynamics of fuzzy spheres at large N (hep-th/0410263)

Takehiro Azuma

High-energy Accelerator Research Organization (KEK)

JPS meeting 2005 at Tokyo University of Science

Mar. 27th 2005, 11:30 $\sim$ 11:45

Contents

1 Introduction **2**

2 The model and its classical solution **3**

3 All order calculation from one-loop effective action **6**

4 Conclusion **10**

1 Introduction

Matrix models on a homogeneous space

Motivations of fuzzy manifold studies:

- **Relation between the non-commutative field theory and the superstring.**
- **Novel regularization scheme alternative to lattice regularization.**
- **Prototype of the curved-space background in the large-N reduced models.**

Matrix models on a homogeneous space G/H:

G= (a Lie group), H = (a closed subgroup of G).

$\mathrm{S}^2 = \mathrm{SU}(2)/\mathrm{U}(1)$, $\mathrm{S}^2 \times \mathrm{S}^2$, $\mathrm{S}^4 = \mathrm{SO}(5)/\mathrm{U}(2)$, $\mathrm{CP}^2 = \mathrm{SU}(3)/\mathrm{U}(2)$, $\cdots$.

These fuzzy manifolds are compact, and thus realized by finite matrices.

The Chern-Simons term is added to accommodate the classical solution of the fuzzy manifolds.

2 The model and its classical solution

3d Yang-Mills-Chern-Simons (YMCS) model

⇒ a toy model with fuzzy sphere solutions:

S. Iso, Y. Kimura, K. Tanaka and K. Wakatsuki, hep-th/0101102.

$$S[A] = N\mathrm{tr}\left(-\frac{1}{4}[A_\mu, A_\nu]^2 + \frac{2i\alpha}{3}\epsilon_{\mu\nu\rho}A_\mu A_\nu A_\rho\right).$$

- Defined in the 3-dimensional Euclidean space ($\mu, \nu, \rho = 1, 2, 3$).
- Convergence of the path integral P. Austing and J. F. Wheater, hep-th/0310170.
- Classical equation of motion: $[A_\nu, [A_\mu, A_\nu]] - i\alpha\epsilon_{\mu\nu\rho}[A_\nu, A_\rho] = 0$.
- fuzzy S^2 classical solutions: $A_\mu = X_\mu = \alpha L_\mu$, (where $[L_\mu, L_\nu] = i\epsilon_{\mu\nu\rho}L_\rho$).

  L_μ = ($N \times N$ representation of the SU(2) Lie algebra).

  Casimir operator: $Q = A_1^2 + A_2^2 + A_3^2 = R^2 1_N$.

  R = (radius of the fuzzy sphere) = $\frac{\alpha}{2}\sqrt{N^2 - 1}$.

First-order phase transition

Monte Carlo simulation launched from fuzzy sphere classical solution:

Critical point at $\alpha_{\rm cr} \simeq \frac{2.1}{\sqrt{N}}$.

- **$\alpha < \alpha_{\rm cr}$: Yang-Mills phase**

  Strong quantum effects.

  behavior like the $\alpha = 0$ case.

  T. Hotta, J. Nishimura and A. Tsuchiya, hep-th/9811220,

$$\langle \frac{S}{N^2} \rangle \simeq \mathrm{O}(1), \ \langle \frac{1}{N} \mathrm{tr}\, A_\mu^2 \rangle \simeq \mathrm{O}(1).$$

- **$\alpha > \alpha_{\rm cr}$: fuzzy sphere phase.**

  Fuzzy sphere configuration is stable.

Yang–Mills phase



Fuzzy sphere phase



Phase transition from the one-loop effective action

The effective action Γ is saturated at the one-loop level at large N.

T. Imai, Y. Kitazawa, Y. Takayama and D. Tomino, hep-th/0307007.

Effective action at one-loop around $A_\mu = tX_\mu$ (where $\tilde{\alpha} = \alpha\sqrt{N}$).

$$\frac{\Gamma_{1-\mathrm{loop}}}{N^2} \simeq \tilde{\alpha}^4 \left(\frac{t^4}{8} - \frac{t^3}{6}\right) + \log t.$$

The local minimum disappears at $\tilde{\alpha} < \tilde{\alpha}_{\mathrm{cr}} = (\frac{8}{3})^{\frac{3}{4}} \simeq 2.086\cdots$.

Consistent with the Monte Carlo simulation.



3 All order calculation from one-loop effective action

The free energy W can be obtained by the extremum of the effective action.

Expansion around $A_\mu = \beta L_\mu$: ($\tilde{\beta} = \beta\sqrt{N}$)

$$\lim_{N\to\infty} \frac{1}{N^2}\Gamma(\tilde{\beta}) = \left(\frac{\tilde{\beta}^4}{8} - \frac{1}{6}\tilde{\alpha}\tilde{\beta}^3\right) + \log\tilde{\beta}.$$

Local minimum for $\tilde{\alpha} > \tilde{\alpha}_{\rm cr} = \sqrt[4]{\frac{512}{27}}$:

$$\begin{aligned}
\tilde{\beta} &= f(\tilde{\alpha}) = \frac{\tilde{\alpha}}{4}\left(1 + \sqrt{1+\delta} + \sqrt{2 - \delta + \frac{2}{\sqrt{1+\delta}}}\right) \\
&= \tilde{\alpha}\left(1 - \frac{2}{\tilde{\alpha}^4} - \frac{12}{\tilde{\alpha}^8} - \frac{120}{\tilde{\alpha}^{12}} - \frac{1456}{\tilde{\alpha}^{16}} - \cdots\right), \text{ where} \\
\delta &= 4\tilde{\alpha}^{-\frac{4}{3}}\left[\left(1 + \sqrt{1 - \frac{512}{27\tilde{\alpha}^4}}\right)^{\frac{1}{3}} + \left(1 - \sqrt{1 - \frac{512}{27\tilde{\alpha}^4}}\right)^{\frac{1}{3}}\right].
\end{aligned}$$

Free energy and observables:

$$\begin{aligned}
\lim_{N\to\infty}\frac{1}{N^2}W &= \left(\frac{1}{8}f(\tilde{\alpha})^4 - \frac{1}{6}\tilde{\alpha}f(\tilde{\alpha})^3\right) + \log f(\tilde{\alpha}) \\
&= -\frac{\tilde{\alpha}^4}{24} + \log\tilde{\alpha} - \frac{1}{\tilde{\alpha}^4} - \frac{14}{3\tilde{\alpha}^8} - \frac{110}{3\tilde{\alpha}^{12}} - \frac{364}{\tilde{\alpha}^{16}} - \cdots, \\
\lim_{N\to\infty}\frac{1}{N^2}\langle S\rangle &= \frac{3}{4} - \frac{1}{24}\tilde{\alpha}f(\tilde{\alpha})^3 \\
&= -\frac{\tilde{\alpha}^4}{24} + 1 \underbrace{+\frac{1}{\tilde{\alpha}^4}}_{\text{agrees with two-loop calculation!}} + \frac{28}{3\tilde{\alpha}^8} + \frac{110}{\tilde{\alpha}^{12}} + \frac{1456}{\tilde{\alpha}^{16}} + \cdots.
\end{aligned}$$

All order calculation of generic observables $\mathcal{O}$

Consider the action $S_\epsilon = S + \epsilon\mathcal{O}$.

Corresponding free energy:

$$\begin{aligned} W_\epsilon &= -\log\left(\int d\tilde{A} e^{-(S+\epsilon\mathcal{O})}\right) = -\log\left(\int d\tilde{A} e^{-S}\right) + \epsilon\frac{\int d\tilde{A}\mathcal{O}e^{-S}}{\int d\tilde{A}e^{-S}} + \mathrm{O}(\epsilon^2) \\ &= W + \epsilon\langle\mathcal{O}\rangle + \mathrm{O}(\epsilon^2). \end{aligned}$$

One-loop effective action (take only 1PI diagrams into account)

$$\Gamma_\epsilon(\tilde{\beta}) = \Gamma(\tilde{\beta}) + \epsilon\Gamma_1(\tilde{\beta}) + \mathrm{O}(\epsilon^2).$$

Its saddle point:

$$\frac{\partial}{\partial\tilde{\beta}}\Gamma_\epsilon(\tilde{\beta}) = 0, \quad \Rightarrow \tilde{\beta} = f(\tilde{\alpha}) + \epsilon g(\tilde{\alpha}) + \mathrm{O}(\epsilon^2).$$

Plugging this solution, we obtain the free energy as

$$W_\epsilon = \Gamma_\epsilon(f(\tilde{\alpha}) + \epsilon g(\tilde{\alpha}) + \cdots) = \Gamma(f(\tilde{\alpha})) + \epsilon\left(\Gamma_1(f(\tilde{\alpha})) + g(\tilde{\alpha})\underbrace{(\frac{\partial\Gamma}{\partial\tilde{\beta}})|_{\tilde{\beta}=f(\tilde{\alpha})}}_{=0}\right) + \mathrm{O}(\epsilon^2).$$

We thus obtain $\langle O\rangle = \Gamma_1(f(\tilde{\alpha}))$.

All order calculation of the spacetime content:

$$\lim_{N\to\infty}\frac{1}{N}\langle\frac{1}{N}\mathrm{tr}\,A_\mu^2\rangle=\frac{\tilde{\alpha}^2}{4}\underbrace{-\frac{1}{\tilde{\alpha}^2}}_{\text{one-loop}}.$$

The one-loop effect comes from tadpole diagrams.

$$\frac{1}{N}\langle\frac{1}{N}\mathrm{tr}\,A^2\rangle=\frac{1}{4}f(\tilde{\alpha})^2=\frac{1}{4}\tilde{\alpha}^2-\frac{1}{\tilde{\alpha}^2}-\frac{5}{\tilde{\alpha}^6}-\frac{48}{\tilde{\alpha}^{10}}-\frac{572}{\tilde{\alpha}^{14}}-\cdots.$$

Other observables:

$$\begin{aligned}\lim_{N\to\infty}\frac{1}{\sqrt{N}}\langle M\rangle &= -\frac{1}{6}f(\tilde{\alpha})^3=-\frac{1}{6}\tilde{\alpha}^3+\frac{1}{\tilde{\alpha}}+\frac{4}{\tilde{\alpha}^5}+\frac{112}{3\tilde{\alpha}^9}+\frac{440}{\tilde{\alpha}^{13}}+\cdots,\\ \lim_{N\to\infty}\left\langle\frac{1}{N}\mathrm{tr}\,(F_{\mu\nu})^2\right\rangle &= 3+\frac{1}{2}\tilde{\alpha}f(\tilde{\alpha})^3=\frac{1}{2}\tilde{\alpha}^4-\frac{12}{\tilde{\alpha}^4}-\frac{112}{\tilde{\alpha}^8}-\frac{1320}{\tilde{\alpha}^{12}}-\cdots.\end{aligned}$$


<S>/N^2
<(1/N) tr A^2>/N
<(1/N) tr F^2>
<M>/N^1/2
α̃
N=8
N=16
N=32
classical
one-loop
all order

4 Conclusion

- **In this talk, we have scrutinized the perturbative dynamics of the 3d YMCS model.**
- **We have obtained the all order results for generic observables at large N.**

Future direction

- **Extension of this technique to the 4-dimensional fuzzy manifolds:**

  fuzzy CP^2 (hep-th/0405277), **fuzzy $\mathrm{S}^2 \times \mathrm{S}^2$** (hep-th/0503***).