

Re-analysis of the supernova data to test the effects of large-scale inhomogeneities

Masaru Adachi, Masumi Kasai

from Hirosaki University

JGRG20 in Kyoto University

September 21 - 25

We re-analyzed
the SN gold set data
from
an inhomogeneous
viewpoint,
and...

**found that the SN data
does not necessarily
require **Dark Energy**,
if...**

if we accept the idea
 Ω_m and H_0 may be
different between
here and there,
namely...

**the inhomogeneous
universe!**

Model without Λ



gives even

better fitting

for $z > 0.1$

Fitting the Distant SNe only: No Λ Necessary?

Since the distant SN data are important to determine the value of Ω_m and Ω_Λ , what happens if we use only the distant SNe, say $z > 0.1$, for the fitting? The following figure shows the best fit results. The left figure shows the best fit open model without Λ , and the right figure also shows the best fit flat model with Λ . The χ^2 values shows the open model without Λ gives “*slightly*” better fitting.



the best fit model for $z > 0.1$	χ^2	deg. of freedom	reduced χ^2
flat ($\Omega_m = 0.41, \Omega_\Lambda = 0.59$)	108.80	$91 - 2 = 89$	1.222
open ($\Omega_m = 0.38, \Omega_\Lambda = 0$)	108.40	$91 - 2 = 89$	1.218

**We also investigated
the effects of local
emptiness using
Dyer-Roeder distance.**

**If you are interested in
our results,
please come to our
poster (No.1).**