

No. 38


The power spectrum of
the magnetic fields generated
by the second-order perturbations
during the pre-recombination era

Department of Physics, Kyoto University

Satoshi Maeda

Collaborators : K.Takahashi(Nagoya Univ.), T.Kobayashi(Tokyo Univ.),
T.Shiromizu(Kyoto Univ.)

Previous results



The magnetic fields generated by the second-order perturbations and the Thomson scattering.

➔ Evolution equation of the magnetic fields

$$(a^3 B^i)' = \frac{1 - \beta^3}{1 + \beta} \frac{\sigma_T}{e} a^2 \rho_\gamma^{(0)} \left[2 \frac{H}{\alpha^{(0)}} a^2 \omega^{(2)i} + \varepsilon^{ijk} \frac{R^{(0)} \partial_j \Delta_b^{(I,1)}}{1 + R^{(0)}} \delta v_{(\gamma b)k}^{(I,1)} \right]$$

Vorticity term

Slip term

$$\left(\begin{array}{l} ' : \text{derivative of the conformal time} \\ \beta = m_e / m_p \\ \omega^{i(2)} : \text{a vorticity of the photon} \end{array} \quad \begin{array}{l} \Delta_b : \text{density perturbations of the baryon} \\ \delta v_{(\gamma b)i}^{(I,1)} = \frac{1}{\alpha^{(0)}} \left[H v_i^{(1)} - \frac{1}{4} \frac{\partial_i \delta \rho_\gamma^{(1)}}{\rho_\gamma^{(0)}} \right] \end{array} \right)$$

Results

- The power spectrum of the magnetic fields by slip terms

