

Introduction to Lattice QCD

(格子QCD入門)

Sinya AOKI

Yukawa Institute for Theoretical Physics, Kyoto University




三者共通講義@原子核三者若手夏の学校

2014年8月4日, パノラマランド木島平、長野県、 JAPAN

1. Introduction

格子QCDで何が計算できるか？

Quarks



Hadrons are made of more fundamental objects, named “quarks”.

1973: Kobayashi and Maskawa predicted existences of 6 types(“flavor”) of quarks.



Kobayashi



Maskawa,
7th director of YITP

2008 Nobel prize



charge $2e/3$

charge $-e/3$

QCD (Quantum ChromoDynamics)


QCD: theory for dynamics of quarks

cf. QED (Quantum Electrodynamics)

$$\mathcal{L} = \bar{q}(x) \gamma^\mu \{ \partial_\mu + ig A_\mu(x) \} q(x) + \frac{1}{4} \{ F_{\mu\nu}^a(x) \}^2$$

gluon quark

quark



$$a = 1 \sim 8$$

$$\bar{q} A_\mu q = \bar{q}^A T_{AB}^a A_\mu^a q^B$$


$$A, B = 1, 2, 3 \text{ (color)}$$

quarks-gluon interaction
(electrons-photon in QED)



$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a - gf^{abc} A_\mu^b A_\nu^c$$

gluon field strength

$$(F_{\mu\nu}^a)^2$$



self-interaction
(absent in QED)



g : unique coupling constant in QCD
universal for all flavors

Some Properties of QCD

Asymptotic freedom

forces becomes weaker at shorter distances

Gross



Politzer




Wilczek



2004 Nobel prize


Quark confinement

forces becomes stronger at longer distances




no isolated quark can be observed

structure of nucleon



quark confinement

Difficulties of QCD





“Free” proton = 3 quarks interacting with each others by exchanging a lot of gluons, so that they move coherently.


Clearly, perturbation theory does not work !

Lattice QCD

We need a non-perturbative method.



1-1. Hadron masses



BMW collaboration
 Sciences 322(2008)1224
 2+1 flavor QCD

an agreement between lattice QCD
 and experiments is good.

Baryon

Quarks	Octet($\frac{1}{2}$)	Decouplet($\frac{3}{2}$)
uuu		Δ^{++}
uud	N p	Δ^+
udd	n	Δ^-
ddd		Δ^0
uus	Σ Σ^+	$(\Sigma^*)^+$
uds	Λ Σ^0, Λ^0	Σ^* $(\Sigma^*)^0$
dds	Σ^-	$(\Sigma^*)^-$
uss	Ξ^0	Ξ^* $(\Xi^*)^0$
dss	Ξ^-	$(\Xi^*)^-$
sss		Ω Ω


Meson

Quarks	PesudoScala(0)	Vector(1)
$\bar{u}u - \bar{d}d$	π π^0	ρ ρ^0
$\bar{d}u, \bar{u}d$	$\pi^\pm$	$\rho^\pm$
$\bar{u}u + \bar{d}d$	η	ω
$\bar{s}d, \bar{d}s$	K $K^0, \bar{K}^0$	$(K^*)^0, (\bar{K}^*)^0$
$\bar{s}u, \bar{u}s$	$K^\pm$	$(K^*)^\pm$
$\bar{s}s$	η_s	ϕ

Iso-spin breaking effects

$m_u \neq m_d$ and QED

Borsanyi et al.
arXiv:1406.4088[hep-lat]
1+1+1+1 flavor QCD





	mass splitting [MeV]	QCD [MeV]	QED [MeV]
$\Delta N = n - p$	1.51(16)(23)	2.52(17)(24)	-1.00(07)(14)
$\Delta \Sigma = \Sigma^- - \Sigma^+$	8.09(16)(11)	8.09(16)(11)	0
$\Delta \Xi = \Xi^- - \Xi^0$	6.66(11)(09)	5.53(17)(17)	1.14(16)(09)
$\Delta D = D^\pm - D^0$	4.68(10)(13)	2.54(08)(10)	2.14(11)(07)
$\Delta \Xi_{cc} = \Xi_{cc}^{++} - \Xi_{cc}^+$	2.16(11)(17)	-2.53(11)(06)	4.69(10)(17)
$\Delta_{CG} = \Delta N - \Delta \Sigma + \Delta \Xi$	0.00(11)(06)	-0.00(13)(05)	0.00(06)(02)

1-2. Weak Matrix Elements

Meson decay constants

$$\langle 0 | \bar{d} \gamma_\mu \gamma_5 u | \pi^+(p) \rangle = i p_\mu \underline{f_\pi}, \quad \langle 0 | \bar{s} \gamma_\mu \gamma_5 u | K^+(p) \rangle = i p_\mu \underline{f_K}.$$

axial-vector current



Lattice

$$f_\pi = 130.2 (1.4) \text{ MeV} \quad (N_f = 2+1),$$

$$f_K = 156.3 (0.9) \text{ MeV} \quad (N_f = 2+1),$$

$$f_K = 158.1 (2.5) \text{ MeV} \quad (N_f = 2).$$

Particle Data Group

$$f_\pi^{(PDG)} = 130.41 (0.20) \text{ MeV},$$


$$f_K^{(PDG)} = 156.1 (0.8) \text{ MeV},$$

Kaon B-parameter B_K

$K_0 - \bar{K}_0$ mixing parameter (indirect CP violation)

$$B_K(\mu) = \frac{\langle \bar{K}^0 | Q_R^{\Delta S=2}(\mu) | K^0 \rangle}{\frac{8}{3} f_K^2 m_K^2}.$$

$$Q^{\Delta S=2} = [\bar{s} \gamma_\mu (1 - \gamma_5) d] [\bar{s} \gamma_\mu (1 - \gamma_5) d]$$



$$\hat{B}_K = \left(\frac{\bar{g}(\mu)^2}{4\pi} \right)^{-\gamma_0/(2\beta_0)} \times \left\{ 1 + \frac{\bar{g}(\mu)^2}{(4\pi)^2} \left[\frac{\beta_1 \gamma_0 - \beta_0 \gamma_1}{2\beta_0^2} \right] \right\} B_K(\mu).$$

$$N_f = 2 + 1 : \quad \hat{B}_K = 0.7661(99),$$

$$N_f = 2 + 1 : \quad B_K^{\overline{\text{MS}}}(2 \text{ GeV}) = 0.5596(72).$$

$K \rightarrow (\pi\pi)_{I=2}$ decay amplitude

T. Blum et al., PRL108(2012)141061
T. Blum et al., PRD86(2012)074513

Lattice

$$\begin{aligned}\text{Re}A_2 &= 1.381(46)_{\text{stat}}(258)_{\text{syst}} 10^{-8} \text{ GeV}, \\ \text{Im}A_2 &= -6.54(46)_{\text{stat}}(120)_{\text{syst}} 10^{-13} \text{ GeV}.\end{aligned}$$

Experiment

$$\text{Re}A_2 = 1.479(4) \times 10^{-8} \text{ GeV}$$

K^+ decays

$$a^{-1} = 1.364 \text{ GeV}, m_\pi = 142 \text{ MeV}, m_K = 506 \text{ MeV}$$

$$W_{2\pi} = 486 \text{ MeV}$$

$\Delta I = 1/2$ rule

P. Boyle et al., PRL110(2013)152001

Lattice

$$\frac{\text{Re}A_0}{\text{Re}A_2} = \begin{cases} 9.1(2.1) & \text{for } m_K = 878 \text{ MeV}, m_\pi = 422 \text{ MeV} \\ 12.0(1.7) & \text{for } m_K = 662 \text{ MeV}, m_\pi = 329 \text{ MeV}. \end{cases}$$

Experiment


$$\frac{\text{Re } A_0}{\text{Re } A_2} = 22.45(6)$$

$$a^{-1} = 1.73 \text{ GeV}$$

1-3. Finite temperature QCD

Phase transition at finite T

hadrons (quark confinement)



quark-gluon plasma (deconfinement)

$T \rightarrow \text{large}$


Borsanyi et al.

arXiv:1312.2193[hep-lat]


2+1 flavor QCD

Equation of states of QCD

Pressure



Entropy & energy density



1-4. Hadron Interactions


Ex. Nuclear Force

1949 Nobel prize
(1st in Japan)





Meson Theory (before quarks) 1935 Hideki Yukawa (1st director of YITP)

- Nucleons interact with each other by exchanging virtual particles.
- the interaction range is proportional to the inverse of the virtual particle's mass
 - -> the virtual particles are heavier than electrons but lighter than nucleons
 - -> (π) "meson"



Coulomb potential

$$V(r) = \frac{e^2}{4\pi} \frac{1}{r}$$



Yukawa potential

$$V(r) = \frac{g^2}{4\pi} \frac{e^{-m_\pi r}}{r}$$


Modern nuclear forces after Yukawa

Nuclear Potential

Yukawa(1935)



One-pion exchange



Multi-pions/heavy mesons



Taketani et al.(1951)




distance between nucleons





Repulsive core

Jastrow(1951)

Nuclear forces in terms of quarks ?




Meson Theory



Quark Theory

Much more difficult than masses.

1st lattice result for NN potential




$a=0.137$ fm $L=4.4$ fm $m_\pi \simeq 0.53$ GeV

Ishii-Aoki-Hatsuda

PRL90(2007)0022001

quenched QCD

potential from experiments





prediction from Yukawa theory

2. Lattice QCD

Lattice QCD


define QCD on a discrete space-time (lattice)




$$U_{\mu}(x) = e^{igaA_{\mu}(x)} = 1 + igaA_{\mu}(x) + \frac{\{igaA_{\mu}(x)\}^2}{2!} + \dots \in \text{SU}(3) \quad \text{SU}(3) \text{ matrix}$$

gluon (lives on link)

infinite numbers of gluons (non-perturbative) !



$$U_{\mu}(x)$$



$$U_{-\mu}(x + \hat{\mu}) \equiv U_{\mu}(x)^{\dagger}$$

continuum QCD

lattice QCD

$$\bar{q}(x)\gamma^\mu\{\partial_\mu + igA_\mu(x)\}q(x) \quad \xleftarrow{a \rightarrow 0} \quad \bar{q}(x)\gamma^\mu \frac{U_\mu(x)q(x + \hat{\mu}) - U_{-\mu}(x)q(x - \hat{\mu})}{2a}$$

quarks(covariant derivative)

$$\bar{q}(x)U_\mu(x)q(x + \hat{\mu}) = \text{---} + \text{---} \begin{array}{c} \vdots \\ \vdots \end{array} + \text{---} \begin{array}{c} \diagup \quad \diagdown \\ \diagup \quad \diagdown \end{array} + \text{---} \begin{array}{c} \diagup \quad \diagdown \\ \vdots \quad \diagdown \end{array} + \dots$$

quark interacts with many gluons in a very short distance !

quark action

$$S_F = \sum_{x,\mu} \bar{q}(x)\gamma^\mu \frac{U_\mu(x)q(x + \hat{\mu}) - U_{-\mu}(x)q(x - \hat{\mu})}{2a} + m \sum_x \bar{q}(x)q(x) \quad \text{gauge invariant}$$

gauge transformation

$$q(x) \rightarrow \Omega(x)q(x)$$

$$\bar{q}(x) \rightarrow \bar{q}(x)\Omega(x)^\dagger$$

$$U_\mu(x) \rightarrow \Omega(x)U_\mu(x)\Omega(x + \hat{\mu})^\dagger$$


continuum QCD

$$\frac{g^2}{2} \text{tr } F_{\mu\nu}(x)^2$$

$\longleftarrow$
 $a \rightarrow 0$

lattice QCD

$$\text{tr } U_\mu(x) U_\mu(x + \hat{\mu}) U_\mu(x + \hat{\nu})^\dagger U_\nu(x)^\dagger \quad \text{gluons}$$



gluon action

$$S_G = \frac{1}{g^2} \sum_x \sum_{\mu \neq \nu} \text{tr } U_\mu(x) U_\mu(x + \hat{\mu}) U_\mu(x + \hat{\nu})^\dagger U_\nu(x)^\dagger \quad \text{gauge invariant}$$

Path integral

continuum QCD

$$\begin{aligned}\langle \mathcal{O}(A_\mu, q, \bar{q}) \rangle &= \frac{1}{Z} \int \mathcal{D}A_\mu \mathcal{D}q \mathcal{D}\bar{q} \mathcal{O}(A_\mu, q, \bar{q}) e^{-S_0 - S_{\text{int}}} \\ &= \frac{1}{Z} \int \mathcal{D}A_\mu \mathcal{D}q \mathcal{D}\bar{q} \mathcal{O}(A_\mu, q, \bar{q}) \sum_{n=0}^{\infty} \frac{(-S_{\text{int}})^n}{n!} e^{-S_0}\end{aligned}$$

perturbative expansion

lattice QCD

$$\langle \mathcal{O}(U_\mu, q, \bar{q}) \rangle = \frac{1}{Z} \int \mathcal{D}U_\mu \mathcal{D}q \mathcal{D}\bar{q} \mathcal{O}(U_\mu, q, \bar{q}) e^{-S_F - S_G}$$

calculate without perturbative expansion

important properties

$$\int \mathcal{D}U_\mu(x) U_\mu(x) = 0$$

gluon is random

$$\int \mathcal{D}U_\mu(x) U_\mu(x) U_\mu(x)^\dagger = \mathbf{1}_{3 \times 3}$$

$$\int \mathcal{D}U_\mu(x) \det U_\mu(x) = 1$$

Strong coupling expansion

$$S_G = O\left(\frac{1}{g^2}\right) \rightarrow 0 \quad g^2 \rightarrow \infty \quad \text{strong coupling limit}$$

quark path-integral


quark $\xrightarrow{U_\mu(x)} = 0$ by U integral quark confinement

meson $\xleftrightarrow[U_\mu(x)^\dagger]{U_\mu(x)} \neq 0$

after U integral


baryon $\xrightarrow{U_\mu(x)^3} \neq 0$

meson and baryon can propagate !



in terms of perturbation theory

If $\frac{1}{g^2}$ is small but non-zero



$$= O\left(\frac{1}{g^2}\right)$$

strong coupling expansion

3 quarks can propagate separately but still coherently,
as a free baryon.


Monte-Carlo simulations

After integral over quarks

$$\begin{aligned}\langle \mathcal{O}(q, \bar{q}, U) \rangle &= \int \mathcal{D}q \mathcal{D}\bar{q} \mathcal{D}U \exp[\bar{q} D(U) q + S_G(U)] \mathcal{O}(q, \bar{q}, U) \\ &= \int \mathcal{D}U \underbrace{\det D(U) e^{S_G(U)}}_{\text{probability of } U \equiv P(U)} \hat{\mathcal{O}}(U)\end{aligned}$$

Importance sampling according to $P(U)$ “Monte-Carlo simulations”

calculate complicated QCD processes by computer simulations




uses of super-computers are required.

Yet calculations are not so easy.

Recently hadron masses have been accurately calculated. (free hadrons)

Hadron mass calculations



creation/annihilation of quark-antiquark pair


“effect of $\det D(U)$ ”

set $\det D(U) = 1$: quenched approximation

$$= C_0 e^{-E_0 |x-y|} + C_1 e^{-E_1 |x-y|} + \dots$$

$$E_n = \sqrt{m_n^2 + p^2}$$


extract the smallest value, E_0 , at large $|x - y|$



hadron mass m_0

Meson propagator

Meson propagator




periodic boundary condition in time

pion is lighter than rho.

Nucleon propagator

Nucleon propagator




Nucleon is lighter than its negative-parity state.

Chiral extrapolation

It is difficult to make quark mass as small as the “experimental” value in numerical simulations. Extrapolations from heavier quark masses are usually made.


Pion mass



quark mass

$$2m_q a = \frac{1}{K} - \frac{1}{K_c}$$

Other hadrons




Continuum extrapolation

$a \rightarrow 0$ limit should be taken.

Nucleon mass

m_N vs. a

Plaquette gauge/Wilson quark, Quenched QCD




continuum extrapolation by fit

$$m_N(a) = m_N(0) + C_1 a$$

$$m_N(a) = m_N + C_1 a^2 + C_2 a^2$$

lattice spacing

The state of arts for hadron masses




an agreement between lattice QCD
and experiments is good.

Baryon

Quarks	Octet($\frac{1}{2}$)	Decouplet($\frac{3}{2}$)
uuu		Δ^{++}
uud	N p	Δ^+
udd	n	Δ^-
ddd		Δ^0
uus	Σ Σ^+	$(\Sigma^*)^+$
uds	Λ Σ^0, Λ^0	Σ^* $(\Sigma^*)^0$
dds		$(\Sigma^*)^-$
uss	Ξ Ξ^0	Ξ^* $(\Xi^*)^0$
dss	Ξ^-	$(\Xi^*)^-$
sss		Ω Ω

Meson

Quarks	PesudoScala(0)	Vector(1)
$\bar{u}u - \bar{d}d$	π π^0	ρ ρ^0
$\bar{d}u, \bar{u}d$	$\pi^\pm$	$\rho^\pm$
$\bar{u}u + \bar{d}d$	η	ω
$\bar{s}d, \bar{d}s$	K $K^0, \bar{K}^0$	$(K^*)^0, (\bar{K}^*)^0$
$\bar{s}u, \bar{u}s$	$K^\pm$	$(K^*)^\pm$
$\bar{s}s$	η_s	ϕ




$$a = 0.09 \text{ fm}$$

$$L = 2.9 \text{ fm} \quad m_\pi L = 2.3$$

$$m_\pi^{\text{min.}} = 156 \text{ MeV}$$


Almost on physical quark mass
(no chiral extrapolation)




chiral extrapolation vs. physical point

Chiral extrapolation sometimes becomes non-trivial due to the chiral-log, as shown in the figure.

Hadron interactions from lattice QCD ?



3. Hadron interactions from lattice QCD



--approaches to nuclear physics from lattice QCD--

3 strategies to nuclear physics from lattice QCD


Extreme

calculate **nuclei** directly from lattice QCD

Ab-Initio but (almost) impossible.

difficult to extract “physics” from results
difficult to apply results to other systems

nuclei propagator



3A quark lines

A: atomic number

large number of contractions/very noisy




some reduction (Doi-Endres, CPC 184(2013)117)

binding energy of $A=3,4$ nuclei

${}^3\text{H}$ ($={}^3\text{He}$)

${}^4\text{He}$



PACS-CS, PRD81(2010)111504, PRD86 (2012) 074514.
NPLQCD, PPNP66(2011)1, arXiv:1004.2935.

signals can be obtained, though results scatter.

Standard

calculate NN phase shift from lattice QCD

Ab-Initio for phase shift.

Results can not be directly applied to nuclear physics.

phase shift → nuclear potential → nuclear structure

Lüscher's finite volume method for the phase shift

two particles in the finite box ($V = L^3$)

energy $E = 2\sqrt{\mathbf{k}^2 + m^2}$ → $\mathbf{k} \neq \frac{2\pi}{L}\mathbf{n} \ (\mathbf{n} \in \mathbb{Z}^3)$

due to the interaction between two particles

→ phase shift $\delta_l(k_n)$


Formula (Ex.) $k \cot \delta_0(k) = \frac{2}{\sqrt{\pi}L} \underline{Z_{00}(1; q^2)}$ $k = |\mathbf{k}|$ $q = \frac{kL}{2\pi} \neq \mathbf{Z}$

generalize zeta-function $Z_{00}(s; q^2) = \frac{1}{\sqrt{4\pi}} \sum_{\mathbf{n} \in \mathbb{Z}^3} (\mathbf{n}^2 - q^2)^{-s}$




$\pi^+\pi^-$ scattering (ρ meson width)

ETMC: Feng-Jansen-Renner, PLB684(2010)





$m_\pi = 290$ MeV



$m_\pi = 330$ MeV

Resonance can be treated in this way.

$\delta_1(E_{\text{cm}})$



2-flavor anisotropic clover fermion

$a_s \sim 0.12$ fm

$m_\pi \sim 400$ MeV

Alternative calculate nuclear potential from lattice QCD strategy in this lecture


Ab-Initio for potential.

“Physics” is clear

nuclear potential → nuclear structure

Difficulties

A. Interactions are much more difficult than masses.



more complicated diagrams,
larger volume,
more Monte-Carlo sampling, etc.

B. Definition of potential in quantum theories ?

classical $V(x)$ → quantum $V(x)$ potential is an input



no classical NN potentials QCD $V_{NN}(x)$? output from QCD

Potentials in QCD ?


What are “potentials” in quantum field theories such as QCD ?

“Potentials” themselves can NOT be directly measured. cf. running coupling in QCD
scheme dependent, Unitary transformation

experimental data of scattering phase shifts




potentials, but not unique



useful to “understand” physics


cf. asymptotic freedom

“Potentials” are useful tools to extract observables such as scattering phase shift.



One may adopt a convenient definition of potentials as long as they reproduce correct physics of QCD.

3-1. Our strategy



Our strategy in lattice QCD

Full details: Aoki, Hatsuda & Ishii, PTP123(2010)89

Consider “elastic scattering”

$$NN \rightarrow NN \quad \cancel{NN \rightarrow NN + \text{others}} \quad (\cancel{NN \rightarrow NN + \pi, NN + \bar{N}N, \dots})$$

energy $W_k = 2\sqrt{\mathbf{k}^2 + m_N^2} < W_{\text{th}} = 2m_N + m_\pi$ Elastic threshold

Quantum Field Theoretical consideration

Unitarity constrains S-matrix below inelastic threshold as

$$S = e^{2i\delta}$$

Ex. Scalar particles

$$\delta(k) = \begin{pmatrix} \delta_0(k) & & & \\ & \delta_1(k) & & \\ & & \delta_2(k) & \\ & & & \dots \end{pmatrix}$$

Step 1


define (Equal-time) Nambu-Bethe-Salpeter (NBS) Wave function

Spin model: Balog et al., 1999/2001


$$\varphi_{\mathbf{k}}(\mathbf{r}) = \langle 0 | \underline{N(\mathbf{x} + \mathbf{r}, 0)} N(\mathbf{x}, 0) | \underline{NN, W_k} \rangle$$

QCD eigen-state

$N(x) = \varepsilon_{abc} q^a(x) q^b(x) q^c(x)$: local operator “scheme”




Asymptotic behavior of NBS wave function



$r = |\mathbf{r}| \rightarrow \infty$
partial wave

$$\varphi_{\mathbf{k}}^l \rightarrow A_l \frac{\sin(kr - l\pi/2 + \delta_l(k))}{kr}$$


NBS wave function



scattering wave function in quantum mechanics

cf. Luescher’s finite volume method

allowed k at L



$\delta_l(k_n)$

Step 2

define non-local but energy-independent “potential” as

$$\mu = m_N/2$$

reduced mass

$$[\epsilon_k - H_0] \varphi_{\mathbf{k}}(\mathbf{x}) = \int d^3y \underbrace{U(\mathbf{x}, \mathbf{y})}_{\text{non-local potential}} \varphi_{\mathbf{k}}(\mathbf{y})$$
$$\epsilon_k = \frac{\mathbf{k}^2}{2\mu} \quad H_0 = \frac{-\nabla^2}{2\mu}$$

(Trivial) proof of “existence”

We can construct a non-local but **energy-independent** potential easily as

$$U(\mathbf{x}, \mathbf{y}) = \sum_{\mathbf{k}, \mathbf{k}'}^{W_k, W_{k'} \leq W_{\text{th}}} [\epsilon_k - H_0] \varphi_{\mathbf{k}}(\mathbf{x}) \eta_{\mathbf{k}, \mathbf{k}'}^{-1} \varphi_{\mathbf{k}'}^\dagger(\mathbf{y}) \quad \eta_{\mathbf{k}, \mathbf{k}'}^{-1}: \text{inverse of } \eta_{\mathbf{k}, \mathbf{k}'} = (\varphi_{\mathbf{k}}, \varphi_{\mathbf{k}'})$$

inner product

For $\forall W_{\mathbf{p}} < W_{\text{th}}$

$$\int d^3y U(\mathbf{x}, \mathbf{y}) \phi_{\mathbf{p}}(\mathbf{y}) = \sum_{\mathbf{k}, \mathbf{k}'} [\epsilon_k - H_0] \varphi_{\mathbf{k}}(x) \eta_{\mathbf{k}, \mathbf{k}'}^{-1} \eta_{\mathbf{k}', \mathbf{p}} = [\epsilon_p - H_0] \varphi_{\mathbf{p}}(x)$$

Remark

Non-relativistic approximation is **NOT** used. We just take the specific (equal-time) frame.

Step 3 expand the non-local potential in terms of derivative as

$$U(\mathbf{x}, \mathbf{y}) = V(\mathbf{x}, \nabla) \delta^3(\mathbf{x} - \mathbf{y})$$

$$V(\mathbf{x}, \nabla) = \underbrace{V_0(r)}_{\text{LO}} + \underbrace{V_\sigma(r)(\sigma_1 \cdot \sigma_2)}_{\text{LO}} + \underbrace{V_T(r)S_{12}}_{\text{LO}} + \underbrace{V_{\text{LS}}(r)\mathbf{L} \cdot \mathbf{S}}_{\text{NLO}} + \underbrace{O(\nabla^2)}_{\text{NNLO}}$$


tensor operator $S_{12} = \frac{3}{r^2}(\sigma_1 \cdot \mathbf{x})(\sigma_2 \cdot \mathbf{x}) - \overset{\text{spins}}{(\sigma_1 \cdot \sigma_2)}$

This expansion is a part of our “scheme” for potentials.

Step 4 extract the local potential at LO as

$$V_{\text{LO}}(\mathbf{x}) = \frac{[\epsilon_k - H_0]\varphi_{\mathbf{k}}(\mathbf{x})}{\varphi_{\mathbf{k}}(\mathbf{x})}$$

Step 5 solve the Schroedinger Eq. in the infinite volume with this potential.



phase shifts and binding energy below inelastic threshold

We can check a size of errors of the LO in the expansion. (See later).

3-2. Nuclear Potential


Extraction of NBS wave function

Standard method

NBS wave function

$$\varphi_{\mathbf{k}}(\mathbf{r}) = \langle 0 | N(\mathbf{x} + \mathbf{r}, 0) N(\mathbf{x}, 0) | NN, W_k \rangle \quad \Rightarrow \quad [\epsilon_k - H_0] \varphi_{\mathbf{k}}(\mathbf{x}) = \int d^3 y U(\mathbf{x}, \mathbf{y}) \varphi_{\mathbf{k}}(\mathbf{y})$$

Potential



4-pt Correlation function

source for NN

$$F(\mathbf{r}, t - t_0) = \langle 0 | T \{ N(\mathbf{x} + \mathbf{r}, t) N(\mathbf{x}, t) \} \underline{\overline{\mathcal{J}}(t_0)} | 0 \rangle$$

complete set for NN

$$\begin{aligned} F(\mathbf{r}, t - t_0) &= \langle 0 | T \{ N(\mathbf{x} + \mathbf{r}, t) N(\mathbf{x}, t) \} \sum_{n, s_1, s_2} \underline{|2N, W_n, s_1, s_2\rangle \langle 2N, W_n, s_1, s_2| \overline{\mathcal{J}}(t_0)} | 0 \rangle + \dots \\ &= \sum_{n, s_1, s_2} A_{n, s_1, s_2} \varphi^{W_n}(\mathbf{r}) e^{-W_n(t-t_0)}, \quad A_{n, s_1, s_2} = \langle 2N, W_n, s_1, s_2 | \overline{\mathcal{J}}(0) | 0 \rangle. \end{aligned}$$

ground state saturation at large t


$$\lim_{(t-t_0) \rightarrow \infty} F(\mathbf{r}, t - t_0) = \underline{A_0 \varphi^{W_0}(\mathbf{r})} e^{-W_0(t-t_0)} + O(e^{-W_{n \neq 0}(t-t_0)})$$

NBS wave function

This is a standard method in lattice QCD and was employed for our first calculation.

normalized 4-pt function

$$R(\mathbf{r}, t) \equiv F(\mathbf{r}, t) / (e^{-m_N t})^2 = \sum_n A_n \varphi^{W_n}(\mathbf{r}) e^{-\Delta W_n t}$$



potential


$$\Delta W_n = W_n - 2m_N = \frac{\mathbf{k}_n^2}{m_N} - \frac{(\Delta W_n)^2}{4m_N}$$

$$-\frac{\partial}{\partial t} R(\mathbf{r}, t) = \left\{ H_0 + U - \frac{1}{4m_N} \frac{\partial^2}{\partial t^2} \right\} R(\mathbf{r}, t)$$

Leading Order

$$\left\{ \underbrace{-H_0}_{1st} - \underbrace{\frac{\partial}{\partial t}}_{2nd} + \underbrace{\frac{1}{4m_N} \frac{\partial^2}{\partial t^2}}_{3rd} \right\} R(\mathbf{r}, t) = \int d^3 r' U(\mathbf{r}, \mathbf{r}') R(\mathbf{r}', t) = \underbrace{V_C(\mathbf{r})}_{total} R(\mathbf{r}, t) + \dots$$

3rd term(relativistic correction)
is negligible.




Ground state saturation is no more required !
(advantage over finite volume method.)


NN potential

2+1 flavor QCD, spin-singlet potential (PLB712(2012)437)

$a=0.09\text{fm}$, $L=2.9\text{fm}$ $m_\pi \simeq 700\text{ MeV}$



phenomenological potential




Qualitative features of NN potential are reproduced !

- (1) attractions at medium and long distances
- (2) repulsion at short distance(repulsive core)


1st paper(quenched QCD): Ishii-Aoki-Hatsuda, PRL90(2007)0022001

selected as one of 21 papers in **Nature Research Highlights 2007**. (One from Physics, Two from Japan, the other is on “iPS” by Sinya Yamanaka et al.)

NN potential



phase shift



It has a reasonable shape. The strength is weaker due to the heavier quark mass.

Need calculations at physical quark mass on K-computer.



Convergence of velocity expansion: estimate 1

If the higher order terms are large, LO potentials determined from NBS wave functions at **different energy** become different.(cf. LOC of ChPT).

Numerical check in quenched QCD

$$m_{\pi} \simeq 0.53 \text{ GeV}$$

K. Murano, N. Ishii, S. Aoki, T. Hatsuda




$$a=0.137\text{fm}, L=4.0 \text{ fm}$$

PTP 125 (2011)1225.



● PBC ($E \sim 0 \text{ MeV}$)

● APBC ($E \sim 46 \text{ MeV}$)


NBS wave functions




potentials



$V_c(r; {}^1S_0)$: PBC v.s. APBC $t=9$ ($x=\pm 5$ or $y=\pm 5$ or $z=\pm 5$)



phase shifts from potentials



Higher order terms turn out to be very small at low energy in our scheme.

Need to be checked at lighter pion mass in 2+1 flavor QCD.


Note: convergence of the velocity expansion can be checked within this method.

(in contrast to convergence of ChPT, convergence of perturbative QCD)

Convergence of velocity expansion: estimate 2

Kurth, Ishii, Doi, Aoki & Hatsuda, JHEP 1312(2013)015

Potential vs Luescher (l=2 pi-pi scattering. Quenched QCD)



both methods
agree very well.

This establishes a validity of the potential method and shows a good convergence of the velocity expansion.

More structure at LO

Tensor potential

$$(H_0 + V_C(r) + V_T(r)S_{12})\psi(\mathbf{r}; 1^+) = E\psi(\mathbf{r}; 1^+)$$


J=1, S=1

mixing between 3S_1 and 3D_1 through the tensor force

3S_1 3D_1

$$\psi(\mathbf{r}; 1^+) = \mathcal{P}\psi(\mathbf{r}; 1^+) + \mathcal{Q}\psi(\mathbf{r}; 1^+)$$

“projection” to L=0“projection” to L=2





$$\begin{aligned} H_0[\mathcal{P}\psi](\mathbf{r}) + V_C(r)[\mathcal{P}\psi](\mathbf{r}) + V_T(r)[\mathcal{P}S_{12}\psi](\mathbf{r}) &= E[\mathcal{P}\psi](\mathbf{r}) \\ H_0[\mathcal{Q}\psi](\mathbf{r}) + V_C(r)[\mathcal{Q}\psi](\mathbf{r}) + V_T(r)[\mathcal{Q}S_{12}\psi](\mathbf{r}) &= E[\mathcal{Q}\psi](\mathbf{r}) \end{aligned}$$

Wave functions

Quenched


Aoki, Hatsuda, Ishii, PTP 123 (2010)89



divided by $Y_{20}(\theta, \phi)$


Potentials

full QCD



$a \simeq 0.091$ fm $L \simeq 2.9$ fm

quenched QCD




$a \simeq 0.137$ fm $L \simeq 4.4$ fm

- no repulsive core in the tensor potential.
- the tensor potential is enhanced in full QCD


Potentials

full QCD




$a \simeq 0.091$ fm $L \simeq 2.9$ fm

quenched QCD



$a \simeq 0.137$ fm $L \simeq 4.4$ fm




- no repulsive core in the tensor potential.
- the tensor potential is enhanced in full QCD



from
R.Machleidt,
Adv.Nucl.Phys.**19**

Fig. 3.7. The contributions from π and ρ (dashed) to the $T = 0$ tensor potential. The solid line is the full potential. The dash-dot lines are obtained when the cutoff is omitted.

Quark mass dependence (full QCD)



- the tensor potential increases as the pion mass decreases.
 - manifestation of one-pion-exchange ?
- both repulsive core and attractive pocket are also grow as the pion mass decreases.

Potentials for the negative parity sector

$$V_{NN}^{(I)}(\vec{r}, \vec{\nabla}) = \underbrace{V_0^{(I)}(r) + V_\sigma^{(I)}(r) \cdot (\vec{\sigma}_1 \cdot \vec{\sigma}_2)}_{\text{LO}} + \underbrace{V_T^{(I)}(r) \cdot S_{12}}_{\text{LO}} + \underbrace{V_{LS}^{(I)}(r) \cdot \vec{L} \cdot \vec{S}}_{\text{NLO}} + O(\nabla^2)$$

$$V_C(r) \equiv V_0(r) + V_\sigma(r) \cdot (\vec{\sigma}_1 \cdot \vec{\sigma}_2)$$

$$= \begin{cases} V_0(r) - 3V_\sigma(r) & \text{for } S=0 \\ V_0(r) + V_\sigma(r) & \text{for } S=1 \end{cases}$$

S=0, P=+ (l=1)	S=1, P=+ (l=0)	S=0, P=- (l=0)	S=1, P=- (l=1)
$V_C(r)$	$V_C(r), V_T(r), V_{LS}(r)$	$V_C(r)$	$V_C(r), V_T(r), V_{LS}(r)$




$^{2S+1}L_J$

- S=1 channel: $^3P_0, ^3P_1, ^3P_2 - ^3F_2$
 - Central & tensor forces in LO
 - Spin-orbit force in NLO

Potentials


Murano et al. (HAL QCD), arXiv:1305.2293[hep-lat]

$$a = 0.16 \text{ fm}, L = 2.5 \text{ fm}, m_\pi = 1100 \text{ MeV}$$




tensor potential is very weak.


3-3. Hyperon Interactions





Origin of the repulsive core ?

quarks are “fermion”  two can not occupy the same position. (“Pauli principle”)


they have 3 colors(red,blue,green), 2 spin($\uparrow \downarrow$), 2 flavors(up,down)

 6 quark can occupy the same position




but allowed color combinations are limited + interaction among quarks

?


 repulsive core ?

What happen if strange quarks are added ?

$\Lambda(uds) - \Lambda(uds)$ interaction





all color combinations are allowed




no repulsive core ?

Octet Baryon interactions

$$\begin{array}{|c|} \hline 8 \\ \hline \end{array} \otimes \begin{array}{|c|} \hline 8 \\ \hline \end{array} = \begin{array}{|c|c|c|} \hline 27 \\ \hline \end{array} \oplus \begin{array}{|c|c|c|} \hline 10^* \\ \hline \end{array} \oplus \begin{array}{|c|} \hline 1 \\ \hline \end{array} \oplus \begin{array}{|c|} \hline 8 \\ \hline \end{array} \oplus \begin{array}{|c|c|c|} \hline 10 \\ \hline \end{array} \oplus \begin{array}{|c|} \hline 8 \\ \hline \end{array}$$



- phase shift available for YN and YY scattering are limited
- plenty of hyper-nucleus data will be soon available at J-PARC




- prediction from lattice QCD
- difference between NN and YN ?

Baryon Potentials in the flavor SU(3) limit

$$m_u = m_d = m_s$$

1. First setup to predict YN, YY interactions not accessible in exp.
2. Origin of the repulsive core (universal or not)



$$8 \times 8 = \underbrace{27 + 8s + 1}_{\text{Symmetric}} + \underbrace{10^* + 10 + 8a}_{\text{Anti-symmetric}}$$

6 independent potentials in flavor-basis

$$\begin{array}{ll} V^{(27)}(r), & V^{(8s)}(r), & V^{(1)}(r) & \longleftarrow & {}^1S_0 \\ V^{(10^*)}(r), & V^{(10)}(r), & V^{(8a)}(r) & \longleftarrow & {}^3S_1 \end{array}$$

3-flavor QCD

$$a=0.12 \text{ fm}$$


Inoue et al. (HAL QCD Coll.), PTP124(2010)591

$$L=2 \text{ fm}$$


Inoue et al. (HAL QCD Coll.), NPA881(2012)28

$$L=2-4 \text{ fm}$$


$L \simeq 4$ fm, $m_\pi \simeq 470$ MeV




same as NN




8s: strong repulsive core. repulsion only.




1: attractive instead of repulsive
core ! attraction only . H-dibaryon.



same as NN



10: strong repulsive core. weak attraction.




8a: weak repulsive core.
strong attraction.

Flavor dependences of BB interactions become manifest in SU(3) limit !

H-dibaryon:


a possible six quark state(uuddss)
predicted by the model but not observed yet.



<http://physics.aps.org/synopsis-for/10.1103/PhysRevLett.106.162001>

Binding baryons on the lattice

April 26, 2011




H-dibaryon in the flavor SU(3) limit

$a=0.12$ fm

Inoue et al. (HAL QCD Coll.), PRL106(2011)162002


Attractive potential
in the flavor singlet channel




possibility of a bound state (H-dibaryon)

$$\Lambda\Lambda - N\Xi - \Sigma\Sigma$$

volume dependence



pion mass dependence




$L=3$ fm is enough for the potential.

lighter the pion mass, stronger the attraction



fit potentials at $L=4$ fm by

$$V(r) = a_1 e^{-a_2 r^2} + a_3 \left(1 - e^{-a_4 r^2}\right)^2 \left(\frac{e^{-a_5 r}}{r}\right)^2$$

Solve Schroedinger equation
in the infinite volume



One bound state (H-dibaryon) exists.




An H-dibaryon exists in the flavor SU(3) limit.

Binding energy = 25-50 MeV at this range of quark mass.


A mild quark mass dependence.

Real world ?

Deuteron



H-dibayon




3-4. Extensions

H-dibaryon with the flavor SU(3) breaking


SU(3) limit

$\Lambda\Lambda - N\Xi - \Sigma\Sigma$



25-50 MeV

H



Real world

$$m_u = m_d \neq m_s$$

$\Sigma\Sigma$



2386 MeV

$N\Xi$



129 MeV

2257 MeV

H ?




25 MeV

$\Lambda\Lambda$



2232 MeV

H ?



S=-2 “Inelastic” scattering

$$m_N = 939 \text{ MeV}, m_\Lambda = 1116 \text{ MeV}, m_\Sigma = 1193 \text{ MeV}, m_\Xi = 1318 \text{ MeV}$$

S=-2 System(I=0)

$$M_{\Lambda\Lambda} = 2232 \text{ MeV} < M_{N\Xi} = 2257 \text{ MeV} < M_{\Sigma\Sigma} = 2386 \text{ MeV}$$

The eigen-state of QCD in the finite box is a mixture of them:

$$|S = -2, I = 0, E\rangle_L = c_1(L)|\Lambda\Lambda, E\rangle + c_2(L)|\Xi N, E\rangle + c_3(L)|\Sigma\Sigma, E\rangle$$

$$E = 2\sqrt{m_\Lambda^2 + \mathbf{p}_1^2} = \sqrt{m_\Xi^2 + \mathbf{p}_2^2} + \sqrt{m_N^2 + \mathbf{p}_2^2} = 2\sqrt{m_\Sigma^2 + \mathbf{p}_3^2}$$

In this situation, we can not directly extract the scattering phase shift in lattice QCD.

Extended method

Let us consider 2-channel problem for simplicity.

NBS wave functions for 2 channels at 2 values of energy:

$$\begin{aligned}\Psi_{\alpha}^{\Lambda\Lambda}(\mathbf{x}) &= \langle 0 | \Lambda(\mathbf{x}) \Lambda(\mathbf{0}) | E_{\alpha} \rangle \\ \Psi_{\alpha}^{\Xi N}(\mathbf{x}) &= \langle 0 | \Xi(\mathbf{x}) N(\mathbf{0}) | E_{\alpha} \rangle\end{aligned}$$

$$\alpha = 1, 2$$

They satisfy

$$(\nabla^2 + \mathbf{p}_{\alpha}^2) \Psi_{\alpha}^{\Lambda\Lambda}(\mathbf{x}) = 0$$

$$(\nabla^2 + \mathbf{q}_{\alpha}^2) \Psi_{\alpha}^{\Xi N}(\mathbf{x}) = 0$$

$$|\mathbf{x}| \rightarrow \infty$$

We define the “potential” from the **coupled channel** Schroedinger equation:


$$\left(\frac{\nabla^2}{2\mu_{\Lambda\Lambda}} + \frac{\mathbf{p}_\alpha^2}{2\mu_{\Lambda\Lambda}} \right) \Psi_\alpha^{\Lambda\Lambda}(\mathbf{x}) = \underbrace{V^{\Lambda\Lambda \leftarrow \Lambda\Lambda}(\mathbf{x})}_{\text{diagonal}} \Psi_\alpha^{\Lambda\Lambda}(\mathbf{x}) + \underbrace{V^{\Lambda\Lambda \leftarrow \Xi N}(\mathbf{x})}_{\text{off-diagonal}} \Psi_\alpha^{\Xi N}(\mathbf{x})$$

$$\left(\frac{\nabla^2}{2\mu_{\Xi N}} + \frac{\mathbf{q}_\alpha^2}{2\mu_{\Xi N}} \right) \Psi_\alpha^{\Xi N}(\mathbf{x}) = \underbrace{V^{\Xi N \leftarrow \Lambda\Lambda}(\mathbf{x})}_{\text{off-diagonal}} \Psi_\alpha^{\Lambda\Lambda}(\mathbf{x}) + \underbrace{V^{\Xi N \leftarrow \Xi N}(\mathbf{x})}_{\text{diagonal}} \Psi_\alpha^{\Xi N}(\mathbf{x})$$

μ : reduced mass

$$\begin{pmatrix} (E_1 - H_0^X) \Psi_1^X(\mathbf{x}) \\ (E_2 - H_0^X) \Psi_2^X(\mathbf{x}) \end{pmatrix} = \begin{pmatrix} \Psi_1^X(\mathbf{x}) & \Psi_1^Y(\mathbf{x}) \\ \Psi_2^X(\mathbf{x}) & \Psi_2^Y(\mathbf{x}) \end{pmatrix} \begin{pmatrix} V^{X \leftarrow X}(\mathbf{x}) \\ V^{X \leftarrow Y}(\mathbf{x}) \end{pmatrix}$$

$$E_\alpha = \frac{\mathbf{p}_\alpha^2}{2\mu_{\Lambda\Lambda}}, \quad \frac{\mathbf{q}_\alpha^2}{2\mu_{\Xi N}} \quad X \neq Y \quad X, Y = \Lambda\Lambda \text{ or } \Xi N$$



$$\begin{pmatrix} V^{X \leftarrow X}(\mathbf{x}) \\ V^{X \leftarrow Y}(\mathbf{x}) \end{pmatrix} = \begin{pmatrix} \Psi_1^X(\mathbf{x}) & \Psi_1^Y(\mathbf{x}) \\ \Psi_2^X(\mathbf{x}) & \Psi_2^Y(\mathbf{x}) \end{pmatrix}^{-1} \begin{pmatrix} (E_1 - H_0^X) \Psi_1^X(\mathbf{x}) \\ (E_2 - H_0^X) \Psi_2^X(\mathbf{x}) \end{pmatrix}$$

Using the coupled channel potentials:

$$\begin{pmatrix} V^{\Lambda\Lambda \leftarrow \Lambda\Lambda}(\mathbf{x}) & V^{\Xi N \leftarrow \Lambda\Lambda}(\mathbf{x}) \\ V^{\Lambda\Lambda \leftarrow \Xi N}(\mathbf{x}) & V^{\Xi N \leftarrow \Xi N}(\mathbf{x}) \end{pmatrix}$$

we solve the coupled channel Schroedinger equation in **the infinite volume** with **an appropriate boundary condition**.

For example, we take the incoming $\Lambda\Lambda$ state by hand.

In this way, we can avoid the mixture of several “in”-states.

$$|S = -2, I = 0, E\rangle_L = c_1(L)|\Lambda\Lambda, E\rangle + c_2(L)|\Xi N, E\rangle + c_3(L)|\Sigma\Sigma, E\rangle$$

Lattice is a tool to extract the interaction kernel (“T-matrix” or “potential”).

Preliminary results from HAL QCD Collaboration


Sasaki for HAL QCD Collaboration

Gauge ensembles

In unit of MeV	<i>Esb 1</i>	<i>Esb 2</i>	<i>Esb 3</i>
π	701 ± 1	570 ± 2	411 ± 2
K	789 ± 1	713 ± 2	635 ± 2
m_π / m_K	0.89	0.80	0.65
N	1585 ± 5	1411 ± 12	1215 ± 12
Λ	1644 ± 5	1504 ± 10	1351 ± 8
Σ	1660 ± 4	1531 ± 11	1400 ± 10
Ξ	1710 ± 5	1610 ± 9	1503 ± 7

u,d quark masses lighter




thresholds



SU(3) breaking effects becomes larger




coupled channel 3x3 potentials

Diagonal elements



Off-diagonal elements


All channels have repulsive core



$\Lambda\Lambda$ and $N\Xi$ phase shift


Preliminary !

Esb1 : $m\pi = 701$ MeV




Bound H-dibaryon
coupled to $N\Xi$

Esb2 : $m\pi = 570$ MeV



H as $\Lambda\Lambda$ resonance
H as bound $N\Xi$

Esb3 : $m\pi = 411$ MeV



H as $\Lambda\Lambda$ resonance
H as bound $N\Xi$

This suggests that H-dibaryon becomes **resonance** at physical point.
Below or above $N\Xi$? Need simulation at physical point.


Physically, it is essential that H-dibaryon is a bound state in the flavor SU(3) limit.

4. Summary


- **Lattice QCD** is a very powerful method to investigate dynamics of quarks
- not only hadron masses but also hadron interactions can be investigated from the 1st principle
- **the potential (HALQCD) method** is new but very useful to investigate not only the nuclear force but also general baryonic interactions in (lattice) QCD.
- the method can be easily applied also to meson-baryon and meson-meson interactions.

Our strategy



Potentials from
lattice QCD




Nuclear Physics
with these potentials



Neutron stars
Supernova explosion



HAL QCD Collaboration



Sinya Aoki (Kyoto U.)

Bruno Charron* (U. Tokyo)

Takumi Doi (Riken)

Faisal Etminan* (Kyoto U.)

Tetsuo Hatsuda (Riken)

Yoichi Ikeda (Riken)

Takashi Inoue (Nihon U.)

Noriyoshi Ishii (U. Tsukuba)

Keiko Murano (Kyoto U.)

Hidekatsu Nemura (U. Tsukuba)



Kenji Sasaki (U. Tsukuba)

Masanori Yamada* (U. Tsukuba)

*PhD Students


Challenge: Three nucleon force (TNF)

Maximum mass of neutron stars



$(\rho_{\max} \sim 6\rho_0)$

sustains neutron stars
against gravitational collapse






TNF from lattice QCD

Doi et al. (HAL QCD), PTP 127 (2012) 723

(1,2) pair $^1S_0, ^3S_1, ^3D_1$ S-wave only

Triton($I = 1/2, J^P = 1/2^+$)

Linear setup



scalar/isoscalar TNF is observed at short distance.

further study is needed to confirm this result.