



Hierarchy in the cosmic structures

Hubble Ultra Deep Field
@ $t=0.1\text{Gyr}$


Cosmic Microwave Background
@ $t=0.4\text{Myr}$



Simulated halo
@ $t=10\text{Gyr}$



Solar system
@ $t=13.7\text{Gyr}$




Yasushi Suto

Department of Physics, The University of Tokyo

Kyoto University GCOE symposium "Links among Hierarchies"

16:20-16:50 February 13, 2012




From our local universe (SDSS) to Cosmic Microwave Background (WMAP)




Hierarchical structures in the universe

**Cosmic
Microwave
Background**



10^{26}m

**Large-scale
structure**



10^{25}m

**Galaxy
cluster**



10^{12}m

10^{23}m

Solar system




Earth



10^7m


Temperature map of Cosmic Microwave Background





$$\Delta T(\theta, \varphi) = \sum_{\ell, m} a_{\ell m} Y_{\ell m}(\theta, \varphi)$$

$$C_{\ell} = \langle |a_{\ell m}|^2 \rangle$$

**Amazing agreement
between (linear)
theory and
observations
⇒ Precision
measurement of
cosmological
parameters
WMAP 5yr
Komatsu et al.
(2008)**



Baryon Acoustic Oscillation imprinted in galaxy distribution





**4,700 Luminous Red
Galaxies @ $z \sim 0.3$ over
3800 deg^2 from Sloan
Digital Sky Survey**

Eisenstein et al. (2005)

Baryon Acoustic Oscillation in galaxy power spectrum: probe of dark energy


- **Sub-percent level accuracy is achieved** with improved perturbation theory of gravitational evolution of dark matter (Taruya et al. 2009,2010)




Formation of galaxies and clusters from dark matter halos

N-body simulations of dark matter halos in Λ CDM

z=4.2




z=1.0




Jing & Suto (2000)

z=2.1




present epoch
(z=0)

$4 \times 10^{14} M_{\text{sun}}$




Apparent diversities of dark matter halos with different mass and environments


Galaxy scales
 $\sim 5 \times 10^{12} M_{\text{sun}}$



Group scales
 $\sim 5 \times 10^{13} M_{\text{sun}}$




Cluster scales
 $\sim 3 \times 10^{14} M_{\text{sun}}$



Jing & Suto (2000)

Universal density profile of dark matter halos: NFW profile




- Regularity behind diversity
- Dark halos exhibit an unexpected scaling law in the nonlinear gravitational evolution process

$$\rho(r) = \frac{\delta_c \rho_{crit}}{(r/r_s)(1+r/r_s)^2}$$


Navarro, Frenk & White (1997)



A pale blue dot ? Not really



Simulated photometric light-curves of Earth




- Adopted Earth data in March
- Spin inclination = 0 (vernal equinox) **Fujii et al. (2010)**
- cloudless

Color-Changing Planets Could Hold Clues to Alien Life (May 13, 2010 in SPACE.com)

- A new way of comparing the color and intensity changes of light reflected off of Earth's surface to the flickers from exoplanets may help reveal the presence of oceans, continents and – possibly – life on alien worlds.
- By comparing the changes in observed hues of an alien planet as it rotates to this distinct Earthly color palette, "we can infer the surface composition of the [exo]planet," said Yuka Fujii, a doctoral student at the University of Tokyo and lead author of a paper published in the May 4 issue of the *Astrophysical Journal*.

<http://www.space.com/scienceastronomy/color-changing-planets-alien-life-100513.html>

Idealized cloudless earth



Fujii et al. (2010,2011)

Input data

- 5 light-curves using anisotropic scattering (BRDF) model
- 2 week observation of a cloudless Earth at 10 pc away

Inversion assumptions

- Ocean, soil, vegetation and snow only (with atmosphere)
- Isotropic scattering assumed

Results

- Estimated areas (symbols) vs Surface classification data (dashed line)
- Reasonably well reproduced.
- Can identify vegetation !

Useful to identify a second earth ?

From deterministic to stochastic

- **Regularities and diversities in astrophysics**
- **Deterministic on large scales and stochastic on small scales**
 - **CMB**: perfect match with linear theory predictions
 - **LSS**: perturbation theory works well
 - **Galaxy clusters**: nonlinear gravitational evolution leads to a class of regularity in density profiles
 - **Galaxies**: huge varieties, roughly classified in the Hubble sequence
 - **Planetary systems**: until 1995, we did not even know if planets outside the Solar system exist at all
 - **Another Earth**: still unknown, our biological world on the earth is just accidental, or a generic outcome of the inflationary cosmology paradigm


“Central dogma” in cosmology or a reductionist’s point of view

- **Inflation will fix the initial condition of the universe**
 - Quantum fluctuations generated during the inflation epoch are responsible for the origin of all the structures in the universe
- **Maybe right in principle, but in reality, can we extract any initial memory of the early universe out the motion of planets in our solar system ?**
 - Competition between the initial condition and chance/stochasticity
 - Analogous to different perspectives between particle physicists and solid-state physicists

Measurement of Spin-Orbit alignment in an Extrasolar Planetary System





- **Joshua N. Winn (MIT)**, R.W. Noyes, M.J. Holman, D.B. Charbonneau, Y. Ohta, A. Taruya, Y. Suto, N. Narita, E.L. Turner, J.A. Johnson, G.W. Marcy, R.P. Butler, & S.S. Vogt
 - **ApJ 631(2005)1215 (astro-ph/0504555)**

HD209458: the first transiting planet discovered in 1999



- Motivated by an analytical perturbation formula for the Rossier-McLaughlin effect by Ohta, Taruya & Suto: ApJ 622(2005)1118

Discovery of a retrograde/polar orbit of HAT-P-7 with Subaru via the RM effect



- Origin of the retrograde/polar orbit is unknown

Formation of retrograde planets

■ Initial condition


- Central star with 1 solar mass
- Three planets all with 1 Jupiter mass
- Circular orbits ($e=0$)
- semi-major axis: 5AU, 7.25AU, 9.5AU
- Inclination: 0.5deg, 1.0deg, 1.5deg
- Random phases **(Nagasawa et al. 2008,2011;**

■ Evolution **see also Beauge and Nesvorny 2011)**



- Simple gravitational few body system
- Orbit-crossing and scale segregation (ejection of a planet)
- formation of a close-in planet (due to tidal dissipation)
- Perturbation from the outer planet significantly changes the dynamics of the inner planet via the Kozai mechanism

Segregation of small and large scales

- **Initially similar scale orbits**
 - semi-major axis: 5AU, 7.25AU, 9.5AU
- **Chaotic evolution and energy exchange**
 - Orbit-crossing and segregation of outer and inner planets
- **Energy and angular-momentum dissipation**
 - formation of a close-in planet (due to tidal dissipation)
- **Interaction between different scales**
 - Perturbation from the outer planet (via the Kozai mechanism) results in a retrograde orbit of the inner planet





Evolution of three planets



0 1 2 3 4 5 6 7 8 time [Myr]

Xue et al.
in prep.

Co-evolution and co-existence of hierarchical structures in the universe



- **Primordial density fluctuations**
 - seeded in the inflationary epoch
- **Gravitational evolution**
 - linear
 - dissipationless nonlinear
- **Non-gravitational evolution**
 - Radiative processes
 - Energy dissipation
 - Angular-momentum transfer
 - Segregation of scales
- **Co-existence**
 - Huge difference among their time-scales

$$t_{grav} \approx \frac{1}{\sqrt{G\rho}} \approx 10\text{Gyr} \sqrt{\frac{10^{-29} \text{ g/cc}}{\rho_{universe}}} \approx 0.1\text{msec} \sqrt{\frac{10^{14} \text{ g/cc}}{\rho_{neutron star}}}$$