



$p\Xi^-$ Correlation in Relativistic Heavy Ion Collisions with Nucleon-Hyperon Interaction from Lattice QCD

Tetsuo Hatsuda¹, Kenji Morita², Akira Ohnishi², Kenji Sasaki²

¹ *iTHEMS Program and Nishina Center, RIKEN, Saitama 351-0198, Japan*

² *Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan*

Abstract

On the basis of the $p\Xi^-$ interaction extracted from (2+1)-flavor lattice QCD simulations at the physical point, the momentum correlation of p and Ξ^- produced in relativistic heavy ion collisions is evaluated. $C_{\text{SL}}(Q)$ defined by a ratio of the momentum correlations between the systems with different source sizes is shown to be largely enhanced at low momentum due to the strong attraction between p and Ξ^- in the $I = J = 0$ channel. Thus, measuring this ratio at RHIC and LHC and its comparison to the theoretical analysis will give a useful constraint on the $p\Xi^-$ interaction.

Keywords: exotic dibaryon, hyperon-nucleon force, Lattice QCD

1. Introduction

The coupled-channel Nambu-Bethe-Salpeter (NBS) wave function measured in lattice QCD [1, 2] can now provide “theoretical” information of hyperon-nucleon and hyperon-hyperon interactions through the HAL QCD method [3, 4, 5, 6]. The energy-independent non-local potentials $U(r, r')$ obtained by the method allow us to calculate the scattering phase shifts and binding energies of two baryons.

These potentials are also useful for analyzing the two-particle momentum correlations in relativistic heavy ion collisions [7]. It was recently studied in [8] that the possible spin-2 $p\Omega^-$ dibaryon state suggested by lattice QCD [9] can be probed by the $p\Omega^-$ momentum correlation at RHIC and LHC. In particular, the ratio of correlation functions between small and large collision systems, $C_{\text{SL}}(Q)$, is shown to be a good measure to extract the strong interaction effect without much contamination from the Coulomb effect [8]. In the present paper, we extend the analysis to the $p\Xi^-$ system in $I = J = 0$ channel which was recently predicted to have large attraction by the lattice QCD simulations at physical quark masses [4].

2. Lattice QCD formulation

We start with the normalized four-point function R in channel α defined by

$$R^\alpha(\vec{r}, t) \equiv \frac{\langle 0 | B_{\alpha_1}(\vec{x} + \vec{r}, t) B_{\alpha_2}(\vec{x}, t) \overline{\mathcal{F}}(0) | 0 \rangle}{\sqrt{Z_{\alpha_1} Z_{\alpha_2}} \exp[-(m_{\alpha_1} + m_{\alpha_2})t]}, \quad (1)$$

where $B_{\alpha_1}(\vec{x}, t)$ and $B_{\alpha_2}(\vec{x}, t)$ are the sink operators for octet baryons. $\sqrt{Z_{\alpha_1}}$ $\sqrt{Z_{\alpha_2}}$ are the corresponding wave-function renormalization factors, and $\mathcal{J}(0)$ is a source operator at zero initial-time to create two baryons. The coupled channel potential is obtained through the linear partial differential equation [2];

$$(D_t^\alpha - H_0^\alpha) R^\alpha(\vec{r}, t) = \int d^3 r' U^{\alpha\beta}(\vec{r}, \vec{r}') \Delta^{\alpha\beta} R^\beta(\vec{r}', t), \quad (2)$$

with $H_0^\alpha = -\frac{\nabla^2}{2\mu^\alpha}$ and $\Delta^{\alpha\beta} = \exp[-(m_{\beta_1} + m_{\beta_2})t] / \exp[-(m_{\alpha_1} + m_{\alpha_2})t]$. D_t^α is a time-derivative operator whose leading-order term reads $-\partial/\partial t$. We introduce a derivative expansion to treat the non-local potential as

$$U^{\alpha\beta}(\vec{r}, \vec{r}') = (V_{\text{LO}}^{\alpha\beta}(\vec{r}) + V_{\text{NLO}}^{\alpha\beta}(\vec{r}) + \dots) \delta(\vec{r} - \vec{r}'). \quad (3)$$

In the following, we truncate the expansion at the leading order.

We employ (2 + 1)-flavor QCD configurations on the $L^4 = 96^4$ lattice with the lattice spacing $a \simeq 0.085\text{fm}$. This corresponds to the physical size, $La = 8.1\text{fm}$, which guarantees that the finite volume effect on $U^{\alpha\beta}(\vec{r}, \vec{r}')$ is negligible. The quark masses are chosen for the system to be almost at the physical point; $m_\pi \simeq 146\text{ MeV}$ and $m_K \simeq 525\text{ MeV}$ [4]. The total number of configurations is 414×4 space-time rotations $\times 48$ wall sources. The baryon masses measured in this setup are listed below.

baryon	N	Λ	Σ	Ξ
mass [MeV]	953 ± 7	1123 ± 3	1204 ± 2	1332 ± 1

3. $p\Xi^-$ potential in $I = 0$ channel

The $S = -2$ baryon-baryon interactions including the $I=0$ $\Lambda\Lambda - N\Xi - \Sigma\Sigma$ coupled-channel system have been recently reported in [4]. In particular, one of the diagonal components $V_{N\Xi, N\Xi}(r)$ in the $(I, J) = (0, 0)$ channel (1S_0) was shown to have large attractive well at intermediate distance and relatively weak repulsive core at short distance, while $V_{N\Xi, N\Xi}(r)$ in the $(I, J) = (0, 1)$ channel (3S_1) has weaker attractive well and stronger repulsive core. Also, $V_{N\Xi, N\Xi}(r)$ in the $I = 1$ channels do not have appreciable attraction. Motivated by these observations, we parametrize the lattice results of $V_{N\Xi, N\Xi}(r)$ in the $I = 0$ channels by a combination of the Gauss and Yukawa functions as shown in Fig.1. Curves with different t correspond to the potentials obtained from $R(\vec{x}, t)$ for different t , so that the t dependence of $V(r)$ reflects typical magnitude of the systematic error of the lattice data. We found that the strong QCD attraction in Fig.1(Left) together with the Coulomb attraction leads to the 1S_0 system close to the unitary region where the inverse of the scattering length is close to zero. On the other hand, the 3S_1 system described by Fig.1(Right) has strong repulsion even with the Coulomb attraction.

4. $p\Xi^-$ momentum correlation

The correlation function of non-identical pair such as $p\Xi^-$ is given in terms of the two-particle distribution $N_{p\Xi}(\mathbf{k}_p, \mathbf{k}_\Xi)$ normalized by a product of the single particle distributions, $N_\Xi(\mathbf{k}_\Xi)N_p(\mathbf{k}_p)$,

$$C(\mathbf{Q}, \mathbf{K}) \equiv \frac{N_{p\Xi}(\mathbf{k}_p, \mathbf{k}_\Xi)}{N_p(\mathbf{k}_p)N_\Xi(\mathbf{k}_\Xi)} \simeq \frac{\int d^4 x_p \int d^4 x_\Xi S_p(x_p, \mathbf{k}_p) S_\Xi(x_\Xi, \mathbf{k}_\Xi) |\Psi_{p\Xi}(\mathbf{r}')|^2}{\int d^4 x_p S_p(x_p, \mathbf{k}_p) \int d^4 x_\Xi S_\Xi(x_\Xi, \mathbf{k}_\Xi)},$$

where relative and total momenta are defined as $\mathbf{Q} = (m_p \mathbf{k}_\Xi - m_\Xi \mathbf{k}_p)/M$ and $\mathbf{K} = \mathbf{k}_p + \mathbf{k}_\Xi$, respectively, with $M \equiv m_p + m_\Xi$. The source functions $S_i(x_i, \mathbf{k}_i) \equiv E_i \frac{dN_i}{d^3 \mathbf{k}_i d^4 x_i}$ (with $i = p, \Xi$ and $E_i = \sqrt{\mathbf{k}_i^2 + m_i^2}$) correspond to the phase space distributions of p and Ξ at freeze-out. The final state interaction after the freeze-out is described by the two-particle wave function $\Psi_{p\Xi}$ with a shifted relative coordinate $\mathbf{r}' = \mathbf{x}_\Xi - \mathbf{x}_p - \mathbf{K}(t_p - t_\Xi)/M$.




Fig. 1. The $N\Xi$ potentials in the $I = 0$ channel fitted to the (2+1)-flavor lattice QCD data at the physical point. Euclidean time used for extracting the lattice QCD potential is denoted by t . (Left) The potential in the $(I, J) = (0, 0)$ channel (1S_0). (Right) The potential in the $(I, J) = (0, 1)$ channel (3S_1).

Here we consider the static source function with spherical symmetry to extract the essential part of physics;

$$S_i(x_i, \mathbf{k}_i) \propto E_i e^{-\frac{x_i^2}{2R_i^2}} \delta(t - t_i), \quad (i = p, \Xi^-), \quad (4)$$

where R_i is a source size parameter. Assuming the equal-time emission $t_p = t_{\Xi^-}$, we obtain

$$C(Q) = \int [dr] \int \frac{d\Omega}{4\pi} |\psi^C(\mathbf{r})|^2 + \frac{1}{8} \int [dr] (|\chi_{sc}^{J=0}(r)|^2 - |\psi_0^C(r)|^2) + \frac{3}{8} \int [dr] (|\chi_{sc}^{J=1}(r)|^2 - |\psi_0^C(r)|^2), \quad (5)$$

where $[dr] = \frac{1}{2\sqrt{\pi}R^3} dr r^2 e^{-\frac{r^2}{4R^2}}$ with $R = \sqrt{(R_p^2 + R_{\Xi^-}^2)/2}$ being the effective size parameter. $\int d\Omega$ is the integration over the solid angle between \mathbf{Q} and \mathbf{r} . Note that $\psi^C(\mathbf{r})$ is the Coulomb wave function characterized by the reduced mass and the Bohr radius of the $p\Xi^-$ system. Its S-wave component is denoted by $\psi_0^C(r)$. The scattering wave functions obtained by solving the Schrödinger equation with both strong interaction and Coulomb interaction are denoted by $\chi_{sc}^{J=0}(r)$ and $\chi_{sc}^{J=1}(r)$ for the 1S_0 channel and 3S_1 channel, respectively. We assume that the $I = 1$ sector does not contribute substantially to $C(Q)$, which is supported by the fact that the $I = 1$ $p\Xi^-$ potential has only short-range repulsion [4]. The factors $1/8 = 1/2 \times 1/4$ and $3/8 = 1/2 \times 3/4$ originate from the isospin and spin multiplicities. Also, we assume that the absorptive contribution by the coupling to the $\Lambda\Lambda$ channel is negligible since it is reported to be weak due to its short range nature [4].

In [8], the ‘‘SL (small-to-large) ratio’’ was introduced: It is defined as a ratio of $C(Q)$ between the systems with different source sizes,

$$C_{\text{SL}}(Q) \equiv C_{R_{p,\Xi}=2.5\text{fm}}(Q)/C_{R_{p,\Xi}=5\text{fm}}(Q), \quad (6)$$

which has good sensitivity to the strong interaction without much contamination from the Coulomb interaction [8]. Shown in Fig.2 is $C_{\text{SL}}(Q)$ of the $p\Xi^-$ system with the Coulomb interaction under the assumption of the static source given in Eq.(4).

The large enhancement of this ratio at small Q originates from the fact that the $p\Xi^-$ system in the 1S_0 channel is close to the unitary region. The result has rather weak dependence on t , which indicates that the systematic errors of the lattice data do not affect the final results significantly. We have also checked that taking the expanding source as discussed in [8] does not change the present result.




Fig. 2. SL (small-to-large) ratio $C_{SL}(Q)$ for the momentum correlation of $p\Xi^-$ system as a function of the relative momentum Q in the case of the static source. Both the strong and Coulomb interactions are taken into account for the $p\Xi^-$ interaction. Different curves correspond to different potentials shown in Fig.1.

5. Summary

The momentum correlation of the $p\Xi^-$ system was presented by employing the $p\Xi^-$ potential extracted from the coupled channel analysis of the (2+1)-flavor lattice QCD data at the physical point. So-called the SL-ratio of the momentum correlation ($C_{SL}(Q)$) was calculated and was shown to have large enhancement at small Q due to the strong attraction between p and Ξ^- in the 1S_0 channel. Measuring this ratio at RHIC and LHC and its comparison to the present theoretical analysis will give useful constraint on the $p\Xi^-$ interaction. Such information is particularly important not only for the nature of the possible H -dibaryon coupled to $p\Xi^-$ [4] but also for the properties of Ξ -hypernuclei [10] and for Ξ^- in the central core of the neutron star [11].

Acknowledgments

This work is supported in part by MEXT Grant-in-Aid for Scientific Research (JP15K17667, 24105008), SPIRE (Strategic Program for Innovative REsearch) Field 5 project and "Priority Issue on Post-K computer" (Elucidation of the Fundamental Laws and Evolution of the Universe). K.M. was supported by JSPS Grant 16K05349, and National Science Center, Poland under grants: Maestro DEC-2013/10/A/ST2/00106. T.H. was partially supported by RIKEN iTHES Project and iTHEMS Program. The authors thank HAL QCD Collaboration for providing us with the data for $p\Xi^-$ interactions and for stimulating discussions.

References

- [1] N. Ishii, S. Aoki and T. Hatsuda, Phys. Rev. Lett. **99** (2007) 022001 [nucl-th/0611096].
- [2] N. Ishii *et al.* [HAL QCD Collaboration], Phys. Lett. B **712** (2012) 437 [arXiv:1203.3642 [hep-lat]].
- [3] T. Doi *et al.* [HAL QCD Collaboration], PoS LATTICE **2016** (2017) 110 [arXiv:1702.01600 [hep-lat]].
- [4] K. Sasaki *et al.* [HAL QCD Collaboration], PoS LATTICE **2016** (2017) 116 [arXiv:1702.06241 [hep-lat]].
- [5] N. Ishii *et al.* [HAL QCD Collaboration], PoS LATTICE **2016** (2017) 127 [arXiv:1702.03495 [hep-lat]].
- [6] H. Nemura *et al.* [HAL QCD Collaboration], PoS LATTICE **2016** (2017) 101 [arXiv:1702.00734 [hep-lat]].
- [7] S. Cho *et al.* [ExHIC Collaboration], arXiv:1702.00486 [nucl-th].
- [8] K. Morita, A. Ohnishi, F. Etminan and T. Hatsuda, Phys. Rev. C **94** (2016) 031901 [arXiv:1605.06765 [hep-ph]].
- [9] F. Etminan *et al.* [HAL QCD Collaboration], Nucl. Phys. A **928** (2014) 89 [arXiv:1403.7284 [hep-lat]].
- [10] K. Nakazawa *et al.*, PTEP **2015** (2015) 033D02.
- [11] see e.g. a review, G. Baym, T. Hatsuda, T. Kojo, C. J. Pethick, P. D. Powell, Y. Song, and T. Takatsuka, (in preparation).



ELSEVIER