

Noncommutative Solitons and Integrable Systems

Masashi HAMANAKA

(Dept. of Math., Nagoya U., JAPAN)

Ref: MH, “NC solitons and integrable systems,”
(to appear soon) [[hep-th/0504001](#)].

cf. MH, “Commuting flows and conservation laws
for NC Lax hierarchies,” [[hep-th/0311206](#)]

[J.Math.Phys.46 \(2005\)](#) to be published on April 1.

1. Introduction

Successful points on NC theories

- Appearance of **new** physical objects
- Description of **real** physics
- Various **successful applications** to D-brane dynamics etc.

NC Solitons play important roles
(Integrable!)

Final goal: NC extension of **all** soliton theories

Integrable equations in diverse dimensions

4	Anti-Self-Dual Yang-Mills eq. (instantons) $F_{\mu\nu} = -\tilde{F}_{\mu\nu}$	NC extension (Successful)
3	Bogomol'nyi eq. (monopoles)	NC extension (Successful)
2 (+1)	KP eq. BCS eq. DS eq. ...	NC extension (This talk)
1 (+1)	KdV eq. Boussinesq eq. NLS eq. Burgers eq. sine-Gordon eq. (affine) Toda field eq. ...	NC extension (This talk)

↑
Dim. of space

Ward's observation: Almost all
integrable equations are
reductions of the ASDYM eqs.

R.Ward, Phil.Trans.Roy.Soc.Lond.A315(85)451

ASDYM eq.



Reductions

KP eq. BCS eq. Ward's chiral model
 KdV eq. Boussinesq eq.
 NLS eq. mKdV eq.
sine-Gordon eq. Burgers eq. ...

(Almost all !?)

e.g. [The book of Mason&Woodhouse]

NC Ward's observation: Almost all
NC integrable equations are
reductions of the NC ASDYM eqs.

MH&K.Toda, PLA316(03)77[hep-th/0211148]

NC ASDYM eq.

Successful

↓ NC Reductions Reductions

NC KP eq. NC BCS eq. NC Ward's chiral model
NC KdV eq. NC Boussinesq eq.
NC NLS eq. NC mKdV eq.
NC sine-Gordon eq. NC Burgers eq. ...

(Almost all !?)

Successful?

A general framework is needed

Program of NC extension of soliton theories



- (i) Confirmation of NC Ward's conjecture
 - NC twistor theory → geometrical origin
 - D-brane interpretations → applications to physics
 - (ii) Completion of NC Sato's theory
 - Existence of "hierarchies" → various soliton eqs.
 - Existence of infinite conserved quantities
 - infinite-dim. hidden symmetry
 - Construction of multi-soliton solutions
 - Theory of tau-functions → structure of the solution spaces and the symmetry
- (i),(ii) → complete understanding of the NC soliton theories

Plan of this talk

1. Introduction
2. Review of soliton theories
3. NC Sato's theory
4. Conservation Laws
5. Exact Solutions and Ward's conjecture
6. Conclusion and Discussion

2. Review of Soliton Theories

- KdV equation : describe shallow water waves



solitary wave = soliton

This configuration satisfies

$$u = 2k^2 \cosh^{-2}(kx - 4k^3 t)$$

$$\dot{u} + u''' + 6u'u = 0 : \text{KdV eq. [Korteweg-de Vries, 1895]}$$

This is a typical integrable equation.

Let's solve it now !

- Hirota's method [PRL27(1971)1192]

$$\dot{u} + u''' + 6u'u = 0 \quad : \text{naively hard to solve}$$

$$\downarrow \quad u = 2\partial_x^2 \log \tau$$

$$\tau\dot{\tau}' - \tau'\dot{\tau} + 3\tau''\tau'' - 4\tau'\tau''' + \tau\tau'''' = 0$$

Hirota's bilinear relation : more complicated ?

A solution: $\tau = 1 + e^{2(kx - \omega t)}, \quad \omega = 4k^3$


$\rightarrow u = 2k^2 \cosh^{-2}(kx - 4k^3 t) : \text{The solitary wave !}$
(1-soliton solution)

- 2-soliton solution

$$\tau = 1 + A_1 e^{2\theta_1} + A_2 e^{2\theta_2} + BA_1 A_2 e^{2(\theta_1 + \theta_2)},$$

$$\theta_i = k_i x - 4k_i^3 t, \quad B = \frac{k_1 - k_2}{k_1 + k_2}$$

Scattering process



The shape and velocity is preserved ! (stable)

The positions are shifted ! (Phase shift)

There are many other soliton eqs.
with (similar) interesting properties

- KP equation (2-dim. KdV equation)

: describe 2-dim shallow water waves

$$u_t + u_{xxx} + 6u_x u + \partial_x^{-1} u_{yy} = 0 \quad \text{KP} \quad \xrightarrow{\partial_y = 0} \quad \text{KdV}$$

$$u_x := \frac{\partial u}{\partial x} \quad \partial_x^{-1} := \int^x dx' \quad \text{etc.}$$

Sato's theorem: [M.Sato&Y.Sato, 1981]

The solution space of KP eq. is an infinite-dim.

Grassmann mfd. (determined by tau -fcns.)

Many other soliton eqs. are obtained from KP.

[See e.g. The book of Miwa-Jimbo-Date (Cambridge UP, 2000)]

3. NC Sato's Theory

- Sato's Theory : one of the most beautiful theory of solitons
 - Based on the existence of hierarchies and tau-functions
- Sato's theory reveals essential aspects of solitons:
 - Construction of exact solutions
 - Structure of solution spaces
 - Infinite conserved quantities
 - Hidden infinite-dim. symmetry

Let's discuss NC extension of Sato's theory

Derivation of soliton equations

- Prepare a Lax operator which is a pseudo-differential operator

$$L := \partial_x + u_2 \partial_x^{-1} + u_3 \partial_x^{-2} + u_4 \partial_x^{-3} + \dots$$

$$u_k = u_k(x^1, x^2, x^3, \dots)$$

- Introduce a differential operator

$$B_m := (L * \dots * L)_{\geq 0}$$

m times

Noncommutativity
is introduced here:

$$[x^i, x^j] = i\theta^{ij}$$

- Define NC (KP) hierarchy:

$$\frac{\partial L}{\partial x^m} = [B_m, L]_*$$

Here all products are
star product:

$$\begin{array}{ll} \partial_m u_2 \partial_x^{-1} + & f_{m2}(u) \partial_x^{-1} + \\ \partial_m u_3 \partial_x^{-2} + & f_{m3}(u) \partial_x^{-2} + \\ \partial_m u_4 \partial_x^{-3} + \dots & f_{m4}(u) \partial_x^{-3} + \dots \end{array}$$



Each coefficient yields
a differential equation.

Negative powers of differential operators

$$\partial_x^n \circ f := \sum_{j=0}^{\infty} \binom{n}{j} (\partial_x^j f) \partial_x^{n-j}$$

$$\frac{n(n-1)(n-2)\cdots(n-(j-1))}{j(j-1)(j-2)\cdots 1}$$

: binomial coefficient
which can be extended
to negative n
→ negative power of
differential operator
(well-defined !)

$$\partial_x^3 \circ f = f\partial_x^3 + 3f\partial_x^2 + 3f\partial_x + f''''$$

$$\partial_x^2 \circ f = f\partial_x^2 + 2f\partial_x + f''$$

$$\partial_x^{-1} \circ f = f\partial_x^{-1} - f\partial_x^{-2} + f\partial_x^{-3} - \dots$$

$$\partial_x^{-2} \circ f = f\partial_x^{-2} - 2f\partial_x^{-3} + 3f\partial_x^{-4} - \dots$$

Star product:

$$f(x) * g(x) := f(x) \exp\left(\frac{i}{2} \theta^{ij} \overleftarrow{\partial}_i \overrightarrow{\partial}_j\right) g(x)$$

which makes theories "noncommutative":

$$[x^i, x^j]_* := x^i * x^j - x^j * x^i = i\theta^{ij}$$

Closer look at NC KP hierarchy

For $m=2$

$$\partial_x^{-1}) \quad \partial_2 u_2 = \underline{2u_3'} + u_2''$$

$$\partial_x^{-2}) \quad \partial_2 u_3 = \underline{2u_4'} + u_3'' + 2u_2 * u_2' + 2[u_2, u_3]_*$$

$$\partial_x^{-3}) \quad \partial_2 u_4 = \underline{2u_5'} + u_4'' + 4u_3 * u_2' - 2u_2 * u_2'' + 2[u_2, u_4]_*$$

⋮

Infinite kind of fields are represented
in terms of one kind of field $u_2 \equiv u$

MH&K.Toda, [hep-th/0309265]

$$u_x := \frac{\partial u}{\partial x}$$

$$\partial_x^{-1} := \int^x dx'$$

For $m=3$

$$\partial_x^{-1}) \quad \partial_3 u_2 = u_2''' + 3u_3'' + 3u_4'' + 3u_2' * u_2 + 3u_2 * u_2'$$

⋮



$$u_t = \frac{1}{4} u_{xxx} + \frac{3}{4} (u_x * u + u * u_x) + \frac{3}{4} \partial_x^{-1} u_{yy} + \frac{3}{4} [u, \partial_x^{-1} u_{yy}]_*$$

(2+1)-dim.
NC KP equation

and other NC equations
(NC KP hierarchy equations)

$$u = u(x^1, x^2, x^3, \dots)$$

$$\begin{array}{ccc} \updownarrow & \updownarrow & \updownarrow \\ x & y & t \end{array}$$

etc.

(KP hierarchy) ^{reductions} \rightarrow (various hierarchies.)

- (Ex.) KdV hierarchy

Reduction condition


$$L^2 = B_2 (=:\partial_x^2 + u) \quad : \text{2-reduction}$$

gives rise to NC KdV hierarchy


which includes (1+1)-dim. NC KdV eq.:

$$u_t = \frac{1}{4}u_{xxx} + \frac{3}{4}(u_x * u + u * u_x)$$

Note $\frac{\partial u}{\partial x_{2N}} = 0$: dimensional reduction in x_{2N} directions



-reduction of NC KP hierarchy yields
wide class of other NC hierarchies

- No-reduction \rightarrow NC KP $(x, y, t) = (x^1, x^2, x^3)$
 - 2-reduction \rightarrow NC KdV $(x, t) = (x^1, x^3)$
 - 3-reduction \rightarrow NC Boussinesq $(x, t) = (x^1, x^2)$
 - 4-reduction \rightarrow NC Coupled KdV ...
 - 5-reduction \rightarrow ...
 - 3-reduction of BKP \rightarrow NC Sawada-Kotera
 - 2-reduction of mKP \rightarrow NC mKdV
 - Special 1-reduction of mKP \rightarrow NC Burgers
 - ...
- Noncommutativity should be introduced into space-time coords
- 

4. Conservation Laws

- Conservation laws: $\partial_t \sigma = \partial_i J^i$ σ : Conserved density
time \nearrow space \nwarrow

Then $Q := \int_{space} dx \sigma$ is a conserved quantity.

$$\because \partial_t Q = \int_{space} dx \partial_t \sigma = \int_{spatial} dS_i J^i = 0$$

infinity

Conservation laws for the hierarchies

$$\partial_m res_{-1} L^n = \partial_x J + \theta^{ij} \partial_j \Xi_i$$

time \nearrow space \nearrow ↑

I have succeeded in the evaluation explicitly !

$res_{-r} L^n$: coefficient
of ∂_x^{-r} in L^n

Noncommutativity should be introduced
in space-time directions only. \rightarrow

$$t \equiv x^m$$

∂_j should be space or time derivative
 \rightarrow ordinary conservation laws !

Infinite conserved densities for the NC soliton eqs. ($n=1,2,\dots$)

$$\sigma_n = \text{res}_{-1} L^n + \theta^{im} \sum_{k=0}^{m-1} \sum_{l=0}^k \binom{k}{l} (\partial_x^{k-l} \text{res}_{-(l+1)} L^n) \diamond (\partial_i \text{res}_k L^m)$$

$t \equiv x^m$ $\text{res}_r L^n$: coefficient of ∂_x^r in L^n

\diamond : Strachan's product (commutative and non-associative)

$$f(x) \diamond g(x) := f(x) \left(\sum_{s=0}^{\infty} \frac{(-1)^s}{(2s+1)!} \left(\frac{1}{2} \theta^{ij} \vec{\partial}_i \vec{\partial}_j \right)^{2s} \right) g(x)$$

MH, JMP46 (2005)
[hep-th/0311206]

This suggests infinite-dimensional symmetries would be hidden.

We can calculate the explicit forms of conserved densities for the wide class of NC soliton equations.

- Space-Space noncommutativity:

NC deformation is slight: $\sigma_n = \text{res}_{-1} L^n$

- Space-time noncommutativity

NC deformation is drastical:

– Example: NC KP and KdV equations $([t, x] = i\theta)$

$$\sigma_n = \text{res}_{-1} L^n - 3\theta((\text{res}_{-1} L^n) \diamond u'_3 + (\text{res}_{-2} L^n) \diamond u'_2)$$

meaningful ?

5. Exact Solutions and Ward's conjecture

- We have found **exact N-soliton solutions** for the wide class of NC hierarchies.

- 1-soliton solutions are all the same as commutative ones because of

$$f(x-vt) * g(x-vt) = f(x-vt)g(x-vt)$$

- Multi-soliton solutions behave in almost the same way as commutative ones **except for phase shifts.**
- Noncommutativity affects **the phase shifts**

Exact N-soliton solutions of the NC KP hierarchy

$L = \Phi \partial_x \Phi^{-1}$ solves the NC KP hierarchy !

$\Phi f := \left| W(y_1, \dots, y_N, f) \right|_{N+1, N+1}$ quasi-determinant of Wronski matrix

$$y_i = \exp \xi(x, \alpha_i) + a_i \exp \xi(x, \beta_i)$$

Etingof-Gelfand-Retakh

$$\xi(x, \alpha) = x_1 \alpha + x_2 \alpha^2 + x_3 \alpha^3 + \dots \quad [\text{q-alg/9701008}]$$

The exact solutions are actually N-soliton solutions !

Noncommutativity might affect the phase shift by $\theta^{ij} \omega_i k_j$

$\therefore \exp(\omega_i t - k_i x) * \exp(\omega_j t - k_j x)$ [MH, work in progress]

$$\cong \exp([t, x]_* \omega_{[i} k_{j]}) \exp((\omega_i + \omega_j)t - (k_i + k_j)x)$$

Exactly solvable!

NC Burgers hierarchy

MH&K.Toda, JPA36(03)11981 [hep-th/0301213]

- NC (1+1)-dim. Burgers equation:

$$\dot{u} = u'' + 2u * u' : \text{Non-linear \&}$$

Infinite order diff. eq. w.r.t. time ! (Integrable?)

NC Cole-Hopf transformation

$$u = \tau^{-1} * \tau' \quad (\xrightarrow{\theta \rightarrow 0} \partial_x \log \tau)$$

(NC) Diffusion equation:

$$\dot{\tau} = \tau'' : \text{Linear \& first order diff. eq. w.r.t. time}$$

(Integrable !)

A solution :

$$\tau = 1 + \sum_{l=1}^N e^{k_l^2 t} * e^{\pm k_l x} = 1 + \sum_{l=1}^N \underline{e^{\frac{i}{2} k_l^3 \theta}} e^{k_l^2 t \pm k_l x}$$

Deformed!

NC Ward's conjecture (NC NLS eq.)

- Reduced ASDYM eq.: $x^\mu \rightarrow (t, x)$

Legare,
[hep-th/0012077]

(i) $B' = 0$

(ii) $C' - \dot{A} + [A, C]_* = 0$

(iii) $A' - \dot{B} + [C, B]_* = 0$

A, B, C: 2 times 2
matrices (gauge fields)



Further
Reduction:

$$A = \begin{pmatrix} 0 & q \\ -\bar{q} & 0 \end{pmatrix}, B = \frac{i}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, C = i \begin{pmatrix} q * \bar{q} & \bar{q}' \\ q' & -\bar{q} * q \end{pmatrix}$$

$$(ii) \Rightarrow \begin{pmatrix} 0 & i\dot{q} - q'' - 2q * \bar{q} * q \\ i\dot{\bar{q}} + \bar{q}'' - 2\bar{q} * q * \bar{q} & 0 \end{pmatrix} = 0$$

NOT traceless

$$i\dot{q} = q'' + 2q * \bar{q} * q \quad : \text{NC NLS eq. !!!}$$

Note: $A, B, C \in u(2) \xrightarrow{\theta \rightarrow 0} su(2)$ **U(1) part is necessary !**

NC Ward's conjecture (NC Burgers eq.)

- Reduced ASDYM eq.: $x^\mu \rightarrow (t, x)$ MH & K.Toda, JPA36
[hep-th/0301213 v2]

$$(i) \quad \dot{A} + \underline{[B, A]}_* = 0$$

$$(ii) \quad \dot{C} - B' + \underline{[B, C]}_* = 0, \quad A, B, C \in \underline{u(1)}$$

$$\cong u(\infty)$$

should remain



Further

Reduction: $A = 0, B = u' - u^2, C = u$

$$(ii) \Rightarrow \dot{u} = u'' + 2u * u'$$

: NC Burgers eq. !!!

Note: Without the commutators $[,]$, (ii) yields:

$$\dot{u} = u'' + \underline{u' * u + u * u'} \quad : \text{neither linearizable nor Lax form}$$

Symmetric

NC Ward's conjecture (NC KdV eq.)

- **Reduced ASDYM eq.:** $x^\mu \rightarrow (t, x)$ MH, (HOT result)
[hep-th/0504001]

(i) $[A, B]_* = 0$

(ii) $A' - C' + [A, C]_* + [D, B]_* = 0$

(iii) $\dot{C} - D' + [C, D]_* = 0$

A, B, C: 2 times 2
matrices (gauge fields)

↓ Further Reduction (different type of Mason-Sparling!):

$$A = \frac{1}{2} \begin{pmatrix} 0 & 0 \\ -u & 0 \end{pmatrix}, B = \begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix}, C = \begin{pmatrix} 0 & 1 \\ -u & 0 \end{pmatrix}, D = \frac{1}{4} \begin{pmatrix} -u' & 2u \\ -u'' - 2u^2 & u' \end{pmatrix}$$

(iii) $\Rightarrow \frac{1}{4} \begin{pmatrix} 0 & 0 \\ 4\dot{u} + u''' + 3(u' * u + u * u') & 0 \end{pmatrix} = 0$

$$\dot{u} + \frac{1}{4} u''' + \frac{3}{4} (u' * u + u * u') = 0 \quad : \text{ NC KdV eq. !!!}$$

Note: $A, B, C \in sl(2, R)$

Bad news for D-brane picture?
Good news for twistor picture !

6. Conclusion and Discussion

- Confirmation of NC Ward's conjecture Going well
 - NC twistor theory → geometrical origin
 - D-brane interpretations → applications to physics
 - Work in progress → [NC book of Mason&Woodhouse ?]
- Completion of NC Sato's theory
 - Existence of ``hierarchies'' → Solved!
 - Existence of infinite conserved quantities Successful
 - infinite-dim. hidden symmetry
 - Construction of multi-soliton solutions Successful
 - Theory of tau-functions → description of the symmetry and the soliton solutions Work in progress