

Anomaly and Superconnection

菅野 颯人

京都大学 基礎物理学研究所 (YITP) D1

Based on arXiv:2106.01591 [hep-th]

Now, published in PTEP. (Selected as an editor's choice in PTEP.)

Work with 杉本茂樹 (基研)



Motivation


QFTを理解したい

- 知っていること
 - 弱結合の理論(例:QED)は摂動論で理解できる
 - 強結合の理論(例:QCD)は良く分からない
- 高エネルギー(UV)の作用が分かっている理論でも、低エネルギー(IR)で強結合の場合、IRでどのような理論になるのか非自明
 - QCD
 - 物性理論
- **アノマリー(量子異常)**はIRの理論を理解するために使える!
 - 't Hooft anomaly matching
 - UVの理論の持つアノマリーをIRの理論(有効作用、EFT)は再現しないとならない
 - IRの理論の形を強く制限する

What is “anomaly”? (1)

Anomaly (Quantum Anomaly)

An classical action has some symmetries, but sometimes these symmetries disappear in quantum theory.



e.g.) $\pi^0 \rightarrow 2\gamma$

- In massless QCD, there is a chiral symmetry $SU(N_f)_L \times SU(N_f)_R$.

N_f : # of flavors

- If there is NO anomaly, π^0 never decays.
- However, π^0 decays into 2γ , because of an anomaly!

Theories what we want to think (1)

Let us consider 4dim action contain fermions.

$$S = \int d^4x \bar{\psi} i \not{D} \psi = \int d^4x \bar{\psi} i \gamma^\mu (\partial_\mu + A_\mu) \psi$$

- This action is massless, so it has a chiral symmetry $U(N_f)_L \times U(N_f)_R$.
- There also be a $U(1)_A$ anomaly.

- Add mass term

- Mass term breaks the chiral symmetry.

$$S = \int d^4x \bar{\psi} \left(i \not{D} + m \right) \psi$$

- Let the mass depend on the spacetime.


- This mass is almost same as the Higgs field.
- How change the symmetry and the anomaly?

$$S = \int d^4x \bar{\psi} \left(i \not{D} + m(x) \right) \psi$$

The spacetime dependent mass

What is “the spacetime dependent mass”?

- e.g.) Domain wall fermions
 - One way to realize chiral fermions on the lattice.
 - Consider 5dim spacetime, and realize 4dim fermions on $m(x) = 0$ subspace.
- Chiral anomalies with Higgs fields
 - If Higgs fields change as bifundamental under the $U(N_f)_L \times U(N_f)_R$ chiral symmetry, the action is invariant for the symmetry.
 - It is known that chiral anomalies are not changed by adding Higgs fields.
 - See Fujikawa-san's text book.



$$S = \int d^4x \bar{\psi} \left(i \not{D} + h(x) \right) \psi$$

The spacetime dependent mass


What is “the spacetime dependent mass”?

- e.g.) Dom

- One wa
- Consid
- fermion

- Chiral an

- If Higg
- $U(N_f)_L$
- invariant
- It is kn
- by add
- See Fu





nions on the
d realize 4di
ce.

elds
ndamental u
etry, the act

lies are not

κ.



$$\int \dots \left(i \not{D} + h(x) \right) \psi$$

Theories what we want to think (2)

How about the spacetime dependent mass?

- The chiral anomaly is changed by the mass!!

- Difference between Higgs and mass
 - Higgs field : bounded
 - Spacetime dependent mass : unbounded

$$S = \int d^d x \bar{\psi} \left(i \not{D} + m(x) \right) \psi$$

- If the mass diverges at some points, it contributes to the anomaly.

- This contribution might be unknown.
- We can find the anomaly in any dimension.

- The anomaly can be written by “superconnection.”

$$\mathcal{A} = \begin{pmatrix} A_R & iT^\dagger \\ iT & A_L \end{pmatrix}$$

Plan

1. Introduction (5)

- What is anomaly?
- Theories what we want to think

2. Derivation (10)

- How to calculate anomalies
- The anomaly for massless case
- The anomaly for massive case
- Superconnection
- The result

3. Application (7)

- Kink, vortex
- With boundary

4. Index theorem (5)

- Index for massive case
- APS index theorem

5. String theory (5)

- Tachyon condensation

6. Conclusion (1)

How to calculate anomalies

['79 Fujikawa]

Fujikawa method

- There are several ways to calculate anomalies.
- Today, we focus on the Fujikawa method.
 - Consider path integral for fermions.
 - Anomaly = Jacobian comes from path integral measure
 - We only consider perturbative anomalies.
- We calculate $\log \mathcal{J}$ for anomalies in the last part of this talk.

$$Z[A] = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S}$$

e.g.) local $U(1)_V$ transformation

$$\begin{aligned} \psi(x) &\rightarrow e^{i\alpha(x)} \psi(x), \\ \bar{\psi}(x) &\rightarrow \bar{\psi}(x) e^{-i\alpha(x)} \end{aligned}$$

$$\begin{aligned} \mathcal{D}\psi \mathcal{D}\bar{\psi} &\rightarrow \mathcal{D}\psi' \mathcal{D}\bar{\psi}' = \mathcal{J} \mathcal{D}\psi \mathcal{D}\bar{\psi} \\ &= e^{-i \int d^4x \alpha(x) \mathcal{A}(x)} \mathcal{D}\psi \mathcal{D}\bar{\psi} \end{aligned}$$

anomaly

$$\log \mathcal{J} = -i \int d^4x \alpha(x) \mathcal{A}(x)$$

Chiral symmetry

Anomalous symmetries we calculate

- $U(N_f)_L \times U(N_f)_R$ chiral symmetry
 - For **even** dimension
 - Because chirality operators exist only even dimensions.
 - Weyl fermions couple to $U(N_f)_L$ background gauge field A_μ^L and $U(N_f)_R$ background gauge field A_μ^R .

$$S = \int d^d x \bar{\psi} i \gamma^\mu \left\{ \partial_\mu + A_\mu^R P_+ + A_\mu^L P_- \right\} \psi = \int d^d x (\bar{\psi}_L, \bar{\psi}_R) i \begin{pmatrix} 0 & \sigma^\mu \\ \sigma^{\mu\dagger} & 0 \end{pmatrix} \left\{ \partial_\mu + \begin{pmatrix} A_\mu^R & 0 \\ 0 & A_\mu^L \end{pmatrix} \right\} \begin{pmatrix} \psi_R \\ \psi_L \end{pmatrix}$$

$$= \int d^d x \left\{ \bar{\psi}_R i \sigma^{\mu\dagger} (\partial_\mu + A_\mu^R) \psi_R + \bar{\psi}_L i \sigma^\mu (\partial_\mu + A_\mu^L) \psi_L \right\}$$

For **odd** dimension

- $U(N_f)$ flavor symmetry
 - For **odd** dimension
 - No perturbative anomaly as usual.
 - Dirac fermions couple to $U(N_f)$ background gauge field.

$$S = \int d^d x \bar{\psi} i \gamma^\mu \left\{ \partial_\mu + A_\mu \right\} \psi$$

For **even** dimension

$$P_\pm = \frac{1 \pm \gamma_{d+1}}{2}$$

The anomaly for massless cases

e.g.) fermions in 4dim

- Mass less case
 - With $U(N_f)_L \times U(N_f)_R$ chiral sym.
 - $U(1)_V$ anomaly is written by the field strengths.
- With a Higgs field
 - With $U(N_f)_L \times U(N_f)_R$ chiral sym.
 - The $U(1)_V$ anomaly is same for massless case.
- How about the massive case?

$$\begin{aligned} S &= \int d^4x \bar{\psi} i \gamma^\mu \left\{ \partial_\mu + \begin{pmatrix} A_\mu^L & 0 \\ 0 & A_\mu^R \end{pmatrix} \right\} \psi \\ &= \int d^4x \left\{ \bar{\psi}_R i \sigma^{\mu\dagger} (\partial_\mu + A_\mu^R) \psi_R + \bar{\psi}_L i \sigma^\mu (\partial_\mu + A_\mu^L) \psi_L \right\} \end{aligned}$$

$$\begin{aligned} \log \mathcal{J} &= \frac{i}{32\pi^2} \int d^4x \alpha(x) \epsilon^{\mu\nu\rho\sigma} \text{tr} [F_{\mu\nu}^R F_{\rho\sigma}^R - F_{\mu\nu}^L F_{\rho\sigma}^L] \\ &= \frac{i}{8\pi^2} \int \alpha(x) \text{tr} [F^R \wedge F^R - F^L \wedge F^L] \end{aligned}$$

$$\begin{aligned} S &= \int d^4x \bar{\psi} (i \not{D} + h(x)) \psi \\ &= \int d^4x \left\{ \bar{\psi}_R i \sigma^{\mu\dagger} D_\mu^R \psi_R + \bar{\psi}_L i \sigma^\mu D_\mu^L \psi_L + \bar{\psi}_L h \psi_R + \bar{\psi}_R h^\dagger \psi_L \right\} \end{aligned}$$

The anomaly for massive case (1)

Let us consider spacetime dependent mass!

- The action for general even dim with $U(N_f)_L \times U(N_f)_R$ symmetry is,

$$\begin{aligned} S &= \int d^d x \bar{\psi} \left[i\gamma^\mu \left\{ \partial_\mu + \begin{pmatrix} A_\mu^R & 0 \\ 0 & A_\mu^L \end{pmatrix} \right\} + \begin{pmatrix} im(x) & 0 \\ 0 & im^\dagger(x) \end{pmatrix} \right] \psi \\ &= \int d^d x \left\{ \bar{\psi}_R i\sigma^{\mu\dagger} D_\mu^R \psi_R + \bar{\psi}_L i\sigma^\mu D_\mu^L \psi_L + \bar{\psi}_L im(x) \psi_R + \bar{\psi}_R im^\dagger(x) \psi_L \right\} \\ &\equiv \int d^d x \bar{\psi} \mathcal{D} \psi \end{aligned}$$

- We assume this \mathcal{D} is the Dirac op. for massive case.
 - \mathcal{D} is non-Hermitian.
- For odd dim case, there is only $U(N_f)$ sym, we put $A_\mu = A_\mu^R = A_\mu^L$ and $m = m^\dagger$.

The anomaly for massive case (2)

Calculate the $U(1)_V$ anomaly for this action by Fujikawa method.

- We **take $m(x)$ divergent**.

$$|m(x^I)| \rightarrow \infty \quad (|x^I| \rightarrow \infty)$$

- I denotes some directions which $m(x)$ changes its values.
- If $m(x)$ does not diverge, the anomaly is same for the massless case.

- We use heat kernel regularization.

$$e^{-\frac{\mathcal{D}^\dagger \mathcal{D}}{\Lambda^2}}, \quad e^{-\frac{\mathcal{D} \mathcal{D}^\dagger}{\Lambda^2}} \quad \Lambda \text{ is the UV cut-off.}$$

- Generalizations

- It is easy to get the anomaly for any dimension.
- It is also easy to get the anomaly for $U(N_f)_L \times U(N_f)_R$, not only for $U(1)_V$.

The anomaly for massive case (3)

e.g.) In 4dim case, the $U(1)_V$ anomaly is, $\tilde{m} = m/\Lambda$

Λ is UV cut-off
comes from
heat kernel
regularization.

$$\begin{aligned} \log \mathcal{J} = & \frac{i}{(2\pi)^2} \int d^4x \alpha(x) \text{tr} \left[\epsilon^{\mu\nu\rho\sigma} \left\{ \frac{1}{8} \left(F_{\mu\nu}^R F_{\rho\sigma}^R - F_{\mu\nu}^L F_{\rho\sigma}^L \right) \right. \right. \\ & + \frac{1}{12} \left(D_\mu \tilde{m}^\dagger D_\nu \tilde{m} F_{\rho\sigma}^R - D_\mu \tilde{m} D_\nu \tilde{m}^\dagger F_{\rho\sigma}^L + F_{\mu\nu}^R D_\rho \tilde{m}^\dagger D_\sigma \tilde{m} \right. \\ & \left. \left. - F_{\mu\nu}^L D_\rho \tilde{m} D_\sigma \tilde{m}^\dagger - D_\mu \tilde{m} F_{\nu\rho}^R D_\sigma \tilde{m}^\dagger + D_\mu \tilde{m}^\dagger F_{\nu\rho}^L D_\sigma \tilde{m} \right) \right. \\ & \left. \left. + \frac{1}{24} \left(D_\mu \tilde{m}^\dagger D_\nu \tilde{m} D_\rho \tilde{m}^\dagger D_\sigma \tilde{m} - D_\mu \tilde{m} D_\nu \tilde{m}^\dagger D_\rho \tilde{m} D_\sigma \tilde{m}^\dagger \right) \right\} \right] e^{-\tilde{m}^\dagger \tilde{m}} \end{aligned}$$

- This result seems very complicated...
- Can we rewrite it more simple way? → **Superconnection!**

Superconnection (1)

[’85 Quillen]

- We define the superconnections for even and odd dimensions.
- This is made by Quillen, who is a mathematician, in 1985.

Even dimension

- Superconnection

$$\mathcal{A} = \begin{pmatrix} A_R & iT^\dagger \\ iT & A_L \end{pmatrix}$$

$A_R : U(N_f)_R$ gauge field (1-form)

$A_L : U(N_f)_L$ gauge field (1-form)

$T : U(N_f)_L \times U(N_f)_R$ bifundamental scalar field (0-form)

- Field strength

$$\mathcal{F} = d\mathcal{A} + \mathcal{A}^2$$

$$\equiv \begin{pmatrix} F^R - T^\dagger T & iDT^\dagger \\ iDT & F^L - TT^\dagger \end{pmatrix}$$

- Supertrace

$$\text{Str} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \text{tr}(a) - \text{tr}(d)$$

Superconnection (2)

[’85 Quillen]

Odd dimension

- Superconnection

$$\mathcal{A} = \begin{pmatrix} A & iT \\ iT & A \end{pmatrix} \quad \begin{array}{l} A : U(N_f) \text{ gauge field (1-form)} \\ T : U(N_f) \text{ adjoint scalar field (0-form)} \end{array}$$

- Field strength $\mathcal{F} \equiv d\mathcal{A} + \mathcal{A}^2$

$$= \begin{pmatrix} F - T^2 & iDT \\ iDT & F - T^2 \end{pmatrix}$$

- Supertrace

$$\text{Str} \begin{pmatrix} a & b \\ b & a \end{pmatrix} = \sqrt{2i} \text{tr}(b)$$

We apply superconnection to write the anomaly.

The result (1)

- We can rewrite the $U(1)_V$ anomaly by superconnection.

$$\log \mathcal{J} = -i \left(\frac{i}{2\pi} \right)^{\frac{d}{2}} \int \alpha(x) \text{Str} [e^{\mathcal{F}}] \Big|_{d\text{-form}} \quad \left| \begin{array}{l} \tilde{m} = m/\Lambda \\ \mathcal{F} = d\mathcal{A} + \mathcal{A}^2 \\ \equiv \begin{pmatrix} F^R - T^\dagger T & iDT^\dagger \\ iDT & F^L - TT^\dagger \end{pmatrix} \\ \text{Str} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \text{tr}(a) - \text{tr}(d) \end{array} \right.$$

$$\mathcal{F} \equiv \begin{pmatrix} F^R - \tilde{m}^\dagger \tilde{m} & i(D\tilde{m})^\dagger \\ iD\tilde{m} & F^L - \tilde{m}\tilde{m}^\dagger \end{pmatrix}$$

- For odd dimension case, put $A_\mu = A_\mu^R = A_\mu^L$ and $m = m^\dagger$. Then, we get $U(1)$ anomaly.
 - In odd dimension, the definition of Str is different from even dim case.

$$\text{Str} \begin{pmatrix} a & b \\ b & a \end{pmatrix} = \sqrt{2i} \text{tr}(b)$$

The result (2)

$$\log \mathcal{J} = -i \left(\frac{i}{2\pi} \right)^{\frac{d}{2}} \int \alpha(x) \text{Str} [e^{\mathcal{F}}] \Big|_{d\text{-form}} \quad \mathcal{F} \equiv \begin{pmatrix} F^R - \tilde{m}^\dagger \tilde{m} & i(D\tilde{m})^\dagger \\ iD\tilde{m} & F^L - \tilde{m}\tilde{m}^\dagger \end{pmatrix}$$

- In this formula, $m(x)$ is included as $\tilde{m}(x)$.
 - Λ is UV cut-off comes from heat kernel regularization.
 - If $m(x)$ is finite, the mass dependence disappears because we need to take $\Lambda \rightarrow \infty$
 - The Λ dependence (or the regulator dependence) of the anomaly disappears after we integrate $\text{Str}[e^{\mathcal{F}}]$ over the spacetime.
- It is easy to check this anomaly is consistent with 4dim massless case.

• In this formula, $m(x)$ is included as $\tilde{m}(x)$.
 • Λ is UV cut-off comes from heat kernel regularization.
 • If $m(x)$ is finite, the mass dependence disappears because we need to take $\Lambda \rightarrow \infty$.
 • The Λ dependence (or the regulator dependence) of the anomaly disappears after we integrate $\text{Str}[e^{\mathcal{F}}]$ over the spacetime.
 • It is easy to check this anomaly is consistent with 4dim massless case.

$$\log \mathcal{J} = \frac{i}{8\pi^2} \int \alpha(x) \text{tr} [F^R \wedge F^R - F^L \wedge F^L]$$

3. Application

Introduction (5)

Derivation (10)

Application (7)


Index theorem (5)

String theory (5)

How can we apply this anomaly?

Mass means a wall for some cases!

- e.g.) Domain wall
 - Can we make domain walls by this $m(x)$?
→ Yes!
- We can make some systems with boundaries.
 - Kink, vortex and general codimension case
 - With boundary
- We also discuss about some index theorems.
 - APS index theorem
 - Callias type index theorem



Kink (1)

Mass kink for our set up

- For example, let's consider 5dim case.
- In our set up, “kink” means this mass configuration.


$$m(y) = uy \quad y = x^5 \quad u \in \mathbb{R}$$

- This “mass” diverges at $y \rightarrow \pm\infty$.
- 5dim fermions with $U(N_f)$ sym, and the mass depends on only y direction.
- The $U(1)$ anomaly is,

$$\log \mathcal{J} = \pm \frac{i}{8\pi^2} \int \alpha(x) \text{tr} [F \wedge F]$$

- Recall 4dim $U(1)_V$ anomaly, Corresponds to the sign of u .


$$\log \mathcal{J} = \frac{i}{8\pi^2} \int \alpha(x) \text{tr} [F^R \wedge F^R - F^L \wedge F^L]$$



Kink (2)


What is the meaning of the anomaly?

- 4dim Weyl fermions are localized at $y = 0$.
 - $u > 0$ corresponds to chirality + (right-handed) fermion, and $u < 0$ corresponds to chirality – (left-handed) fermion.



Domain wall fermion

- These Weyl fermions correspond to domain wall fermions.
 - But the regularization is different, so that I don't know the correspondence in detail.



Vortex

Next, we check codim-2 case.

- Vortex is 2dim topological object.

- Let us consider $2r + 2$ dim.

$$m(z) = uz \mathbf{1}_{N \times N}$$

$$z = x^{\mu=2r+1} - ix^{\mu=2r+2}$$

- $m(z)$ depends on 2 directions, and it is complex valued “mass”.

- This mass diverges at $|z| \rightarrow \infty$.

- For simplicity, we put $A_L = A_R$ in $2r + 2$ dim.

- The $U(1)_V$ anomaly is,

$$\log \mathcal{J} = -i \left(\frac{i}{2\pi} \right)^r \int \alpha(x) \text{Str} [e^F] \Big|_{2r\text{-form}}$$

- This is $2r$ dim $U(1)$ anomaly with $U(N_f)_R$ gauge field.

- If you want to get chirality – (left-handed) result, use $m(\bar{z}) = u\bar{z}$, instead.

General defects

We can apply this formula to general codimension cases.

- When we think d dim system with n dim topological defects, we get $d - n$ dim $U(1)$ anomalies.
 - If $d - n$ is odd, we get nothing because odd dim mass less fermions are anomaly-free.
 - The mass configurations for general codimension is,

$$m(x) = u \sum_{I=1}^n \Gamma^I x^I$$
$$\gamma^I = \Gamma^I \quad (n = \text{odd})$$
$$\gamma^I = \begin{pmatrix} 0 & \Gamma^I \\ \Gamma^{I\dagger} & 0 \end{pmatrix} \quad (n = \text{even})$$

- This results correspond to “tachyon condensation” in string theory.
 - We will discuss about it in section 5.


With boundary (1)

Let us make some boundaries.

- Fermions are massive = boundary

Odd dimension ($2r$ dim)

- We realize localized fermions at $[0, L]$.
- The bulk is anomaly-free.
- The anomaly is,




$$m(y) = \mu(y)1_N = \begin{cases} (m_0 + u'(y - L))1_N & (L < y) \\ m_0 1_N & (0 \leq y \leq L) \\ (m_0 + uy)1_N & (y < 0) \end{cases}$$

$$\log \mathcal{J} = i\kappa_- \int_{y=0} \alpha [\text{ch}(F)]_{2r} + i\kappa_+ \int_{y=L} \alpha [\text{ch}(F)]_{2r}$$

$$\kappa_- = \frac{1}{2} \text{sgn}(u) , \quad \kappa_+ = \frac{1}{2} \text{sgn}(u')$$

With boundary (2)

Even dimension



$$m(x) = \mu(y)g(x) = \begin{cases} u'(y-L)g(x) & (L < y) \\ 0 & (0 \leq y \leq L) \\ uyg(x) & (y < 0) \end{cases}$$

- The anomaly is,

$$\log \mathcal{J} = -i \int_{0 < y < L} \alpha [\text{ch}(F_+) - \text{ch}(F_-)]_{2r} - i \int_{y=L} \alpha[\omega]_{2r-1} + i \int_{y=0} \alpha[\omega]_{2r-1}$$

- ω is Chern-Simons form.
- Anomaly from bulk + CS

4. Index theorem

Introduction (5)

Derivation (10)

Application (7)

Index theorem (5)

String theory (5)

Index for massive Dirac op. (1)

- We will discuss index theorems for the massive Dirac operator \mathcal{D} .
 - We just consider flat spacetime.

$$\mathcal{D} = i\gamma^\mu \left\{ \partial_\mu + \begin{pmatrix} A_\mu^R & 0 \\ 0 & A_\mu^L \end{pmatrix} \right\} + \begin{pmatrix} im(x) & 0 \\ 0 & im^\dagger(x) \end{pmatrix} \quad S = \int d^d x \bar{\psi}(x) \mathcal{D} \psi(x)$$

$$\text{Ind}(\mathcal{D}) = \dim \ker(\mathcal{D}) - \dim \ker(\mathcal{D}^\dagger)$$

Chern character

- We need to define Chern character for the superconnection.
- The Chern character for massive case is,

$$\text{ch}(\mathcal{F}) = \sum_{k \geq 0} \left(\frac{i}{2\pi} \right)^{\frac{k}{2}} \text{Str} [e^{\mathcal{F}}] \Big|_{k\text{-form}} \quad \mathcal{F} \equiv \begin{pmatrix} F^R - \tilde{m}^\dagger \tilde{m} & i(D\tilde{m})^\dagger \\ iD\tilde{m} & F^L - \tilde{m}\tilde{m}^\dagger \end{pmatrix}$$

Index for massive Dirac op. (2)


- We can write the $U(1)$ anomaly by the Chern character.

$$\log \mathcal{J} = -i \left(\frac{i}{2\pi} \right)^{\frac{d}{2}} \int \alpha(x) \text{Str} [e^{\mathcal{F}}] \Big|_{d\text{-form}} = -i \int \alpha(x) \text{ch}(\mathcal{F})$$

- The index for the massive Dirac operator is, $\text{Ind}(\mathcal{D}) = \int \text{ch}(\mathcal{F})$
 - For closed manifolds, this is Atiyah-Singer index theorem.

- Let us consider $2r$ dimensional system with boundary.

- We can make the boundary with the mass.
- The index will be Atiyah-Patodi-Singer (APS) index.
- Let's check the index!




APS index theorem (1)

[’75 Atiyah-Patodi-Singer]

APS index theorem

- APS index is the index for open manifold with boundaries.
 - APS index = bulk index + η -invariant on the boundaries
 - In APS paper, they introduce APS boundary condition, which is non-local boundary condition for fermions.
- It is known that APS index is realized with local boundary condition.
 - [’17 Fukaya-Onogi-Yamaguchi] [hep-th/1710.03379]
 - If you use domain walls for boundaries, you can use local boundary condition for fermions.




- Let us consider both boundary conditions in our set up.
 - We can realize boundaries by mass; this is very similar to DW set up.

APS index theorem (2)

APS index in our set up

- The index is,
$$\text{Ind}(\mathcal{D}) = \lim_{\Lambda \rightarrow \infty} \int_{y_- < y < y_+} [\text{ch}(\mathcal{F})]_{2r} + \frac{1}{2} [\eta(H_y)]_{y=y_-}^{y=y_+}$$
 - This is the APS index theorem for the massive Dirac operators in $2r$ dim with boundaries at $y = y_{\pm}$.
 - We just consider $m(x)$ depends on only one direction y .
- Let us consider APS index for $2r$ dim system in previous section.
 - If you take $y_+ = L$ and $y_- = 0$, then you will get APS index with APS boundary condition.
 - But in this case, mass does not work because the Dirac op. is massless in $0 < y < L$.




$$\text{Ind}(\mathcal{D}|_{[0,L]}) = \int_{0 < y < L} [\text{ch}(F_+) - \text{ch}(F_-)]_{2r} - \frac{1}{2} \left[\eta(i\mathcal{D}_+^{(2r-1)}) - \eta(i\mathcal{D}_-^{(2r-1)}) \right]_{y=0}^{y=L}$$

APS index theorem (3)

APS index with mass

- Let us put $y_+ = \infty$ and $y_- = -\infty$.
 - Boundaries come from the mass.
 - APS index is,

$$\text{Ind}(\mathcal{D}) = \int_{0 < y < L} [\text{ch}(F_+) - \text{ch}(F_-)]_{2r} + \int_{y=L} [\omega]_{2r-1} - \int_{y=0} [\omega]_{2r-1}$$



- This is APS index with “physicist-friendly” boundary condition.
 - cf. [’17 Fukaya-Onogi-Yamaguchi]
 - In our set up, massive \simeq domain wall.
- To apply this form, we get a relation between eta invariant and Chern-Simons form ω .

$$\int [\omega]_{2r-1} = \frac{1}{2} \left(\eta(i\mathcal{D}_-^{(2r-1)}) - \eta(i\mathcal{D}_+^{(2r-1)}) \right) \pmod{\mathbb{Z}}$$

5. String theory

Introduction (5)

Derivation (10)

Application (7)

Index theorem (5)

String theory (5)

String theory


Let us check the relation between this anomaly and string theory.

- Consider type IIA or IIB string theory with D-branes.
 - Open strings have their ends on D-branes.
 - Excitation modes of these open strings \rightarrow Fields on D-branes
 - Open strings on D_p -branes \rightarrow QFT in $p + 1$ dim
- In some cases, excitation modes of the strings have tachyon modes.
 - Lowest excitation modes are $m^2 < 0$. (Tachyon mode)
 - Non-BPS states have tachyons.
 - These tachyonic modes are unstable. \rightarrow **Tachyon condensation**
 - See Sen's review [hep-th/9904207].




Tachyon condensation (1)

- Tachyonic modes are unstable, so the tachyons have non-zero VEV.
 - Non-trivial configuration of tachyon is also realizable.



e.g.) D -brane and anti D -brane (\bar{D} -brane) system

- Non-BPS state
- Tachyonic modes appear in $D - \bar{D}$ string.
 - The shape of tachyon potential is known. $V(T) = e^{-T^\dagger T}$
 - If tachyon configuration is trivial, the D -branes disappear.




Tachyon condensation (2)

Kink on tachyon in $D_p - \bar{D}_p$ system

- Tachyonic kinks for this system is,

$$T(x) = u \sum_{I=1}^n \Gamma^I x^I \quad \gamma^I = \begin{pmatrix} 0 & \Gamma^I \\ \Gamma^{I\dagger} & 0 \end{pmatrix}$$

- We get D_{p-n} -branes from this tachyon.
 - If D_{p-n} -branes are non-BPS, tachyons still exist on the D -branes.
 - In this case, tachyon condensation occur again.



Tachyon condensation (3)

- The superconnection is used in the context of tachyon condensation.
 - This structure comes from RR-coupling of D-branes.

cf.) ['98 Witten] [hep-th/9810188]

['99 Kennedy-Wilkins] [hep-th/9905195]

['01 Kuraus-Larsen] [hep-th/0012198]

['01 Takayanagi-Terashima-Uesugi] [hep-th/0012210]

$$S = T_{D9} \int C \wedge \text{Str } e^{2\pi\alpha' i\mathcal{F}}$$

- The tachyon configuration is given by ['98 Witten].
 - In this paper, relation between tachyon condensation and **K-theory** is discussed.
 - This tachyon configuration comes from ['64 Atiyah-Bott-Shapiro]

$$T(x) = u \sum_{I=1}^n \Gamma^I x^I$$

Relation between the anomaly and string


- This tachyon configuration is same for the mass defect in section 3!

$$T(x) = u \sum_{I=1}^n \Gamma^I x^I$$

$$m(x) = u \sum_{I=1}^n \Gamma^I x^I$$

- This anomaly can be understood from string theory.
 - Fermions are found where D -branes intersect.
 - This is similar to flavor symmetry on holographic QCD model.

(Sakai-Sugimoto model)



Conclusion

- We discussed about perturbative anomaly with **spacetime dependent mass**.
 - If value of the mass **diverge**, non-trivial contribution of the mass appears.
- The anomaly can be written by **superconnection**.

$$\log \mathcal{J} = -i \left(\frac{i}{2\pi} \right)^{\frac{d}{2}} \int \alpha(x) \text{Str} [e^{\mathcal{F}}] \Big|_{d\text{-form}} \quad \mathcal{F} \equiv \begin{pmatrix} F^R - \tilde{m}^\dagger \tilde{m} & i(D\tilde{m})^\dagger \\ iD\tilde{m} & F^L - \tilde{m}\tilde{m}^\dagger \end{pmatrix}$$

- There are some applications.
 - Kink, vortex, ...
 - With boundary
 - Index theorem