


QCD and the Spin of the Proton



Anthony W. Thomas

**High Energy Strong Interactions - YITP
Kyoto University : August 27th 2010**




Australian Government
Australian Research Council

Outline

- A reminder: the proton “spin crisis”
and the “spin problem”: $L^q + S^q + J^q = 0.5$
- Progress over the last 20 years
- The resolution of the problem
 - one-gluon-exchange
 - the pion cloud
 - input from lattice QCD
- Information from lattice QCD – and uncertainties
- Future outlook



The EMC “Spin Crisis”



Up to standard pQCD coefficients
(series in $\alpha_s(Q^2)$):

$$\int_0^1 dx g_1^p(x) = \frac{(\Delta u - \Delta d)}{12} + \frac{(\Delta u + \Delta d - 2\Delta s)}{36}$$

$$+ \frac{(\Delta u + \Delta d + \Delta s)}{9}$$

(up to QCD radiative corrections)

g_A^3 : from β decay of n

naively fraction of proton
'spin' carried by its quarks


g_A^8 : hyperon β decay

$$\Sigma_{inv} = \Sigma (Q^2 = \infty)$$

Δu = fraction of proton spin carried by u and anti-u quarks, etc..

What do we expect ?

Most quark models start with 3 quarks in the 1s-state of a confining potential: proton spin is ALL carried by its quarks: $\Sigma = 100\%$



N.B. Given low values of $m_{u,d}$ the quark motion is relativistic and lower Dirac components have spin down: $\Sigma \sim 65\%$

The Beginning

Volume 206, number 2

PHYSICS LETTERS B

19 May 1988

A MEASUREMENT OF THE SPIN ASYMMETRY AND DETERMINATION OF THE STRUCTURE FUNCTION g_1 IN DEEP INELASTIC MUON-PROTON SCATTERING

European Muon Collaboration

Aachen, CERN, Freiburg, Heidelberg, Lancaster, LAPP (Annecy), Liverpool, Marseille, Mons, Oxford, Rutherford, Sheffield, Turin, Uppsala, Warsaw, Wuppertal, Yale


J. ASHMAN ^a, B. BADELEK ^{b,1}, G. BAUM ^{c,2}, J. BEAUFAYS ^d, C.P. BEE ^e, C BENCHOUK ^f,

(93 authors)

The spin asymmetry in deep inelastic scattering of longitudinally polarised muons by longitudinally polarised protons has been measured over a large x range ($0.01 < x < 0.7$). The spin-dependent structure function $g_1(x)$ for the proton has been determined and its integral over x found to be $0.114 \pm 0.012 \pm 0.026$, in disagreement with the Ellis–Jaffe sum rule. Assuming the validity of the Bjorken sum rule, this result implies a significant negative value for the integral of g_1 for the neutron. These values for the integrals of g_1 lead to the conclusion that the total quark spin constitutes a rather small fraction of the spin of the nucleon.


$$\Sigma = 14 \pm 3 \pm 10 \% :$$

i.e. 86% of spin of p NOT carried by its quarks
and *possibly none*



Ancient History of the Spin Crisis

- EMC Spin Paper: 22 Dec 87 - 19 May 88
- Schreiber-Thomas CBM: 17 May 88 - 8 Dec 88
- Myhrer-Thomas OGE: 13 June 88 - 1 Sept 88
(neither paper could explain reduction to only 14%!)
 - Efremov-Teryaev Anomaly: 25 May 88
 - Altarelli-Ross Anomaly: 29 June 88 - 29 Sept 88
 - Carlitz, Collins, Mueller and many, many others...




Possible Role of Polarized Glue in the Proton

$$\Sigma_{\text{naïve}} \rightarrow \Sigma_{\text{naïve}} - \frac{N_f \alpha_s(Q^2) \Delta G(Q^2)}{2\pi}$$

and

QCD evolution: $\alpha_s(Q^2) \Delta G(Q^2)$ does not vanish as $Q^2 \rightarrow \infty$

and polarized gluons would resolve crisis




HOW MUCH: $\Delta G \sim +4$ no physical explanation of such a huge value ever offered !

This spurred a tremendous experimental effort


- **DIS measurements of spin structure functions of polarized p, d, ^3He (and ^6Li) at SLAC, CERN, Hermes, JLab**
- **Direct search for high- p_T hadrons as well as inclusive jet and π^0 production at Hermes, COMPASS, RHIC to directly search for effects of polarized glue in the p**
- **This effort has lasted the past 20 years, with great success**

COMPASS compared with earlier data





- very good agreement with SMC (the only other experiment at low x)
- factor 10–20 improvement of statistical errors compared to SMC


STAR result - Sarsour DNP Oct 07




- NLO pQCD describes inclusive jet cross section at RHIC
- Within GRSV framework, 2005 results constrain ΔG to less than 65% of the proton spin with 90% confidence
- Significant increase in precision in Run 2006 data provides even stronger constraints on gluon polarization



PHENIX Result: From A_{LL} to ΔG



Calc. by W.Vogelsang and M.Stratmann



“std” scenario, $\Delta G(Q^2=1 \text{ GeV}^2)=0.4$, is excluded by data on >3 sigma level



Where is the Spin of the proton?




- Modern data (Hermes, COMPASS) yields:
 $\Sigma = 0.33 \pm 0.03 \pm 0.05$

(c.f. $0.14 \pm 0.03 \pm 0.10$ originally)

- In addition, there is little or no polarized glue
 - COMPASS: $g^D_1 = 0$ to $x = 10^{-4}$
 - A_{LL} (π^0 and jets) at PHENIX & STAR: $\Delta G \sim 0$
 - Hermes, COMPASS and JLab: $\Delta G / G$ small
- Hence: axial anomaly plays at most a small role in explaining the spin crisis
- Return to alternate explanation lost in 1988 in rush to explore the anomaly

Ancient History of the Spin Crisis

- **EMC Spin Paper:** 22 Dec 87 - 19 May 88
 - **Brodsky et al. Skyrme:** 22 Feb 88 - 19 May 88
 - **Schreiber-Thomas CBM:** 17 May 88 - 8 Dec 88
 - **Myhrer-Thomas OGE:** 13 June 88 - 1 Sept 88
- (neither paper could explain reduction to only 14%!)
• **Efremov-Teryaev Anomaly:** 25 May 88
- **Altarelli-Ross Anomaly:** 29 June 88 - 29 Sept 88



OGE Hyperfine Interaction


- **Essentially every quark model needs this QCD based interaction for spectroscopy – beginning with de Rujula, Georgi, Glashow, Isgur, Karl.....**
- **N- Δ , Σ - Λ splitting etc...
(MIT bag, constituent quark model(s))**
- **As soon as this is included one must also calculate the corresponding exchange current corrections**
- **First done for magnetic moments and non-singlet axial charges by Hogaasen and Myhrer**

One-Gluon-Exchange Correction


- Further reduces the fraction of spin carried by the quarks in the bag model (naively 0.65)

- $\Sigma \rightarrow \Sigma - 3G$; with $G \sim 0.05$
 $\Sigma \rightarrow 0.65 - 0.15 = 0.5$

- Effect is to transfer quark spin to quark (relativity) and anti-quark (OGE) **orbital angular momentum**



(c)




(d)

Effect of the Pion Cloud

- Probability to find a bare N is $Z \sim 70\%$

- Biggest Fock Component is $N \pi \sim 20-25\%$ and $2/3$ of the time N spin points down



- Next biggest is $\Delta \pi \sim 5-10\%$

- To this order (i.e. including terms which yield LNA and NLNA contributions):

- Spin gets renormalized by a factor :

$$Z - \frac{1}{3} P_{N\pi} + \frac{15}{9} P_{\Delta\pi} \sim 0.75 - 0.8$$

$$\text{Hence: } \Sigma = 0.65 \rightarrow 0.49 - 0.52$$

Support for Pion Cloud Picture

- Most spectacular example is the prediction* of $\bar{d} > \bar{u}$, because of the pion cloud ($p \rightarrow n \pi^+$)


$$\int_0^1 dx [\bar{d} - \bar{u}] = 2 P_{N\pi} / 3 - P_{\Delta\pi} / 3$$
$$\epsilon 0.11 - 0.15$$

(in excellent agreement with latest data)

J.J. Kelly

* Thomas, Phys. Lett. B126 (1983) 97


- Charge distribution of the neutron
- Natural understanding of quark mass dependence of data from lattice QCD (later)



Can one add OGE and Pion Corrections?

- Prime phenomenological need for OGE interaction is the hyperfine splitting of hadron masses,
- In early days of chiral models believed some of this hyperfine splitting came from pion self-energy differences
- Maybe double counting to include correction to Σ from both pions and OGE??
- Modern understanding: *NO!* – from analysis of data in quenched (QQCD) and full QCD, from Lattice - implies 50 MeV (or less) of $m_{\Delta} - m_N$ in this way

- Lattice data (from **MILC Collaboration**) : red triangles
- Green boxes: fit evaluating σ 's on same finite grid as lattice
- Lines are exact, continuum results



Young *et al.*, hep-lat/0111041; Phys. Rev. D66 (2002) 094507

Thomas Jefferson National Accelerator Facility




Nucleon - Δ Splitting

Lattice analysis implies pions give 40 ± 20 MeV

PHYSICAL REVIEW D 66, 094507 (2002)

- Hence most of the N- Δ splitting comes from OGE – as in most quark models

- Thus the value of α_s used in the bag model calculation of the exchange current correction is more or less unchanged



and... one can add the pion and OGE corrections

Final Result for Quark Spin

$$\Sigma = (Z - P_{N\pi}/3 + 5 P_{\Delta\pi}/3) (0.65 - 3 G)$$

$$= (0.7, 0.8) \text{ times } (0.65 - 0.15) = (0.35, 0.40)$$

c.f. Experiment: $0.33 \pm 0.03 \pm 0.05$

- ALL effects, relativity and OGE and the pion cloud swap quark spin for valence orbital angular momentum and anti-quark orbital angular momentum (>60% of the spin of the proton)

Myhrer & Thomas, hep-ph/0709.4067


The Balance Sheet – fraction of total spin

	$2 L_{u+\bar{u}}$	$2 L_{d+\bar{d}}$	Σ
Non-relativistic			1.0
Relativity (e.g. Bag)	0.46	-0.11	0.65
Plus OGE	0.52	-0.02	0.50
Plus pion	0.50	0.12	0.38

At model scale: $L_u + S_u = 0.25 + 0.42 = 0.67 = J_u$
: $L_d + S_d = 0.06 - 0.22 = -0.16 = J_d$

LHPC Lattice Results


- At first glance shocking : $L^u \sim -0.1$ and $L^d \sim +0.1$
(c.f. $+0.25$ and $+0.06$ in our “resolution”)
- N.B. Disconnected terms missing
no anomaly, sea wrong : NO idea of the size of the error!



What we “see” changes with spatial resolution


>1 fm

Nucleons




0.1 — 1 fm

Constituent quarks
and glue



< 0.1 fm

“bare” quarks
and glue






In particular: $J^q(L^q)$ is not scale invariant – what scale?

- Known since mid-70s (Le Yaouanc et al., Parisi, Bell, Jaffe...) that connection between quark models and QCD must be at low scale
- This is because momentum fraction carried by quarks is monotonically decreasing with Q^2 and in models quarks carry all the momentum
- Used (for example) by Glück-Reya to model HERA data to 10^5 GeV^2 , starting with valence dominated distributions (in LO) at $\mu^2 = 0.23 \text{ GeV}^2$ (Phys Lett 359, 205 (1995))

More Modern (Confining) NJL Calculations




Cloët et al.,
Phys. Lett. B621, 246 (2005)
($\mu = 0.4 \text{ GeV}$)



Solution of the LO Evolution Equations

L^u and L^d both small and cross-over rapidly: AWT, PRL 101 (2008) 102003


- model independent !



Update

- **Recently (Bass-AWT Phys Lett B684 (2010) 216)
update to check g_A^8 and ensure that g_A^3 is correct**
- **$g_A^8 = 0.46 \pm 0.05$ (not 0.57 : 20% SU(3) breaking)**
- **This implies that value of Σ extracted from
experiment should be $0.36 \pm 0.03 \pm 0.05$**
- **To be compared with calculated $\Sigma = 0.42 \pm 0.07$
(no polarized gluon correction included)**
- **In this case we find: $J^{u,d,s} = (+0.66, -0.17, +0.01)$
at the model scale and $L^{u,d,s} = (+0.23, +0.045, +0.015)$**


NLO Evolution – input: Bass-Thomas update



LQCD Calculation: Hägler et al. (LHPC) PR D77 094502

$$\mathcal{O}_\Gamma(x) = \int \frac{d\lambda}{4\pi} e^{i\lambda x} \bar{q}\left(\frac{-\lambda n}{2}\right) \Gamma \mathcal{P} e^{-ig \int_{-\lambda/2}^{\lambda/2} d\alpha n \cdot A(\alpha n)} q\left(\frac{\lambda n}{2}\right)$$

$$\langle P', \Lambda' | \mathcal{O}_\Gamma(x) | P, \Lambda \rangle = \langle \langle \not{n} \rangle \rangle H(x, \xi, t) + \frac{n_\mu \Delta_\nu}{2m} \times \langle \langle i\sigma^{\mu\nu} \rangle \rangle E(x, \xi, t),$$




OPE)

$$\langle P' | \mathcal{O}^{\mu_1} | P \rangle = \langle \langle \gamma^{\mu_1} \rangle \rangle A_{10}(t) + \frac{i}{2m} \langle \langle \sigma^{\mu_1 \alpha} \rangle \rangle \Delta_\alpha B_{10}(t),$$

$$\langle P' | \mathcal{O}^{\{\mu_1 \mu_2\}} | P \rangle = \bar{P}^{\{\mu_1} \langle \langle \gamma^{\mu_2\} \rangle \rangle A_{20}(t) + \frac{i}{2m} \bar{P}^{\{\mu_1} \langle \langle \sigma^{\mu_2\} \alpha \rangle \rangle \Delta_\alpha B_{20}(t) + \frac{1}{m} \Delta^{\{\mu_1} \Delta^{\mu_2\}} C_{20}(t),$$

where: $\mathcal{O}_{[\gamma_5]}^{\{\mu_1 \dots \mu_n\}} = \bar{q}(0) \gamma^{\{\mu_1} [\gamma_5] i \vec{D}^{\mu_2} \dots i \vec{D}^{\mu_n\} q(0)$

$$J^q = [A_{20}(0) + B_{20}(0)] / 2$$





LQCD Needs Chiral Extrapolation and $t \rightarrow 0$


Lattice groups typically use dim-reg over large range of m_π^2

— we know this is beyond range of convergence and therefore suspect (prefer FRR)

— also extrapolate $B_{20}(t)$ *linearly* in t over $(0, 1.2)$ GeV^2



LHPC Results cont'.



Results:

$$J^u - J^d = + 0.21 \pm 0.04$$


$$L^u - L^d = - 0.42 \pm 0.04$$

$$L^u = - 0.19 \pm 0.02$$

$$L^d = + 0.23 \pm 0.02$$





(modulo disconnected terms)

- small errors rely on fit with low # parameters over huge range of t , m_π
- volume dependence taken to be small





Check Using Finite Range Regulator

As in $G_M^s(Q^2)$ study of Wang et al., PR D75 (2007) 073012



FRR Treatment of GPD moments*



Results similar to LHPC analysis
e.g. $J^u - J^d \sim 0.22$

BUT errors may be larger ...
Also, extrapolation in t using
a dipole rather than a linear
function increases J^{u-d} by 25%

*Wang & Thomas, PR D81 (2010) 114015

Implicit g_A^3 – possibly the major problem

- In order to extract L^{u-d} from lattice simulation one needs to subtract $g_A^3 / 2$ from J^{u-d}
- **BUT** we have no experience whatsoever concerning volume dependence of the value of g_A^3 implicit in the GPD moment calculation
- The calculation of g_A itself is hard enough!!




How Big Must the Lattice be?

Universally assumed $L > 4 m_\pi^{-1}$ is enough

NOT SO:
we must have
 $m_\pi (L/2 - R) \gg 1$
with $R \sim 0.8$ fm

i.e. $L > 2R + 4/m_\pi$


or $L > 6$ fm
when
 $m_\pi \sim 200$ MeV




Thomas et al.,
hep-lat/0502002

Volume Dependence of g_A Dramatic

- As well as $m_\pi \rightarrow m_\pi^{\text{phys}}$ and $t \rightarrow 0$, one has to deal with finite a and $L \rightarrow \infty$. Latter is especially problematic given that J^{u-d} includes $\Delta u - \Delta d$: RBC-UKQCD studies have shown strong volume dependence for this
 - no detailed study yet for GPDs




Yamazaki et al. [RBC + UKQCD]
arXiv: 0801.4016



Thomas, Ashley et al.,
J Phys G Conf. 9 (2005) 321

Effect on L^{u-d}


- If $J^{u-d} = 0.23 \pm 0.04$ (or 0.27 ± 0.04 – using dipole in t)
BUT g_A^3 (implicit) = (say) 1.1, then
 $L^{u-d} = -0.32 \pm 0.09$ (or -0.28 ± 0.09) at 4 GeV^2
- This agrees very well with value obtained at NLO
by Thomas: $L^{u-d} = -0.27$; also compatible with Wakamatsu



Axial Form Factor – Finite Volume Simulation

- State-of-the-art lattice calculations find axial form factor much too hard **axial radius $\sim 1/2$ physical value**
- Recent work in chiral quark model finds this in small volume – mainly from integral of

$$- \sigma(\partial^i \pi_j)$$



- This strongly supports the argument that g_A (implicit) may be considerably lower than empirical value

Hall, Thomas, Young, to be published

Summary

- Two decades of experiments have given us important new insight into spin structure of the p
- U(1) axial anomaly appears to play little role in resolving the problem
 - not as severe as in original EMC paper
- Instead, important details of the non-perturbative structure of the nucleon DO resolve the “crisis”
 - OGE hyperfine interaction
 - chiral symmetry: pion cloud
 - relativistic motion of quarks

**Ingredients of
a minimal
description of
proton structure**

Summary (cont.)

- **Important consequence for quark model:**
a large fraction of the proton spin is carried as orbital angular momentum by valence quarks and by anti-quarks in the proton
- **Effect of QCD Evolution is to:**
 - flip ordering of L^u and L^d
 - reduce the magnitude of orbital angular momentum
 - restore agreement between data, LQCD and Myhrer-Thomas explanation
- **Study of GPDs at JLab at 12 GeV may eventually provide the primary tool to verify this (also transversity?)**

Summary (cont.)

- For the time being lattice QCD offers best hope of a determination of L^u and L^d
- However:
 - L^{u+d} uncertain because of omission of disconnected terms;
 - L^{u-d} uncertain because of need to extrapolate in t and m_π over large distance *and* by need to subtract implicit value of g_A which may have significant finite volume errors
- For reasonable guess at finite volume effect L^{u-d} agrees very well with model of Myhrer and Thomas
- Model of Wakamatsu, χ QSM, is also compatible, even though value of L^{u-d} at model scale is completely different
- Much larger lattice volumes and smaller pion masses should resolve the problem

