


Nuclear Reactions

Shape, interaction, and excitation structures of nuclei ← scattering expt.
cf. Experiment by Rutherford (α scatt.)




http://www.th.phys.titech.ac.jp/~muto/lectures/QMII11/QMII11_chap21.pdf


K. Muto (TIT)

notation:


$A(a,b)B$ reaction



✓ elastic scattering




fundamental interaction
between a and A




K. Sekiguchi et al., PRC89('14)064007

$A(a,b)B$ reaction




✓ elastic scattering




fundamental interaction
between a and A


✓ inelastic scattering



excitation spectrum
of a nucleus A




$A(a,b)B$ reaction




transfer reactions

✓ transfer reaction
 (below: an example of pick-up reaction)




level scheme of ${}^{207}\text{Pb}$

✓ transfer reaction
 (below: an example of stripping reaction)



level scheme of ${}^{209}\text{Pb}$


✓ fusion reaction




- interaction between a and A
- structure of a and A

hypernucleus production reactions

✓(K^-, π^-) reaction




✓(π^+, K^+) reaction




excitation spectrum
of a hypernucleus A_Λ

$^{12}\text{C} (\pi^+, K^+) {}^{12}\Lambda\text{C}$ reaction



O. Hashimoto and H. Tamura,
Prog. in Part. and Nucl. Phys. 57 ('06)564


✓(e,e'K⁺) reaction



S.N. Nakamura et al.,
PRL110('13)012502


T. Gogami,
Ph.D. Thesis (Tohoku U.)
2014

$^{12}\text{C}(\text{e},\text{e}'\text{K}^+) \ ^{12}_{\Lambda}\text{B}$



L. Tang et al., PRC90('14)034320


Cross sections



incident beam

flux = the number of particles crossing unit area per unit time

$$j = \rho_P \cdot v$$




event rate (the number of event per unit time per target nucleus)
: proportional to the incident flux

$$R = N_T \cdot \sigma \cdot j$$

← cross section

Cross sections



event rate (the number of event per unit time per target nucleus)
: proportional to the incident flux

$$\longrightarrow R = N_T \cdot \sigma \cdot j$$


← cross section

differential cross sections (angular distribution)

$$dR(\theta, \phi) = N_T \cdot \frac{d\sigma}{d\Omega} \cdot j \cdot d\Omega,$$
$$\sigma = \int d\Omega \frac{d\sigma}{d\Omega}$$

units: 1 barn = $10^{-24} \text{ cm}^2 = 100 \text{ fm}^2$ (1 mb = $10^{-3} \text{ b} = 0.1 \text{ fm}^2$)

Cross sections (experiments)



the target thickness

$$dR(\theta, \phi) = N_T \cdot \frac{d\sigma}{d\Omega} \cdot j \cdot d\Omega.$$

beam intensity: $I = j \cdot S$

the number of target nucleus: $N_T = S \cdot t \cdot \rho_T$





$$dR(\theta, \phi) = I \cdot \frac{d\sigma}{d\Omega} \cdot t \rho_T \cdot d\Omega \cdot \epsilon$$



detection efficiency

Cross sections (theory)

$A(a,b)B$ reaction





center of mass frame





$$\frac{d\sigma}{d\Omega} = \frac{R}{j_{in}}$$

Cross sections

✓ laboratory frame



✓ center of mass frame



□ transformation ← energy and momentum conservations

$$\tan \theta_{\text{lab}} = \frac{\sin \theta_{\text{cm}}}{\gamma + \cos \theta_{\text{cm}}}, \quad d\Omega_{\text{lab}} = \frac{|1 + \gamma \cos \theta_{\text{cm}}|}{(1 + \gamma^2 + 2\gamma \cos \theta_{\text{cm}})^{3/2}} d\Omega_{\text{cm}}$$

$$E_{\text{cm}} = \frac{M_A}{M_a + M_A} E_{\text{lab}}, \quad \gamma = \sqrt{\frac{M_a M_b}{M_A M_B} \frac{E_{\text{cm}}}{E_{\text{cm}} + Q}}$$


Born approximation

$$\psi_i(r) = e^{i\mathbf{p}_i \cdot \mathbf{r}/\hbar}$$



$$V(r)$$

$$\psi_f(r) = e^{i\mathbf{p}_f \cdot \mathbf{r}/\hbar}$$



$$\left(-\frac{\hbar^2}{2\mu} \nabla^2 + \underline{V(r)} - E \right) \psi(r) = 0$$

perturbation

transition rate for elastic scattering:

$$\begin{aligned} W_{fi} &= \frac{2\pi}{\hbar} \int \frac{d\mathbf{p}_f}{(2\pi\hbar)^3} |\langle \psi_f | V | \psi_i \rangle|^2 \delta(E_f - E_i) \\ &= \frac{\mu p_i}{4\pi^2 \hbar^4} \int d\Omega |\tilde{V}(\mathbf{q})|^2 \end{aligned}$$

$$\tilde{V}(\mathbf{q}) = \int dr e^{i(\mathbf{p}_i - \mathbf{p}_f) \cdot \mathbf{r}/\hbar} V(r) \equiv \int dr e^{-i\mathbf{q} \cdot \mathbf{r}} V(r)$$


Born approximation

$$\psi_i(r) = e^{ip_i \cdot r / \hbar}$$



$$V(r)$$

$$\psi_f(r) = e^{ip_f \cdot r / \hbar}$$



$$W_{fi} = \frac{\mu p_i}{4\pi^2 \hbar^4} \int d\Omega |\tilde{V}(\mathbf{q})|^2$$


$$\tilde{V}(\mathbf{q}) = \int dr e^{i(\mathbf{p}_i - \mathbf{p}_f) \cdot \mathbf{r} / \hbar} V(r) \equiv \int dr e^{-i\mathbf{q} \cdot \mathbf{r}} V(r)$$

incident flux: $j_{\text{inc}} = \rho_i v = p_i / \mu$

↷

$$\sigma = \frac{W_{fi}}{j_{\text{inc}}} = \int d\Omega \frac{\mu^2}{4\pi^2 \hbar^4} |\tilde{V}(\mathbf{q})|^2$$

$$= \frac{d\sigma}{d\Omega}$$




$$q\hbar = 2p_i \sin \frac{\theta}{2}$$

momentum
transfer



Electron scattering



$$V(r) = -e^2 \int d\mathbf{r}' \frac{\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}$$


$$\begin{aligned} \frac{d\sigma}{d\Omega} &= \frac{Z_P^2 e^4}{(4E \sin^2 \theta/2)^2} |F(\mathbf{q})|^2 \\ &= \left(\frac{d\sigma_{\text{Ruth}}}{d\Omega} \right) |F(\mathbf{q})|^2 \end{aligned}$$

Form factor


$$F(\mathbf{q}) = \int e^{-i\mathbf{q} \cdot \mathbf{r}} \rho(\mathbf{r}) d\mathbf{r}$$

* relativistic correction:


$$\begin{aligned} \frac{d\sigma_{\text{Ruth}}}{d\Omega} &\rightarrow \frac{d\sigma_{\text{Mott}}}{d\Omega} \\ &= \frac{d\sigma_{\text{Ruth}}}{d\Omega} \cdot \left(1 - \frac{v^2}{c^2} \sin^2 \frac{\theta}{2} \right) \\ &\sim \frac{d\sigma_{\text{Ruth}}}{d\Omega} \cdot \cos^2 \frac{\theta}{2} \quad (v \rightarrow c) \end{aligned}$$



cf. electron scattering off unstable nuclei (SCRIT)




K. Tsukada et al.,
PRL118, 262501 (2017)



proton radius puzzle

$$\begin{aligned} F(q) &= \int e^{-i\mathbf{q}\cdot\mathbf{r}} \rho(\mathbf{r}) d\mathbf{r} \\ &\sim \int \left(1 - i\mathbf{q} \cdot \mathbf{r} - \frac{(qr)^2}{2} \cos^2 \theta + \dots \right) \rho(\mathbf{r}) d\mathbf{r} \\ &\sim Z \left(1 - \frac{q^2}{6} \langle r^2 \rangle + \dots \right) \end{aligned}$$




Distorted Wave Born approximation (DWBA)

$$\left(-\frac{\hbar^2}{2\mu} \nabla^2 + \underline{V(r) - E} \right) \psi(r) = 0$$

perturbation

→ $\left(-\frac{\hbar^2}{2\mu} \nabla^2 + V_0(r) + \underline{V(r) - V_0(r) - E} \right) \psi(r) = 0$

perturbation





- ✓ inelastic scattering
- ✓ transfer reactions

Optical model

Reaction processes

- Elastic scatt.
- Inelastic scatt.
- Transfer reaction
- Compound nucleus formation (fusion)



Loss of incident flux
(absorption)


Optical potential

$$V_{\text{opt}}(r) = V(r) - iW(r) \quad (W > 0)$$

$$\rightarrow \nabla \cdot j = \dots = -\frac{2}{\hbar} W |\psi|^2$$


(note) Gauss's theorem

$$\int_S \mathbf{j} \cdot \mathbf{n} dS = \int_V \nabla \cdot \mathbf{j} dV$$



$$\left(-\frac{\hbar^2}{2\mu} \nabla^2 + \frac{Z_P Z_T e^2}{r} + V_{\text{opt}}(r) - E \right) \psi(r) = 0$$


Woods-Saxon + volume & surface
imaginary parts



H. Sakaguchi et al.,
PRC26 (1982) 944

Appendix: DWBA in ocean acoustics

Fishfinder



(backward) scattering of (ultra-)sonic waves due to fish etc.

$$dR(\theta, \phi) = N_T \cdot \frac{d\sigma}{d\Omega} \cdot j \cdot d\Omega$$




$$N_T = \frac{\frac{dR}{d\Omega}}{j \cdot \frac{d\sigma}{d\Omega}}$$

one can know the number of fish N_T if one knows the differential cross sections

Use of the distorted wave Born approximation to predict scattering by inhomogeneous objects: Application to squid


Benjamin A. Jones,^{a)} Andone C. Lavery, and Timothy K. Stanton


Department of Applied Ocean Physics and Engineering, Woods Hole Oceanographic Institution,
Woods Hole, Massachusetts 02543-1053



Modeling of squid


DWBA: local wave number
inside a squid





Krill (オキアミ)


measurement



Impulse approximation

example: ${}^A_Z(K^-, \pi^-) {}^A_{\Lambda} Z$ reaction

- ✓ high energy
- ✓ single scattering approximation




$$T_{fi} \sim \left\langle \psi_{\pi^-} \left| \left\langle \Psi_{\Lambda}^A \left| \sum_j v_{eff}(j) \right| \Psi_{AZ} \right\rangle \right| \psi_{K^-} \right\rangle$$

$$\frac{d\sigma}{d\Omega} \sim \underbrace{\alpha_{\text{kin}} \left(\frac{d\sigma}{d\Omega} \right)_{K^- n \rightarrow \pi^- \Lambda}}_{\text{kinematical factor}} N_{\text{eff}}(\theta; i \rightarrow f)$$

kinematical
factor elementary process



$$N_{\text{eff}}(\theta; i \rightarrow f) \sim \left| \int dr \psi_{\pi^-}^*(r) \varphi_{j_\Lambda l_\Lambda m_\Lambda}^{(\Lambda)*}(r) \varphi_{j_n l_n m_n}^{(n)}(r) \psi_{K^-}(r) \right|^2$$

- Plane wave impulse approximation (PWIA)
- Distorted wave impulse approximation (DWIA)



O. Hashimoto and H. Tamura,
Prog. in Part. and Nucl. Phys. 57 ('06)564

T. Motoba et al., PRC38('88)1322




$$m_n + m_K = 1432 \text{ MeV} \quad Q > 0$$

$$m_\pi + m_\Lambda = 1255.3 \text{ MeV} \quad Q < 0$$

$$m_\pi + m_n = 1079.2 \text{ MeV} \quad Q < 0$$


$$m_K + m_\Lambda = 1609.4 \text{ MeV} \quad Q < 0$$



O. Hashimoto and H. Tamura,
Prog. in Part. and Nucl. Phys. 57 ('06)564




T. Motoba et al., PRC38('88)1322

relation between q and Δl



$$l \sim kb \text{ (classically)}$$

$$\rightarrow \Delta l \sim b(p' - p) = bq$$



O. Hashimoto and H. Tamura,
Prog. in Part. and Nucl. Phys. 57 ('06)564



T. Motoba et al., PRC38('88)1322

$$\Delta l \sim b(p' - p) = bq$$

Absorption cross sections

Reaction processes

- Elastic scatt.
- Inelastic scatt.
- Transfer reaction
- Compound nucleus formation (fusion)



Loss of incident flux
(absorption)


reaction cross sections

total scattering cross section - elastic cross section

$$\sigma_R = \sigma_{\text{tot}} - \sigma_{\text{el}}$$


- fusion
- inelastic
- transfer

Interaction cross sections and halo nuclei



interaction cross section σ_I
= cross section for the change
of Z a/o N in the incident nucleus


transmission method




$$\sigma_R = -\frac{1}{t} \ln \left(\frac{N_{\text{out}}}{N_{\text{in}}} \right)$$

$$t = \rho_T \cdot d \cdot \epsilon$$

Interaction cross sections and halo nuclei





interaction cross section σ_I
= cross section for the change
of Z a/o N in the incident nucleus



$$\sigma_I \sim \pi [R_I(P) + R_I(T)]^2 \longrightarrow R_I(P)$$


Discovery of halo nuclei



I. Tanihata, T. Kobayashi, O. Hashimoto
et al., PRL55('85)2676; PLB206('88)592




Reaction cross sections




Glauber theory (optical limit approximation : OLA)

$$\sigma_R \sim 2\pi \int_0^\infty b db \left[1 - \exp \left(-\sigma_{NN} \int d^2 s \rho_P^{(z)}(s) \rho_T^{(z)}(s - b) \right) \right]$$

- straight-line trajectory (high energy scattering)
- adiabatic approximation
- simplified treatment for multiple scattering: $(1 - x)^N \rightarrow e^{-Nx}$



Density distribution which explains the experimental σ_R



M. Fukuda et al., PLB268('91)339

$$\sigma_R \sim 2\pi \int_0^\infty bdb \left[1 - \exp \left(-\sigma_{NN} \int d^2s \rho_P^{(z)}(s) \rho_T^{(z)}(s-b) \right) \right]$$