

Dispersion Relation of Charmonia above T_c

Masakiyo Kitazawa (Osaka U.)

Ikeda, Asakawa, MK, PRD95 (2017) 014504




Charm Quarks in HIC


Impurity of QGP
= unique experimental probe

- production only in first stage
- small abundance

- J/p suppression
- transport property
- heavy-quark potential




ALI-DER-65274



Figs. from arXiv:1506.03981

Charmonia above T_c


- Property of charmonia at rest
 - Melting temperature
 - Mass shift?



Charmonia above T_c

- Property of charmonia at rest


- Melting temperature
- Mass shift?



- Property of moving charmonia

- Dispersion relation
- residue
- decay rate

In heavy-ion collisions,
charmonia are typically moving!



Charmonia above T_c

- Property of charmonia at rest

- Melting temperature



- Mass shift?

- Property of moving charmonia


- Dispersion relation

- residue

- decay rate



**Let's study
on the lattice**




In heavy-ion collisions,
charmonia are typically moving!

Nonzero-p Spectral Func.

In vacuum :Lorentz symmetry

- Tensor structure (V) $\rho_{\mu\nu}(\omega, \vec{p}) = \left(\frac{p_\mu p_\nu}{p^2} - g_{\mu\nu} \right) \rho_V(p)$
- Bound-state pole $\sim Z \delta(\omega^2 - E(\vec{p})^2) = \frac{Z}{2E(\vec{p})} \delta(\omega - 2E(\vec{p}))$
- Dispersion relation $E(\vec{p}) = \sqrt{m^2 + \vec{p}^2}$




Nonzero-p Spectral Func.

In vacuum : Lorentz symmetry

- Tensor structure (V) $\rho_{\mu\nu}(\omega, \vec{p}) = \left(\frac{p_\mu p_\nu}{p^2} - g_{\mu\nu} \right) \rho_V(p)$
- Bound-state pole $\sim Z \delta(\omega^2 - E(\vec{p})^2) = \frac{Z}{2E(\vec{p})} \delta(\omega - 2E(\vec{p}))$
- Dispersion relation $E(\vec{p}) = \sqrt{m^2 + \vec{p}^2}$


In medium

- Transverse and longitudinal splitting
 $\rho_{\mu\nu}(\omega, \vec{p}) = \rho_T(\omega, \vec{p}) \Lambda_T + \rho_L(\omega, \vec{p}) \Lambda_L$
- Dispersion relation can be modified
- Z no longer be a constant



Maximum Entropy Method


Lattice data



$$G(\tau) = \int_0^\infty d\omega \frac{\cosh(1/2T - \tau)\omega)}{\sinh(\omega/2T)} \rho(\omega)$$


“ill-posed problem”

Spectral Function



Maximum Entropy Method

Lattice data



Bayes
theorem




Prior probability

- Shannon-Jaynes entropy
- default model $m(\omega)$




Probability
of $\rho(\omega)$
 $P[\rho(\omega), \alpha]$

Spectral Function



Maximum Entropy Method

Lattice data




Bayes
theorem



Prior probability


- Shannon-Jaynes entropy
- default model $m(\omega)$



Probability of $\rho(\omega)$

$$P[\rho(\omega), \alpha]$$

Spectral Function



expectation value

$$\langle \rho(\omega) \rangle_P$$


$$\langle \mathcal{O} \rangle_P = \int d\alpha \int [d\rho] P[\rho, \alpha] \mathcal{O}$$

Error in MEM

MEM error = variance in $P[\rho(\omega), \alpha]$ space

$$W = \int d\omega f(\omega) \rho(\omega)$$

- exp. val.: $\langle W \rangle_P = \int d\omega f(\omega) \langle \rho(\omega) \rangle_P$
- error: $\Delta W = \sqrt{(W - \langle W \rangle_P)^2}$




Error in MEM

MEM error = variance in $P[\rho(\omega), \alpha]$ space

$$W = \int d\omega f(\omega) \rho(\omega)$$

- exp. val.: $\langle W \rangle_P = \int d\omega f(\omega) \langle \rho(\omega) \rangle_P$
- error: $\Delta W = \sqrt{\langle (W - \langle W \rangle_P)^2 \rangle}$

$$f(\omega') = \delta(\omega' - \omega) \Rightarrow \begin{aligned} \rho_{\text{out}}(\omega) &= \langle \rho(\omega) \rangle_P \\ \Delta \rho_{\text{out}}(\omega) &= \sqrt{\langle (\delta \rho(\omega))^2 \rangle_P} \end{aligned} \Rightarrow \text{typically huge}$$



Error in MEM

MEM error = variance in $P[\rho(\omega), \alpha]$ space

$$W = \int d\omega f(\omega) \rho(\omega)$$

- exp. val.: $\langle W \rangle_P = \int d\omega f(\omega) \langle \rho(\omega) \rangle_P$
- error: $\Delta W = \sqrt{(W - \langle W \rangle_P)^2}$

$$f(\omega') = \delta(\omega' - \omega) \Rightarrow \begin{aligned} \rho_{\text{out}}(\omega) &= \langle \rho(\omega) \rangle_P \\ \Delta \rho_{\text{out}}(\omega) &= \sqrt{\langle (\delta \rho(\omega))^2 \rangle_P} \end{aligned} \Rightarrow \text{typically huge}$$


NOTE

- SPC obtained by MEM is just an image. No robust meaning.
- MEM error is more conservative than statistical one.

Defining Peak Position

Center of weight in a range $[\omega_{\min} : \omega_{\max}]$

$$\bar{E} = \frac{\int_{\omega_{\min}}^{\omega_{\max}} d\omega \omega (\rho(\omega) / \omega^2)}{\int_{\omega_{\min}}^{\omega_{\max}} d\omega \rho(\omega) / \omega^2}$$




- ❑ Represent peak position for a sufficiently sharp peak
- ❑ **Error analysis in MEM is possible!**
- ❑ $[\omega_{\min}, \omega_{\max}]$ dependence has to be checked

Defining Peak Position

Center of weight in a range $[\omega_{\min} : \omega_{\max}]$

$$\bar{E} = \frac{\int_{\omega_{\min}}^{\omega_{\max}} d\omega \omega (\rho(\omega) / \omega^2)}{\int_{\omega_{\min}}^{\omega_{\max}} d\omega \rho(\omega) / \omega^2}$$



- ❑ Represent peak position for a sufficiently sharp peak
- ❑ **Error analysis in MEM is possible!**
- ❑ $[\omega_{\min}, \omega_{\max}]$ dependence has to be checked

- ❑ Residue: $\bar{Z} = \int_{\omega_1}^{\omega_2} d\omega 2\omega \rho(\omega)$

Lattice Setup

- quenched simulation
- Wilson fermion / gauge
- anisotropic lattice ($a_\sigma/a_\tau=4$)

$$\beta = 7.0, \gamma_F = 3.476, \kappa_\sigma = 0.8282$$

$$a_\sigma = 0.00975[\text{fm}], a_\sigma/a_\tau = 4$$

Asakawa, Hatsuda, 2004

BlueGene/Q@KEK


fermion part: Iroiro++

N_τ	T/T_c	N_σ	$L_\sigma[\text{fm}]$	N_{conf}
40	1.86	64	2.5	500x8
46	1.62	64	2.5	500x8
50	1.49	64	2.5	500x8
96	0.78	64	2.5	500x8

8 measurements
on each conf.

- Large spatial volume \rightarrow high momentum resolution
- Large N_τ / high statistics \rightarrow high MEM precision

Spectral Func. @ $T=0.78T_c$





- ρ_T and ρ_L channels degenerate
- although correlators are different

$$\vec{p} = (p, 0, 0)$$

$$G_L = \frac{\omega^2 - p^2}{\omega^2} G_{11}$$

$$G_T = G_{22} = G_{33}$$

Spectral Func. @ $T=0.78T_c$



- ρ_T and ρ_L channels degenerate
- although correlators are different


$$\vec{p} = (p, 0, 0)$$

$$G_L = \frac{\omega^2 - p^2}{\omega^2} G_{11}$$


$$G_T = G_{22} = G_{33}$$

Spectral Function @ $p=0$


Vector channel



PS channel




Spectral Func. @ $T=1.62T_c$



- ρ_T and ρ_L channels seem to degenerate.
- Peak exists for all momentum.

Dispersion Relation

Energy interval $[\omega_{\min}, \omega_{\max}]$ for disp. rel.





$\omega_{\min} = 3\text{GeV}$

ω_{\max} : first minimum

Dispersion Relation

● PS: $0.78T_c$
■ PS: $1.49T_c$
◆ PS: $1.62T_c$





- Clear mass enhancement in medium.
- Dispersion relation is consistent with the Lorentz covariant form even at $T=1.62T_c$.

mass $E(p=0)$

T/T_c	0.78	1.49	1.62
J/ψ	3.24(6)	4.30(16)	4.47(16)
η_c	3.19(5)	4.24(31)	4.49(48)

[GeV]



Dependence on $[\omega_{\min} : \omega_{\max}]$



- ω_{\max} dependence is well suppressed.
- No ω_{\min} dependence for $\omega_{\min} < 3$ GeV.

Test: N_t Dependence


Special thanks to A. Rothkopf



- Peak position does not shift with the change of N_t

correlator	$G(\tau, 0, T)$	$G^{\text{rec}}(\tau, 0, 2T; T)$
\bar{m}	3.24(6)	3.40(90)

Res due



- No p dependence of Z even for $T=1.62T_c$
- No T/L splitting in vector channel

Summary

- We analyzed the **peak positions in SPC with MEM error** by **defining** them in terms of the **center of weight**.
- Charmonia have **significant mass enhancement**.
- Dispersion relations are consistent with **Lorentz covariant form even at $T=1.62T_c$** .

Future Work

- much finer p resolution
- m_q dependence
- comparison with potential models and etc.