



# Jet-driven supernovae in the multi-messenger era

Philipp Mösta


GRAPPA / University of  
Amsterdam  
p.moesta@uva.nl

YITP, Oct 1, 2019







**(Binary) black holes**  
accretion disks  
EM counterparts



**Core-collapse  
supernovae**  
neutrinos  
turbulence



**Extreme core-collapse**  
hyperenergetic  
superluminous  
IGRBs



**Binary neutron stars**  
gravitational waves  
EM counterparts  
sGRBs

**Magnetic fields in  
high-energy astro**

# New era of transient science

- Current (PTF, DeCAM, ASAS-SN) and upcoming wide-field time domain astronomy (ZTF, LSST, ...) -> wealth of data
- adv LIGO / gravitational waves detected
- Computational tools at dawn of new exascale era



Image: PTF/ZTF/COO




Image: LSST

# New era of transient science

- Current (PTF, DeCAM, ASAS-SN) and upcoming wide-field time domain astronomy (ZTF, LSST, ...) -> wealth of data
- adv LIGO / gravitational waves detected
- Computational tools at dawn of new exascale era

Transformative years ahead for our understanding of these events



Image: PTF/ZTF/COO




Image: LSST

# Astrophysics of core-collapse supernovae



M82/Chandra/NASA


Galaxy evolution/feedback



# Observing core-collapse supernovae

- EM waves (optical/UV/X/Gamma):  
secondary information,  
late-time probes of engine


Red Supergiant  
Betelgeuse  
D ~200 pc  
HST



# Observing core-collapse supernovae

- Neutrinos
- EM waves (optical/UV/X/Gamma):  
secondary information,  
late-time probes of engine


Red Supergiant  
Betelgeuse  
D ~200 pc  
HST




# Observing core-collapse supernovae

- Gravitational waves
- Neutrinos
- EM waves (optical/UV/X/Gamma):  
secondary information,  
late-time probes of engine

Red Supergiant  
Betelgeuse  
D ~200 pc  
HST




# Core collapse basics



Nuclear equation of state stiffens at nuclear density

Inner core ( $\sim 0.5 M_{\odot}$ )  
-> protoneutron star + shockwave


# Core collapse basics



Nuclear equation of state stiffens at nuclear density

Inner core ( $\sim 0.5 M_{\odot}$ )  
-> protoneutron star + shockwave


Reviews:  
Bethe'90  
Janka+'12



Outer core accretes onto shock & protoneutron star with  $O(1) M_{\odot}/s$

Shock stalls at  $\sim 100\text{ km}$


# Core collapse basics



Nuclear equation of state stiffens at nuclear density

Inner core ( $\sim 0.5 M_{\odot}$ )  
-> protoneutron star + shockwave


Reviews:  
Bethe'90  
Janka+'12



Core-collapse  
supernova problem:  
How to revive the  
shockwave?


# Core collapse basics

Neutrino mechanism



# Core collapse basics

Neutrino mechanism



# Core collapse basics


Neutrino mechanism

$L_\nu$

Theory incomplete!

accretion


shock



# Core collapse basics


3D Volume  
Visualization of

**Entropy**



Roberts+16

# Extreme Supernovae and GRBs






- 11 long GRB - core-collapse supernova associations.
- All GRB-SNe are stripped envelope, show outflows  $v \sim 0.1c$
- But not all stripped-envelope supernovae come with GRBs
- Trace low metallicity environments
- Some SLSNe share same characteristics

# The engine(s) driving these transients

Superluminous

Hyperenergetic SNe

IGRBs




**Engine?**


**Progenitor**


```
graph TD; A([Progenitor]) --> B([Engine?]);
```

The diagram consists of two cyan ovals with dark blue outlines. The top oval is smaller and contains the word 'Progenitor'. A white arrow points from the bottom of this oval to the top of a larger oval below it, which contains the text 'Engine?'.

**Engine?**








**Establish mapping**

**progenitor -> engine -> observations**


# Core collapse basics



Nuclear equation of state stiffens at nuclear density

Inner core ( $\sim 0.5 M_{\odot}$ )  
-> protoneutron star + shockwave

Reviews:  
Bethe'90  
Janka+'12



## Engine formation?

# Protomagnetar powered explosions



**Rapid Rotation + B-field amplification**

**Results in ms-period proto-magnetar**

**2D: Energetic bipolar explosions**  
Energy in rotation up to  $10^{52}$  erg

# MHD-supernova vs collapsar

MHD-supernova / magnetorotational  
supernova: outflows driven by  
protomagnetar



# MHD-supernova vs collapsar

**MHD-supernova / magnetorotational supernova:** outflows driven by protomagnetar

**Collapsar:** Compact object (likely black hole) + accretion disk -> outflows driven by disk wind



# MHD-supernova vs collapsar

**MHD-supernova / magnetorotational supernova:** outflows driven by protomagnetar

**Collapsar:** Compact object (likely black hole) + accretion disk -> outflows driven by disk wind

**Two different engines with different signatures!**



# MHD-supernova vs collapsar

**MHD-supernova / magnetorotational supernova:** outflows driven by protomagnetar

**Collapsar:** Compact object (likely black hole) + accretion disk -> outflows driven by disk wind

**Two different engines with different signatures!**

Could be realized in same progenitor system but at different times



# A multiphysics challenge

Magneto-Hydrodynamics

→ Gas/plasma dynamics

# A multiphysics challenge


Magneto-Hydrodynamics

→ Gas/plasma dynamics

General Relativity

→ Gravity

# Dynamical gravity / Numerical Relativity



$$G^{\mu\nu} = \frac{8\pi G}{c^4} T^{\mu\nu}$$

- 12 first-order hyperbolic *evolution* equations
- 4 elliptic *constraint* equations
- 4 coordinate gauge degrees of freedom:  $\alpha$ ,  $\beta^i$

# A multiphysics challenge

Magneto-Hydrodynamics

→ Gas/plasma dynamics

General Relativity

→ Gravity

Nuclear and Neutrino Physics

→ Nuclear EOS, nuclear reactions &  $\nu$  interactions

# A multiphysics challenge

Magneto-Hydrodynamics

→ Gas/plasma dynamics

General Relativity

→ Gravity

Nuclear and Neutrino Physics

→ Nuclear EOS, nuclear reactions &  $\nu$  interactions

Boltzmann Transport Theory

→ Neutrino transport

# A multiphysics challenge

Fully coupled!

Magneto-Hydrodynamics

→ Gas/plasma dynamics

General Relativity

→ Gravity

Nuclear and Neutrino Physics


→ Nuclear EOS, nuclear reactions &  $\nu$  interactions

Boltzmann Transport Theory

→ Neutrino transport

**All four forces!**

# A multiphysics challenge



**All four forces!**

**Additional Complication: Core-Collapse Supernovae are 3D**

- rotation
- fluid and MHD instabilities, multi-D structure, spatial scales


**Need 21st century tools:**

- cutting edge numerical algorithms
- sophisticated open-source software infrastructure
- peta/exa scale computers



How do we form magnetars?

# One proposed channel: MRI + dynamo



# MRI Basics


- Weak field instability
- Requires negative angular velocity gradient
- Can build up magnetic field exponentially fast
- Extensively researched in accretion disks: ability to modulate angular momentum transport and grow large scale field

# What's the situation in core-collapse?

Stability criterion:

$$-8\Omega^2 < \omega_{\text{BV}}^2 + r \frac{d\Omega^2}{dr} < 0$$


[Balbus&Hawley 91,98, Akiyama+03, Obergaulinger+09]



# Global 3D MHD turbulence simulations

- 10 billion grid points (Millenium simulation used 10 billion particles)
- 130 thousand cores on Blue Waters
- 2 weeks wall time
- 60 million compute hours
- 10000 more expensive than any previous simulations

Do MRI and dynamo build up dynamically relevant global field?



PM+ 15 Nature

**BLUE WATERS**  
SUSTAINED PETASCALE COMPUTING



# 3D magnetic field structure


$dx=500m$


$dx=200m$

$dx=100m$

$dx=50m$

$t = 0.00 \text{ ms}$





PM+15 Nature




# R-process nucleosynthesis in magnetar-driven explosions

# 3D explosions dynamics very different!

PM+ 14


← 2000 km →

$t = -3.00$  ms



← 2000 km →

$t = -3.00$  ms




Octant Symmetry (no odd modes)  
identical to 2D


Full 3D

# What's going on here?

with Sherwood Richers (Caltech)



PM, Richers+ 14



- $m=1$  spiral instability
- consistent with MHD kink instability; should hold independent of initial B-field strength

$$\tau_{\text{fgm}} \approx \frac{4a\sqrt{\pi\rho}}{B_{\text{tor}}} \approx 1\text{ms}$$

$$\lambda_{\text{fgm}} \approx \frac{4\pi a B_z}{B_{\text{tor}}} \approx 5\text{km}$$


3D Volume  
Visualization of

$t = -3.00 \text{ ms}$


# Entropy

PM+ 14

# Implications for long Gamma-Ray Bursts



dual-lobe 'slow'  
explosion




**Continued accretion ->  
Black hole engine possible!**

# Neutron-rich nucleosynthesis in supernovae


## Creating the heaviest elements

Jet-driven explosions proposed as site for r-process

- Low electron fraction
- Medium entropy
- Low density
- High temperature





# Making the heaviest elements




PM+ 18  
 Halevi, PM+ 18

# R-process in jet-driven supernovae




Winteler+12

Nobuya's talk  
yesterday!



Nishimura+15



# R-process in jet-driven supernovae




# R-process nucleosynthesis in supernovae

$B = 10^{13} \text{ G}$


See Goni's talk on  
Thursday morning!




# R-process in jet-driven supernovae




# R-process in jet-driven supernovae



# R-process in jet-driven supernovae





# R-process nucleosynthesis in supernovae




PM, Roberts, Halevi+ 18

$B = 10^{12} \text{ G full 3D}$



# R-process nucleosynthesis in supernovae



# From simulations to observations

## Observations:

- new transients classes and subclasses
- need detailed predictions to constrain engines

## Simulations

- initial 3D simulations open up diverse outcomes
- magnetic fields crucial component for signatures

**Need mapping:**

**progenitor** -> **engine** -> **observations**

# From simulations to observations

## State of the art now:

Detailed simulations  
full physics  
0.1-1s  
~10000km

engine formation/dynamics  
gravitational waves  
nucleosynthesis

# From simulations to observations

## State of the art now:

Detailed simulations  
full physics  
0.1-1s  
~10000km

# From simulations to observations

## State of the art now:

Detailed simulations  
full physics  
0.1-1s  
~10000km

## Current frontier:

- 1) engine model from full-physics simulations
- 2) simplified simulations with engine model to shock breakout

explosion geometry  
explosion energy  
nucleosynthesis  
basic engine model

# From simulations to observations

## State of the art now:

Detailed simulations  
full physics  
0.1-1s  
~10000km


## Current frontier:

- 1) Engine model from full-physics simulations
- 2) Simplified simulations with engine model to shock breakout

Full 3D, full physics



Full star



# From simulations to observations

## State of the art now:

Detailed simulations  
full physics  
0.1-1s  
~10000km

## Next five years:

full-scale simulations  
full physics  
shock breakout

## Current frontier:

- 1) Engine model from full-physics simulations
- 2) Simplified simulations with engine model to shock breakout

detailed light curves  
detailed spectra  
connect observations and engines  
map progenitor params

# Summary

R-process from jet-driven supernovae in explosion phase only  
for strong seed magnetization

Need to look at late time behavior when neutrino luminosity  
has gone down

Follow evolution to remnant formation / shock breakout

**Mapping:**

**progenitor** -> **engine** -> **observations**

# Discussion points

What is going to help us disentangle r-process different channels?

Does it make sense to focus on better theoretical modeling right now?

**Mapping:**

**progenitor** -> **engine** -> **observations**