

# Heavy exotics and kinematical effects

Feng-Kun Guo

Institute of Theoretical Physics, Chinese Academy of Sciences

*New Frontiers in QCD 2018 — Confinement, Phase Transition, Hadrons and  
Hadron Interactions, YITP, May 28 – June 29, 2018*

---

Based on:

M. Bayar, F. Aceti, FKG, E. Oset, PRD94(2016)074039; (initiated here!)

FKG, U.-G. Meißner, W. Wang, Z. Yang, PRD92(2015)071502(R);

J.-J. Xie, FKG, PLB774(2017)108;




M. Albaladejo, FKG, C. Hidalgo-Duque, J. Nieves, PLB755(2016)337

# The search of resonances




In practice, resonance hunting is normally the search of peaks.

Some famous peaks:

$Z_b(10610)$  and  $Z_b(10650)$




$Z_c(3900)$ ,  $X(5568)$ ,  $P_c(4380, 4450)$



# Resonances are not always peaks

However, ...


Resonances do not always appear as peaks:



J. R. Taylor, *Scattering Theory — The Quantum Theory on Nonrelativistic Collisions*

# Peaks are not always resonances

- Hadron resonances due to **QCD dynamics**  $\Rightarrow$  poles in the  $S$ -matrix:




- Kinematic** effects  $\Rightarrow$  (normally) branching points of the  $S$ -matrix

- ☞ normal two-body threshold cusp
- ☞ triangle singularity
- ☞ ...

traps/tools in hadron spectroscopy

# Peaks are not always resonances

- Hadron resonances due to **QCD dynamics**  $\Rightarrow$  poles in the  $S$ -matrix:





- **Kinematic** effects  $\Rightarrow$  (normally) branching points of the  $S$ -matrix

- $\Rightarrow$  normal two-body threshold cusp
- $\Rightarrow$  triangle singularity
- $\Rightarrow$  ...

traps/tools in hadron spectroscopy

# Threshold cusp

- There is **always** a cusp at an  $S$ -wave threshold



$$= \text{disc } \mathcal{A}(s) \propto C^*(s) \frac{q_{\text{cm}}(s)}{\sqrt{s}} B(s) \theta(s - (m_1 + m_2)^2)$$



# Threshold cusp: a well-known example

- Cusp effect as a useful tool for precise measurement:

☞ example of the cusp in  $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$

☞ strength of the cusp measures the **interaction strength!**


Meißner, Müller, Steininger (1997); Cabibbo (2004); Colangelo, Gasser, Kubis, Rusetsky (2006); ...



a precise measurement with an uncertainty of about 2%

# $\pi^+\pi^-$ cusp in $\Upsilon(3S) \rightarrow \Upsilon(1S)\pi^0\pi^0$

X.-H. Liu, FKG, E. Epelbaum, EPJC73(2013)2284




measurable at Belle-II

Bin width	Events	$6 \times 10^4$	$6 \times 10^5$	$3 \times 10^6$	$6 \times 10^6$
0.1 MeV	$\chi^2/\text{dof}$	1.21	1.09	1.16	0.88
	$a_0 - a_2$	$0.293 \pm 0.036$	$0.260 \pm 0.012$	$0.2717 \pm 0.0048$	$0.2661 \pm 0.0036$
0.2 MeV	$\chi^2/\text{dof}$	0.72	1.15	1.05	1.12
	$a_0 - a_2$	$0.286 \pm 0.035$	$0.251 \pm 0.014$	$0.2722 \pm 0.0048$	$0.2621 \pm 0.0038$
0.5 MeV	$\chi^2/\text{dof}$	0.93	0.54	1.27	1.30
	$a_0 - a_2$	$0.262 \pm 0.026$	$0.256 \pm 0.012$	$0.2659 \pm 0.0051$	$0.2693 \pm 0.0035$
1 MeV	$\chi^2/\text{dof}$	1.05	0.78	1.17	0.69
	$a_0 - a_2$	$0.221 \pm 0.054$	$0.291 \pm 0.010$	$0.2658 \pm 0.0054$	$0.2661 \pm 0.0037$
2 MeV	$\chi^2/\text{dof}$	0.59	1.06	1.05	1.37
	$a_0 - a_2$	$0.260 \pm 0.040$	$0.262 \pm 0.012$	$0.2592 \pm 0.0055$	$0.2632 \pm 0.0037$



# Triangle singularity



$$\frac{1}{2m_A} \sqrt{\lambda(m_A^2, m_1^2, m_2^2)} \equiv \boxed{p_{2,\text{left}} = p_{2,\text{right}}} \equiv \gamma(\beta E_2^* - p_2^*)$$

**on-shell** momentum of  $m_2$  at the **left** and **right** cuts in the  $A$  rest frame

$$\beta = |\vec{p}_{23}|/E_{23}, \quad \gamma = 1/\sqrt{1 - \beta^2}$$


Bayar et al., PRD94(2016)074039

- $p_2 > 0, p_3 = \gamma(\beta E_3^* + p_2^*) > 0 \Rightarrow m_2$  and  $m_3$  move in the same direction
- velocities in the  $A$  rest frame:  $v_3 > \beta > v_2$

$$v_2 = \beta \frac{E_2^* - p_2^*/\beta}{E_2^* - \beta p_2^*} < \beta, \quad v_3 = \beta \frac{E_3^* + p_2^*/\beta}{E_3^* + \beta p_2^*} > \beta$$

- Conditions (Coleman–Norton theorem): Coleman, Norton (1965); Bronzan (1964)
  - ☞ all three intermediate particles can go on shell simultaneously
  - ☞  $\vec{p}_2 \parallel \vec{p}_3$ , particle-3 can catch up with particle-2 (as a classical process)
- needs very special kinematics  $\Rightarrow$  process dependent! (contrary to pole position)

# Triangle singularity



$$\frac{1}{2m_A} \sqrt{\lambda(m_A^2, m_1^2, m_2^2)} \equiv \boxed{p_{2,\text{left}} = p_{2,\text{right}}} \equiv \gamma (\beta E_2^* - p_2^*)$$

**on-shell** momentum of  $m_2$  at the **left** and **right** cuts in the  $A$  rest frame

$$\beta = |\vec{p}_{23}|/E_{23}, \quad \gamma = 1/\sqrt{1 - \beta^2}$$


Bayar et al., PRD94(2016)074039

- $p_2 > 0, p_3 = \gamma (\beta E_3^* + p_2^*) > 0 \Rightarrow m_2$  and  $m_3$  move in the same direction
- velocities in the  $A$  rest frame:  $v_3 > \beta > v_2$

$$v_2 = \beta \frac{E_2^* - p_2^*/\beta}{E_2^* - \beta p_2^*} < \beta, \quad v_3 = \beta \frac{E_3^* + p_2^*/\beta}{E_3^* + \beta p_2^*} > \beta$$

- Conditions (Coleman–Norton theorem): Coleman, Norton (1965); Bronzan (1964)
  - ☞ all three intermediate particles can go **on shell simultaneously**
  - ☞  $\vec{p}_2 \parallel \vec{p}_3$ , particle-3 can catch up with particle-2 (as a classical process)
- needs very special kinematics  $\Rightarrow$  **process dependent!** (contrary to pole position)

## TS: some details (I)



Consider the scalar three-point loop integral

$$I = i \int \frac{d^4 q}{(2\pi)^4} \frac{1}{[(P - q)^2 - m_1^2 + i\epsilon] (q^2 - m_2^2 + i\epsilon) [(p_{23} - q)^2 - m_3^2 + i\epsilon]}$$



Rewriting a propagator into two poles:

$$\frac{1}{q^2 - m_2^2 + i\epsilon} = \frac{1}{(q^0 - \omega_2 + i\epsilon)(q^0 + \omega_2 - i\epsilon)} \quad \text{with} \quad \omega_2 = \sqrt{m_2^2 + \vec{q}^2}$$

focus on the positive-energy poles

$$I \simeq \frac{i}{8m_1 m_2 m_3} \int \frac{dq^0 d^3 \vec{q}}{(2\pi)^4} \frac{1}{(P^0 - q^0 - \omega_1 + i\epsilon)(q^0 - \omega_2 + i\epsilon)(p_{23}^0 - q^0 - \omega_3 + i\epsilon)}$$

## TS: some details (II)



Contour integral over  $q^0 \Rightarrow$


$$\begin{aligned}
 I &\propto \int \frac{d^3 \vec{q}}{(2\pi)^3} \frac{1}{[P^0 - \omega_1(q) - \omega_2(q) + i\epsilon][p_{23}^0 - \omega_2(q) - \omega_3(\vec{p}_{23} - \vec{q}) + i\epsilon]} \\
 &\propto \int_0^\infty dq \frac{q^2}{P^0 - \omega_1(q) - \omega_2(q) + i\epsilon} f(q)
 \end{aligned}$$

The second cut:

$$f(q) = \int_{-1}^1 dz \frac{1}{p_{23}^0 - \omega_2(q) - \sqrt{m_3^2 + q^2} + p_{23}^2 - 2p_{23}qz + i\epsilon}$$




### Relation between singularities of integrand and integral

- singularity of integrand does **not necessarily** give a singularity of integral:  
integral contour may be deformed to avoid the singularity
- Two cases that a singularity cannot be avoided:
  - ⇨ endpoint singularity
  - ⇨ pinch singularity




### Relation between singularities of integrand and integral

- singularity of integrand does **not necessarily** give a singularity of integral:  
integral contour may be deformed to avoid the singularity
- Two cases that a singularity cannot be avoided:
  - ☞ **endpoint singularity**
  - ☞ **pinch singularity**



## TS: some details (IV)



$$I \propto \int_0^\infty dq \frac{q^2}{P^0 - \omega_1(q) - \omega_2(q) + i\epsilon} f(q)$$

$$f(q) = \int_{-1}^1 dz \frac{1}{A(q, z)} \equiv \int_{-1}^1 dz \frac{1}{p_{23}^0 - \omega_2(q) - \sqrt{m_3^2 + q^2 + p_{23}^2 - 2p_{23}qz} + i\epsilon}$$

Singularities of the **integrand of  $I$**  in the rest frame of initial particle ( $P^0 = M$ ):

- 1st cut:  $M - \omega_1(l) - \omega_2(l) + i\epsilon = 0 \Rightarrow$   

$$q_{\text{on}\pm} \equiv \pm \left( \frac{1}{2M} \sqrt{\lambda(M^2, m_1^2, m_2^2)} + i\epsilon \right)$$
- 2nd cut:  $A(q, \pm 1) = 0 \Rightarrow$  endpoint singularities of  $f(q)$


$$z = +1: \quad q_{a+} = \gamma(\beta E_2^* + p_2^*) + i\epsilon, \quad q_{a-} = \gamma(\beta E_2^* - p_2^*) - i\epsilon,$$

$$z = -1: \quad q_{b+} = \gamma(-\beta E_2^* + p_2^*) + i\epsilon, \quad q_{b-} = -\gamma(\beta E_2^* + p_2^*) - i\epsilon$$

$$\beta = |\vec{p}_{23}|/E_{23}, \quad \gamma = 1/\sqrt{1 - \beta^2} = E_{23}/m_{23}$$

$E_2^*(p_2^*)$ : energy (momentum) of particle-2 in the cmf of the (2,3) system

## TS: some details (IV)



$$I \propto \int_0^\infty dq \frac{q^2}{P^0 - \omega_1(q) - \omega_2(q) + i\epsilon} f(q)$$

$$f(q) = \int_{-1}^1 dz \frac{1}{A(q, z)} \equiv \int_{-1}^1 dz \frac{1}{p_{23}^0 - \omega_2(q) - \sqrt{m_3^2 + q^2 + p_{23}^2 - 2p_{23}qz} + i\epsilon}$$

Singularities of the **integrand of  $I$**  in the rest frame of initial particle ( $P^0 = M$ ):

- **1st cut:**  $M - \omega_1(l) - \omega_2(l) + i\epsilon = 0 \Rightarrow$   

$$q_{\text{on}\pm} \equiv \pm \left( \frac{1}{2M} \sqrt{\lambda(M^2, m_1^2, m_2^2)} + i\epsilon \right)$$
- **2nd cut:**  $A(q, \pm 1) = 0 \Rightarrow$  **endpoint singularities of  $f(q)$**

$$z = +1: \quad q_{a+} = \gamma(\beta E_2^* + p_2^*) + i\epsilon, \quad q_{a-} = \gamma(\beta E_2^* - p_2^*) - i\epsilon,$$

$$z = -1: \quad q_{b+} = \gamma(-\beta E_2^* + p_2^*) + i\epsilon, \quad q_{b-} = -\gamma(\beta E_2^* + p_2^*) - i\epsilon$$

$$\beta = |\vec{p}_{23}|/E_{23}, \quad \gamma = 1/\sqrt{1 - \beta^2} = E_{23}/m_{23}$$

$E_2^*(p_2^*)$ : energy (momentum) of particle-2 in the cmf of the (2,3) system




# TS: some details (V)


All singularities of the integrand of  $I$ :

$$q_{0n+}, \quad q_{a+} = \gamma (\beta E_2^* + p_2^*) + i \epsilon, \quad q_{a-} = \gamma (\beta E_2^* - p_2^*) - i \epsilon,$$


$$q_{0n-} < 0, \quad q_{b-} = -q_{a+} < 0 \text{ (for } \epsilon = 0), \quad q_{b+} = -q_{a-},$$




(a)



(b)



(c)




2-body threshold  
singularity at

$$m_{23} = m_2 + m_3$$

triangle singularity at


$$q_{0n+} = q_{a-}$$

$q_{0n+}$ :  $p_{2,\text{left}}$ ,  $q_{a-}$ :  $p_{2,\text{right}}$  in page 8



$$M_1 = (4380 \pm 8 \pm 29) \text{ MeV},$$

$$M_2 = (4449.8 \pm 1.7 \pm 2.5) \text{ MeV},$$



$$\Gamma_1 = (205 \pm 18 \pm 86) \text{ MeV},$$

$$\Gamma_2 = (39 \pm 5 \pm 19) \text{ MeV}.$$


- Quantum numbers not fully determined, for ( $P_c(4380)$ ,  $P_c(4450)$ ):  
( $3/2^-, 5/2^+$ ), ( $3/2^+, 5/2^-$ ), ( $5/2^+, 3/2^-$ ), ... (more see later slides)
- In  $J/\psi p$  invariant mass distribution, with **hidden charm**  
 $\Rightarrow$  **pentaquarks if they are really hadron states**
- Narrow pentaquark-like structures with hidden-charm had been predicted 5 years before (07.2010):  
*Prediction of narrow  $N^*$  and  $\Lambda^*$  resonances with hidden charm above 4 GeV,*  
J.-J. Wu, R. Molina, E. Oset, B.-S. Zou, *Phys. Rev. Lett.* **105** (2010) 232001
- Pentaquark candidates! thus important to study in great details

# Coincidence of $P_c(4450)$ with kinematic singularities

- Mass:  $M_{P_c(4450)} = (4449.8 \pm 1.7 \pm 2.5) \text{ MeV}$
- Trivial observation:  $P_c(4450)$  coincides with the  $\chi_{c1}p$  threshold:

$$M_{P_c(4450)} - M_{\chi_{c1}} - M_p = (0.9 \pm 3.1) \text{ MeV}$$

- Non-trivial observation: there is a **triangle singularity** at the same time! Solving the equation  $p_{2,\text{left}} = p_{2,\text{right}} \Rightarrow$  to have a TS at  $M_{J/\psi p} = M_{\chi_{c1}} + M_p$ , we need  $M_{\Lambda^*} \simeq 1.89 \text{ GeV}$   
On shell  $\Rightarrow \Lambda^*$  must be unstable, the TS is then a finite peak



More possible relevant TSs, see



X.-H. Liu, Q. Wang, Q. Zhao, PLB757(2015)231

# Coincidence of $P_c(4450)$ with kinematic singularities

- Mass:  $M_{P_c(4450)} = (4449.8 \pm 1.7 \pm 2.5) \text{ MeV}$
- Trivial observation:  $P_c(4450)$  coincides with the  $\chi_{c1}p$  threshold:

$$M_{P_c(4450)} - M_{\chi_{c1}} - M_p = (0.9 \pm 3.1) \text{ MeV}$$

- Non-trivial observation: there is a **triangle singularity** at the same time! Solving the equation  $p_{2,\text{left}} = p_{2,\text{right}} \Rightarrow$  to have a TS at  $M_{J/\psi p} = M_{\chi_{c1}} + M_p$ , we need  $M_{\Lambda^*} \simeq 1.89 \text{ GeV}$   
On shell  $\Rightarrow \Lambda^*$  must be unstable, the TS is then a finite peak





More possible relevant TSs, see

X.-H. Liu, Q. Wang, Q. Zhao, PLB757(2015)231

# Trajectories of triangle singularities in complex energy plane

Dalitz plot for  $\Lambda_b \rightarrow \chi_{c1} p K^-$ :



numbers: assumed masses for  $\Lambda^*$

**blue**: proton and  $\chi_{c1}$  are **parallel**, in the **2nd Riemann sheet**

**green**: proton and  $\chi_{c1}$  are **anti-parallel**



$$M_{\Lambda_b} = 5.62 \text{ GeV}, M_{\chi_{c1}} = 3.51 \text{ GeV}, \quad \sqrt{s} \equiv M(\chi_{c1} p)$$

$$M_{K^- p, A} = M_{\Lambda_b} - M_{\chi_{c1}}, \quad M_{K^- p, B} = \sqrt{\frac{M_{\Lambda_b}^2 M_p + M_K^2 M_{\chi_{c1}}}{M_{\chi_{c1}} + M_p} - M_{\chi_{c1}} M_p}$$

# TS for $P_c(4450)$

FKG et al., PRD92(2015)071502(R); X.-H. Liu, Q. Wang, Q. Zhao, PLB757(2015)231

- When  $M_{\Lambda^*} = 1.89$  GeV, TS is located **exactly at the  $\chi_{c1}p$  threshold, 4.449 GeV!**
- **Four-star baryon  $\Lambda(1890)$ :  $J^P = 3/2^+$ ,  $\Gamma$ : 60 – 200 MeV**
- triangle loop with  **$S$ -wave  $\chi_{c1}p$ :  $J^P = \frac{1}{2}^+$  or  $\frac{3}{2}^+$**





- impossible to produce a narrow peak for  $\chi_{c1}p$  in other partial waves

Bayar et al., PRD94(2016)074039

# TS for $P_c(4450)$

FKG et al., PRD92(2015)071502(R); X.-H. Liu, Q. Wang, Q. Zhao, PLB757(2015)231

- When  $M_{\Lambda^*} = 1.89$  GeV, TS is located exactly at the  $\chi_{c1}p$  threshold, 4.449 GeV!
- Four-star baryon  $\Lambda(1890)$ :  $J^P = 3/2^+$ ,  $\Gamma$ : 60 – 200 MeV
- triangle loop with  $S$ -wave  $\chi_{c1}p$ :  $J^P = \frac{1}{2}^+$  or  $\frac{3}{2}^+$



- impossible to produce a narrow peak for  $\chi_{c1}p$  in other partial waves

Bayar et al., PRD94(2016)074039



## TS for $P_c(4450)$ : Comments

- Position of the TS completely fixed; shape also largely fixed
- but, strength of the TS is **unknown**
- operative in  $J/\psi\pi$  quantum numbers  $J^P = \frac{1}{2}^+$  or  $\frac{3}{2}^+$

$J^P(4380, 4450)$	$(\sqrt{\Delta(-2\ln\mathcal{L})})^2$	$P_c(4380)$		$P_c(4450)$	
		$M_0$	$\Gamma_0$	$M_0$	$\Gamma_0$
(3/2-, 5/2+) solution					
3/2-, 5/2+	--	4359	151	4450.1	49
$\Delta$ from (3/2-, 5/2+) solution					
5/2+, 3/2-	-3.6 <sup>2</sup>	10	-7	-1.6	-6
5/2-, 3/2+	-2.7 <sup>2</sup>	-4	-9	-3.6	-2
3/2-, 5/2+	-	-	-	-	-

from a reanalysis of the LHCb data using an extended  $\Lambda^*$  model

N. Jurik, CERN-THESIS-2016-086

- does not exclude the possibility of the existence of a pentaquark in addition

# TS for $P_c(4450)$ : Comments

- Position of the TS completely fixed; shape also largely fixed
- but, strength of the TS is **unknown**
- operative in  $J/\psi\pi$  quantum numbers  $J^P = \frac{1}{2}^+$  or  $\frac{3}{2}^+$

$J^P(4380, 4450)$	$(\sqrt{\Delta(-2\ln\mathcal{L})})^2$	$P_c(4380)$		$P_c(4450)$	
		$M_0$	$\Gamma_0$	$M_0$	$\Gamma_0$
(3/2-, 5/2+) solution					
3/2-, 5/2+	--	4359	151	4450.1	49
$\Delta$ from (3/2-, 5/2+) solution					
5/2+, 3/2-	-3.6 <sup>2</sup>	10	-7	-1.6	-6
5/2-, 3/2+	-2.7 <sup>2</sup>	-4	-9	-3.6	-2
3/2-, 5/2+	-	-	-	-	-

from a reanalysis of the LHCb data using an extended  $\Lambda^*$  model

N. Jurik, CERN-THESIS-2016-086

- does not exclude the possibility of the existence of a pentaquark in addition



# How to distinguish a TS from a genuine resonance?

- Schmid theorem:

C. Schmid, Phys. Rev. 154 (1967) 1363

see also, A. V. Anisovich, V. V. Anisovich, Phys. Lett. B 345 (1995) 321

Triangle singularity **cannot** produce an additional peak in the invariant mass distribution of the **elastic channel** when neglecting inelasticity



Nearby the effective singularity:

$$\mathcal{A}_{(a)+(b)}(s) \sim [1 + 2i\rho(s)T(s)] \mathcal{A}_{(a)}(s) = e^{2i\delta_{\chi_{c1}p}(s)} \mathcal{A}_{(a)}(s)$$

here  $\delta_{\chi_{c1}p}$  is the elastic  $\chi_{c1}p$  scattering phase shift

- corrections from **coupled channels**

A. Szczepaniak, PLB757(2016)61

# How to distinguish a TS from a genuine resonance?


- determining quantum numbers unambiguously:  
TS as discussed here requires the  $\chi_{c1}p$  in *S*-wave  $\Rightarrow J^P = \frac{1}{2}^+$  or  $\frac{3}{2}^+$
- processes (such as photoproduction) with a **different kinematics**  
Q. Wang, X.-H. Liu, Q. Zhao, PRD92(2015)034022;  
V. Kubarovsky, M. Voloshin, PRD92(2015)031502;  
M. Karliner, J. L. Rosner, PLB752(2015)329; ...
- measuring the process  $\Lambda_b^0 \rightarrow \chi_{c1} p K^-$   
☞ if a narrow near-threshold peak in  $\chi_{c1} p \Rightarrow$  a real exotic resonance  
recently measured by LHCb in PRL119(2017)062001, no invariant mass distribution reported:

$$\mathcal{B}(\Lambda_b^0 \rightarrow \chi_{c1} p K^-) = (7.4 \pm 0.4 \pm 0.4 \pm 0.6_{-0.7}^{+1.0}) \times 10^{-5}$$
$$\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi p K^-) = (3.01 \pm 0.22_{-0.27}^{+0.43}) \times 10^{-4}$$

With LHC Run-1 data, statistics not enough N. Jurik, Mitsuyoshi Tanaka Dissertation Award  
Talk at the APS April Meeting 2018

# $P_s$ searching





- A  $\phi p$  bound state was predicted in several models with a mass  $\sim 2$  GeV  
 H. Gao, T.S.H. Lee, V. Marinov, PRC63(2001)022201;  
 F. Huang, Z.-Y. Zhang, Y.-W. Yu, PRC73(2006)025207;  
 H. Gao, H. Huang, T. Liu, J. Ping, F. Wang, Z. Zhao, PRC95(2017)055202
- Lattice evidence for strangonium-nucleon bound state at a large quark mass  
 $m_{u,d,s}^{\text{Lat.}} = m_s^{\text{ph.}}$  ( $M_\pi^{\text{Lat.}} \simeq 805$  MeV) S.R. Beane et al. [NPLQCD], PRD91(2015)114503
- Bump observed at  $\sqrt{s} \sim 2$  GeV by LEPS and CLAS in  $\gamma p \rightarrow \phi p$   
 LEPS, PRL95(2005)182001; CLAS, PRC89(2014)055208, PRC90(2014)019901
- Suggestion to search for  $P_s$  in  $\Lambda_c \rightarrow \pi^0 \phi p$  R. Lebed, PRD92(2015)114030



- No clear evidence was found in Belle searching Belle, PRD96(2017)051102(R)


# TS and $P_s$ in $\Lambda_c \rightarrow p\phi\pi^0$

J.-J. Xie, FKG, PLB774(2017)108



Model I: the  $BV$  interaction model ( $P_s$  generated) of [A. Ramos, E. Oset, PLB727\(2013\)287](#);

Model II: no resonance, constant interaction; Model III: phase space




- TS produces a bump at around 2.02 GeV, width mainly from that of  $K^*$
- $P_s$ , if exists, could distort the line shape, but difficult to be distinguished from TS in this process
- A measurement of  $\Lambda_c \rightarrow \Sigma^* K^*$  can help constrain the TS strength


# $Z_b/Z_c$ : threshold cusps?

- Models of  $Z_b(10610, 10650)$ ,  $Z_c(3900, 4020)$  as threshold cusps

👉 Bugg, Swanson:



D. Bugg, EPL96(2011)11002; E. Swanson, PRD91(2015)034009



👉 Initial pion radiation: D.-Y.Chen, X.Liu, PRD84(2011)094003; PRD84(2011)034032; Chen, Liu, Matsuki, PRD84(2011)074032; PRL110(2013)232001; ...





# $Z_b/Z_c$ : threshold cusps?


- Models of  $Z_b(10610, 10650)$ ,  $Z_c(3900, 4020)$  as threshold cusps

👉 Bugg, Swanson:

D. Bugg, EPL96(2011)11002; E. Swanson, PRD91(2015)034009





👉 Initial pion radiation: D.-Y.Chen, X.Liu, PRD84(2011)094003; PRD84(2011)034032; Chen, Liu, Matsuki, PRD84(2011)074032; PRL110(2013)232001; ...



## $Z_b/Z_c$ : threshold cusps?

- But  $Z_c(3900)[Z_b]$  as a narrow peak in  $DD^*[B\bar{B}^*]$  distribution cannot be only due to cusp: **prominent cusp  $\Rightarrow$  strong int.  $\Rightarrow$  pole!**

FKG, Hanhart, Wang, Zhao, PRD91(2015)051504



Black curve: up to 1 loop with  $C_\Lambda G_\Lambda(E_{\text{th}}) = -1/2$ ,  
no narrow peak any more!



$g_Y [1 + C_\Lambda G_\Lambda(E) + C_\Lambda G_\Lambda(E)C_\Lambda G_\Lambda(E) + \dots]$  produces a pole

- so far, triangle diagrams not considered (see next slides)

# $Z_b/Z_c$ : threshold cusps?

- But  $Z_c(3900)[Z_b]$  as a narrow peak in  $DD^*[B\bar{B}^*]$  distribution cannot be only due to cusp: **prominent cusp  $\Rightarrow$  strong int.  $\Rightarrow$  pole!**

FKG, Hanhart, Wang, Zhao, PRD91(2015)051504



Black curve: up to 1 loop with  $C_\Lambda G_\Lambda(E_{th}) = -1/2$ ,  
no narrow peak any more!



$g_Y [1 + C_\Lambda G_\Lambda(E) + C_\Lambda G_\Lambda(E)C_\Lambda G_\Lambda(E) + \dots]$  produces a pole

- so far, triangle diagrams not considered (see next slides)

## $Z_b/Z_c$ : threshold cusps?

- But  $Z_c(3900)[Z_b]$  as a narrow peak in  $DD^*[B\bar{B}^*]$  distribution cannot be only due to cusp: **prominent cusp  $\Rightarrow$  strong int.  $\Rightarrow$  pole!**

FKG, Hanhart, Wang, Zhao, PRD91(2015)051504



Black curve: up to 1 loop with  $C_\Lambda G_\Lambda(E_{\text{th}}) = -1/2$ ,  
**no narrow peak any more!**



$g_Y [1 + C_\Lambda G_\Lambda(E) + C_\Lambda G_\Lambda(E)C_\Lambda G_\Lambda(E) + \dots]$  produces a pole

- so far, **triangle diagrams** not considered (see next slides)

## $Z_b/Z_c$ : threshold cusps?

- But  $Z_c(3900)[Z_b]$  as a narrow peak in  $DD^*[B\bar{B}^*]$  distribution cannot be only due to cusp: **prominent cusp**  $\Rightarrow$  **strong int.**  $\Rightarrow$  **pole!**

FKG, Hanhart, Wang, Zhao, PRD91(2015)051504




Black curve: up to 1 loop with  $C_\Lambda G_\Lambda(E_{\text{th}}) = -1/2$ ,  
**no narrow peak any more!**

$g_Y [1 + C_\Lambda G_\Lambda(E) + C_\Lambda G_\Lambda(E)C_\Lambda G_\Lambda(E) + \dots]$  produces a pole

- so far, **triangle diagrams** not considered (see next slides)

## $Z_c(3900)$ : triangle diagram

- Consider the triangle loop:




- For  $E_{\text{cm}} = 4.26$  GeV, TS in the unphysical region


- Enhancement very sensitive to the cm energy

## $Z_c(3900)$ : triangle diagram

- Consider the triangle loop:




- For  $E_{\text{cm}} = 4.26$  GeV, TS in the unphysical region




- Enhancement very sensitive to the cm energy

# $Z_c(3900)$ : triangle diagram



- Consider the triangle loop:



- For  $E_{\text{cm}} = 4.26$  GeV, TS in the unphysical region



- Enhancement very sensitive to the cm energy








# More about $Z_c(3900)$

Triangle + coupled-channel FSI

Albaladejo, FKG, Hidalgo-Duque, Nieves, PLB755(2016)337



# More about $Z_c(3900)$






$M_{Z_c}$ (MeV)	$\Gamma_{Z_c}/2$ (MeV)	Ref.	Final state
$3899 \pm 6$	$23 \pm 11$	[1] (BESIII)	$J/\psi \pi$
$3895 \pm 8$	$32 \pm 18$	[2] (Belle)	$J/\psi \pi$
$3886 \pm 5$	$19 \pm 5$	[3] (CLEO-c)	$J/\psi \pi$
$3884 \pm 5$	$12 \pm 6$	[4] (BESIII)	$\bar{D}^* D$
$3882 \pm 3$	$13 \pm 5$	[5] (BESIII)	$\bar{D}^* D$
$3894 \pm 6 \pm 1$	$30 \pm 12 \pm 6$	$\Lambda_2 = 1.0$ GeV	$J/\psi \pi, \bar{D}^* D$
$3886 \pm 4 \pm 1$	$22 \pm 6 \pm 4$	$\Lambda_2 = 0.5$ GeV	$J/\psi \pi, \bar{D}^* D$
$3831 \pm 26^{+7}_{-28}$	virtual state	$\Lambda_2 = 1.0$ GeV	$J/\psi \pi, \bar{D}^* D$
$3844 \pm 19^{+12}_{-21}$	virtual state	$\Lambda_2 = 0.5$ GeV	$J/\psi \pi, \bar{D}^* D$

resonance pole

or virtual state

# $Z_c(3900)$ : Interpreting lattice results by Prelovsek et al.

Albaladejo, Fernandez-Soler, Nieves, EPJC76(2016)573




- Lat. ( $M_\pi = 266$  MeV): [Prelovsek et al., PRD91\(2015\)014504](#)  
“no additional eigenstate” corresponding to  $Z_c$
- In finite volume ( $L = 2$  fm): consistent with lattice energy levels, but **with a pole in continuum!**

## $Y(4260) \rightarrow Z_c \pi$ : TS or not?


- Importance of TS in  $Y(4260) \rightarrow Z_c \pi$  already noticed, but  $Z_c$  pole still needed  
Q.Wang, Hanhart, Q.Zhao, PRL111(2013)132002; PLB725(2013)106
- however, debate continues: whether  $Z_c$  pole is needed seems still inconclusive  
Pilloni et al. (JPAC), PLB772(2017)200

# $Y(4260) \rightarrow Z_c \pi$ : TS or not?

- Importance of TS in  $Y(4260) \rightarrow Z_c \pi$  already noticed, but  $Z_c$  pole still needed  
Q.Wang, Hanhart, Q.Zhao, PRL111(2013)132002; PLB725(2013)106
- however, debate continues: whether  $Z_c$  pole is needed seems still inconclusive  
Pilloni et al. (JPAC), PLB772(2017)200



Pole+TS:




# $Y(4260) \rightarrow Z_c \pi$ : TS or not?

- Importance of TS in  $Y(4260) \rightarrow Z_c \pi$  already noticed, but  $Z_c$  pole still needed



Q.Wang, Hanhart, Q.Zhao, PRL111(2013)132002; PLB725(2013)106

- however, debate continues: whether  $Z_c$  pole is needed seems still inconclusive


Pilloni et al. (JPAC), PLB772(2017)200



Pole+TS:




Only TS:



But...

New data from BESIII in the  $J/\psi\pi^+\pi^-$  channel

BESIII, PRL119(2017)072001



## To-do list

- To search for resonances in processes with different kinematics, and to measure the quantum numbers
- To estimate the strength of the TS contributions, [see, e.g., many papers by Eulogio](#)
- Analysis framework incorporating kinematic singularities
- **Not just traps, but also tools:**  
TS enhancement  $\Rightarrow$  enhanced production;  
 $S$ -wave  $\Rightarrow$  quantum number filter



- To search for resonances in processes with different kinematics, and to measure the quantum numbers
- To estimate the strength of the TS contributions, [see, e.g., many papers by Eulogio](#)
- Analysis framework incorporating kinematic singularities
- **Not just traps, but also tools:**
  - TS enhancement  $\Rightarrow$  enhanced production;
  - $S$ -wave  $\Rightarrow$  quantum number filter

THANK YOU FOR YOUR  
ATTENTION!

# Backup slides

## Triangle singularity – literature

- Some recent works using **triangle singularity** to explain (part of) peak structures [ $\eta(1405/1475)$ ,  $a_1(1420)$ , ...]:

J.-J. Wu, X.-H. Liu, Q. Zhao and B.-S. Zou, PRL108(2012)081803;

X.-G. Wu, J.-J. Wu, Q. Zhao and B.-S. Zou, PRD87(2013)014023(2013);

Q. Wang, C. Hanhart and Q. Zhao, PLB725(2013)106;

M. Mikhasenko, B. Ketzner and A. Sarantsev, PRD91(2015)094015;

X.-H. Liu, M. Oka and Q. Zhao, PLB753(2016)297;

A. P. Szczepaniak, PLB747(2015)410; PLB757(2016)61;

F. Aceti, L.-R. Dai and E. Oset, PRD94(2016)096015;

A. E. Bondar and M. B. Voloshin, PRD93(2016)094008

V. R. Debastiani, F. Aceti, W.-H. Liang, E. Oset, PRD95(2017)034015

.....

Recent reviews:



Q.Zhao, JPS Conf.Proc.13(2017)010008; FKG et al., RMP90(2018)015004

Recent lecture notes by one of the key players:

I. J. R. Aitchison, arXiv:1507.02697 [hep-ph], *Unitarity, Analyticity and Crossing Symmetry in Two- and Three-hadron Final State Interactions*

J.-J. Wu, X.-H. Liu, Q. Zhao and B.-S. Zou, PRL108(2012)081803;

X.-G. Wu, J.-J. Wu, Q. Zhao and B.-S. Zou, PRD87(2013)014023(2013)



Unique consequence: huge isospin breaking, **vary narrow  $f_0(980)$  peak  $\sim 10$  MeV**

