


# Excited $\Omega_c$ 's as Possible exotic states

Hyun-Chul Kim

Department of Physics, Inha University  
Incheon, Korea


NFQCD@YITP 15 June, 2018, Kyoto, Japan

# LHCb Findings: New five $\Omega_{cs}$



Five  $\Omega_{cs}$  were announced by LHCb Coll.

# Belle Findings: Four $\Omega_{cs}$ confirmed



Four  $\Omega_{cs}$  were confirmed by Belle Coll.

# LHCb Findings: New five Omega\_cs

The Widths are rather small, even if we consider the fact that heavy baryons have smaller widths than light ones.

Resonance	Mass ( MeV)	$\Gamma$ ( MeV)	Yield	$N_\sigma$
$\Omega_c(3000)^0$	$3000.4 \pm 0.2 \pm 0.1^{+0.3}_{-0.5}$	$4.5 \pm 0.6 \pm 0.3$	$1300 \pm 100 \pm 80$	20.4
$\Omega_c(3050)^0$	$3050.2 \pm 0.1 \pm 0.1^{+0.3}_{-0.5}$	$0.8 \pm 0.2 \pm 0.1$	$970 \pm 60 \pm 20$	20.4
		$< 1.2 \text{ MeV, 95\% CL}$		
$\Omega_c(3066)^0$	$3065.6 \pm 0.1 \pm 0.3^{+0.3}_{-0.5}$	$3.5 \pm 0.4 \pm 0.2$	$1740 \pm 100 \pm 50$	23.9
$\Omega_c(3090)^0$	$3090.2 \pm 0.3 \pm 0.5^{+0.3}_{-0.5}$	$8.7 \pm 1.0 \pm 0.8$	$2000 \pm 140 \pm 130$	21.1
$\Omega_c(3119)^0$	$3119.1 \pm 0.3 \pm 0.9^{+0.3}_{-0.5}$	$1.1 \pm 0.8 \pm 0.4$	$480 \pm 70 \pm 30$	10.4
		$< 2.6 \text{ MeV, 95\% CL}$		
$\Omega_c(3188)^0$	$3188 \pm 5 \pm 13$	$60 \pm 15 \pm 11$	$1670 \pm 450 \pm 360$	
$\Omega_c(3066)_{\text{fd}}^0$			$700 \pm 40 \pm 140$	
$\Omega_c(3090)_{\text{fd}}^0$			$220 \pm 60 \pm 90$	
$\Omega_c(3119)_{\text{fd}}^0$			$190 \pm 70 \pm 20$	

LHCb Collaboration, 2017

$\Omega_c$ Excited State	3000	3050	3066	3090	3119	3188
Yield	$37.7 \pm 11.0$	$28.2 \pm 7.7$	$81.7 \pm 13.9$	$86.6 \pm 17.4$	$3.6 \pm 6.9$	$135.2 \pm 43.0$
Significance	$3.9\sigma$	$4.6\sigma$	$7.2\sigma$	$5.7\sigma$	$0.4\sigma$	$2.4\sigma$
LHCb Mass	$3000.4 \pm 0.2 \pm 0.1$	$3050.2 \pm 0.1 \pm 0.1$	$3065.5 \pm 0.1 \pm 0.3$	$3090.2 \pm 0.3 \pm 0.5$	$3119 \pm 0.3 \pm 0.9$	$3188 \pm 5 \pm 13$
Belle Mass (with fixed $\Gamma$ )	$3000.7 \pm 1.0 \pm 0.2$	$3050.2 \pm 0.4 \pm 0.2$	$3064.9 \pm 0.6 \pm 0.2$	$3089.3 \pm 1.2 \pm 0.2$	-	$3199 \pm 9 \pm 4$

Belle Collaboration, PRD 97 (2018) 051102



# Baryon in pion mean fields

- \* A **baryon** can be viewed as a state of  $N_c$  quarks bound by mesonic **mean fields** (E. Witten, NPB, 1979 & 1983).

Its mass is proportional to  $N_c$ , while its width is of order  $O(1)$ .

- Mesons are weakly interacting (Quantum fluctuations are suppressed by  $1/N_c$ :  $O(1/N_c)$ ).


## Meson mean-field approach (Chiral Quark-Soliton Model)

- \* Baryons as a state of  $N_c$  quarks bound by mesonic mean fields.

$$S_{\text{eff}} = -N_c \text{Tr} \ln (i\cancel{D} + iMU\gamma^5 + i\hat{m})$$


- \* **Key point: Hedgehog Ansatz**

$$\pi^a(\mathbf{r}) = \begin{cases} n^a F(r), & n^a = x^a/r, \quad a = 1, 2, 3 \\ 0, & a = 4, 5, 6, 7, 8. \end{cases}$$



- It breaks spontaneously  $SU(3)_{\text{flavor}} \otimes O(3)_{\text{space}} \rightarrow SU(2)_{\text{isospin+space}}$

# Light baryons in pion mean fields




$$\langle J_B J_B^\dagger \rangle_0 \sim e^{-N_c E_{\text{val}} T}$$

Presence of  $N_c$  quarks will polarize the vacuum or create mean fields.

$N_c$  valence quarks  $\longrightarrow$  Vacuum polarization or meson mean fields

*self-consistent way*

# Light baryons in pion mean fields



$$E_{cl} = N_c E_{val} + E_{sea}$$




Classical Nucleon mass is described by the  $N_c$  valence quark energy and sea-quark energy.

$$\frac{\delta E_{cl}}{\delta U} = 0$$



$M_{cl}$




$P(r)$

$P(r)$ : Soliton profile function or Soliton field

*~ Solving Eq. of motion*



# Light baryons in pion mean fields



- u,d quark levels are classified by the grand spin  $K^P$ .

- s quark levels are classified by the grand spin  $J^P$ .

Grand spin:  $K = J + T$

$$J = L + S$$


# Lowest-lying light baryons

$$K = J + T = 0, \quad T_8 = \frac{N_c}{2\sqrt{3}}$$

Right hypercharge

$$Y' = \frac{N_c}{3}$$

- $N_c$  quark gives the baryon number in the XQSM.



Ground state

# Success of the XQSM in the light baryon sector

- Connection to QCD via the instanton vacuum (natural scale)  
 $\rho \approx 600 \text{ MeV}$
- The mass splittings of the lowest-lying hyperons
- **All different types of baryon form factors**
- Parton distribution amplitudes (u-d asymmetry, transversity)
- Quasi-parton distribution amplitudes
- GPDs






Heavy baryon as a baryon with  $N_c - 1$  light valence quarks

# Singly heavy baryons in $SU(3)$

- \* In the heavy quark mass limit, a heavy quark spin is conserved, so light-quark spin is also conserved.
- \* In this limit, a heavy quark can be considered as **a color static source**.
- \* Dynamics is governed by light quarks.

$$3 \otimes 3 = \bar{3} \oplus 6$$



# Heavy baryons

- \* Valence quarks are bound by the meson mean fields.
- \* Light quarks govern a heavy-light quark system.
- \* Heavy quarks can be simply viewed as **static color sources**.

$$K = J + T = 0, \quad T_8 = \frac{N_c - 1}{2\sqrt{3}} \quad \text{Ground-state heavy baryons}$$

$$\text{Right hypercharge} \quad Y' = \frac{N_c - 1}{3}$$

A heavy quark: Static color source to make a heavy baryon color singlet.


D. Diakonov, arXiv:1003.2157 [hep-ph].



# Heavy baryons



$$K^P = 0^+$$

Dirac Sea



Heavy quark  
as a color static source

# Heavy baryons



Heavy quark  
as a color static source

$N_c - 1$  light quarks govern a singly heavy baryon.

# Heavy baryons

$N_c - 1$  quarks represent heavy-baryon spectra.


$$Y' = \frac{N_c - 1}{3}$$

Grand spin:  $K = 0 \rightarrow T = J$

- The lowest rotationally excited states  $\mathbf{3} \times \mathbf{3} = \bar{\mathbf{3}} + \mathbf{6}$

- \*  $T=0$  for a anti-triplet:  $J=0$  for it. Combining a charm quark with spin  $1/2$ , we have one anti-triplet.

- \*  $T=1$  for a sextet:  $J=1$ . We have **two** sextets with a charm quark. ( $1/2, 3/2$ ).



# SU(3) symmetry breaking

- The collective Hamiltonian for SU(3) symmetry breaking

$$H_{\text{br}} = \alpha D_{88}^{(8)} + \beta Y + \frac{\gamma}{\sqrt{3}} \sum_{i=1}^3 D_{8i}^{(8)} J_i$$

In the light-quark sector, we have already fixed these dynamical parameters as

# SU(3) symmetry breaking

- The collective Hamiltonian for SU(3) symmetry breaking

$$H_{\text{br}} = \alpha D_{88}^{(8)} + \beta Y + \frac{\gamma}{\sqrt{3}} \sum_{i=1}^3 D_{8i}^{(8)} J_i$$

In the light-quark sector, we have already fixed these dynamical parameters as



$$\alpha = -\frac{2m_s}{3}\sigma - \beta Y' = -(255.03 \pm 5.82) \text{ MeV}$$

$$\beta = -\frac{m_s K_2}{I_2} = -(140.04 \pm 3.20) \text{ MeV}$$

$$\gamma = \frac{2m_s K_1}{I_1} + 2\beta = -(101.08 \pm 2.33) \text{ MeV}$$

$$\alpha \rightarrow \bar{\alpha} = \frac{N_c - 1}{N_c} \alpha$$

# Hyperfine mass splittings (only new parameter)




$$\frac{\kappa}{m_c} = (68.1 \pm 1.1) \text{ MeV}$$

$$\frac{\kappa}{m_b} = (20.3 \pm 1.0) \text{ MeV}$$

Remind you that all the parameters are the same as in the light baryon sector except for the hyperfine interaction.

# Hyperfine mass splittings (only new parameter)



Hyperfine splitting between different spin states


$$H_{LQ} = \frac{2}{3} \frac{\kappa}{m_Q M_{\text{sol}}} \mathbf{S}_L \cdot \mathbf{S}_Q = \frac{2}{3} \frac{\varkappa}{m_Q} \mathbf{S}_L \cdot \mathbf{S}_Q$$

$$\frac{\varkappa}{m_c} = (68.1 \pm 1.1) \text{ MeV}$$

$$\frac{\varkappa}{m_b} = (20.3 \pm 1.0) \text{ MeV}$$

Remind you that all the parameters are the same as in the light baryon sector except for the hyperfine interaction.

# Hyperfine mass splittings (only new parameter)



Hyperfine splitting between different spin states

$$H_{LQ} = \frac{2}{3} \frac{\kappa}{m_Q M_{\text{sol}}} \mathbf{S}_L \cdot \mathbf{S}_Q = \frac{2}{3} \frac{\kappa}{m_Q} \mathbf{S}_L \cdot \mathbf{S}_Q$$

The ratio can be determined by the center values of the sextet masses

$$\frac{\kappa}{m_c} = (68.1 \pm 1.1) \text{ MeV}$$

$$\frac{\kappa}{m_b} = (20.3 \pm 1.0) \text{ MeV}$$

Remind you that all the parameters are the same as in the light baryon sector except for the hyperfine interaction.



# Results for the charmed baryon masses

$\mathcal{R}_J^Q$	$B_c$	Mass	Experiment [17]	Deviation $\xi_c$
$\bar{\mathbf{3}}_{1/2}^c$	$\Lambda_c$	$2272.5 \pm 2.3$	$2286.5 \pm 0.1$	$-0.006$
	$\Xi_c$	$2476.3 \pm 1.2$	$2469.4 \pm 0.3$	$0.003$
$\mathbf{6}_{1/2}^c$	$\Sigma_c$	$2445.3 \pm 2.5$	$2453.5 \pm 0.1$	$-0.003$
	$\Xi'_c$	$2580.5 \pm 1.6$	$2576.8 \pm 2.1$	$0.001$
	$\Omega_c$	$2715.7 \pm 4.5$	$2695.2 \pm 1.7$	$0.008$
$\mathbf{6}_{3/2}^c$	$\Sigma_c^*$	$2513.4 \pm 2.3$	$2518.1 \pm 0.8$	$-0.002$
	$\Xi_c^*$	$2648.6 \pm 1.3$	$2645.9 \pm 0.4$	$0.001$
	$\Omega_c^*$	$2783.8 \pm 4.5$	$2765.9 \pm 2.0$	$0.006$

The results are in remarkable agreement with the experimental data.

$$\xi_c = (M_{\text{th}}^{B_c} - M_{\text{exp}}^{B_c})/M_{\text{exp}}^{B_c}$$

# Results for the bottom baryon masses

$\mathcal{R}_J^Q$	$B_b$	Mass	Experiment [17]	Deviation $\xi_b$
$\overline{\mathbf{3}}_{1/2}^b$	$\Lambda_b$	$5599.3 \pm 2.4$	$5619.5 \pm 0.2$	$-0.004$
	$\Xi_b$	$5803.1 \pm 1.2$	$5793.1 \pm 0.7$	$0.002$
$\mathbf{6}_{1/2}^b$	$\Sigma_b$	$5804.3 \pm 2.4$	$5813.4 \pm 1.3$	$-0.002$
	$\Xi'_b$	$5939.5 \pm 1.5$	$5935.0 \pm 0.05$	$0.001$
	$\Omega_b$	$6074.7 \pm 4.5$	$6048.0 \pm 1.9$	$0.004$
$\mathbf{6}_{3/2}^b$	$\Sigma_b^*$	$5824.6 \pm 2.3$	$5833.6 \pm 1.3$	$-0.002$
	$\Xi_b^*$	$5959.8 \pm 1.2$	$5955.3 \pm 0.1$	$0.001$
	$\Omega_b^*$	$6095.0 \pm 4.4$	—	—

$$\xi_b = (M_{\text{th}}^{B_b} - M_{\text{exp}}^{B_b}) / M_{\text{exp}}^{B_b}$$


# Results for the bottom baryon masses

$\mathcal{R}_J^Q$	$B_b$	Mass	Experiment [17]	Deviation $\xi_b$
$\bar{\mathbf{3}}_{1/2}^b$	$\Lambda_b$	$5599.3 \pm 2.4$	$5619.5 \pm 0.2$	$-0.004$
	$\Xi_b$	$5803.1 \pm 1.2$	$5793.1 \pm 0.7$	$0.002$
$\mathbf{6}_{1/2}^b$	$\Sigma_b$	$5804.3 \pm 2.4$	$5813.4 \pm 1.3$	$-0.002$
	$\Xi'_b$	$5939.5 \pm 1.5$	$5935.0 \pm 0.05$	$0.001$
	$\Omega_b$	$6074.7 \pm 4.5$	$6048.0 \pm 1.9$	$0.004$
$\mathbf{6}_{3/2}^b$	$\Sigma_b^*$	$5824.6 \pm 2.3$	$5833.6 \pm 1.3$	$-0.002$
	$\Xi_b^*$	$5959.8 \pm 1.2$	$5955.3 \pm 0.1$	$0.001$
	$\Omega_b^*$	$6095.0 \pm 4.4$	—	—

**Prediction from the present work**

$$\xi_b = (M_{\text{th}}^{B_b} - M_{\text{exp}}^{B_b}) / M_{\text{exp}}^{B_b}$$

# Excited anti-triplets and sextets




$$K = J + T \neq 0$$

$$J = |T - K|, \dots |T + K|$$

# Excited anti-triplets and sextets

Grand spin:  $K^p = 1^-$

- \* Quantization of excited baryons yield **two** anti-triplet and **FIVE** sextets.




$$K = J + T = 1$$

$$J = |T - K|, \dots, |T + K|$$


- \* **T=0** for an anti-triplet:  $J=1$  for it. Combining a charm quark with spin  $1/2$ , we have **two** anti-triplets  $(1/2)$  and  $(3/2)$ .
- \* **T=1** for a sextet:  $J=0, 1, 2$  for 6. We have **five** sextets with a charm quark  $(1/2)$ ,  $(1/2, 3/2)$ , and  $(3/2, 5/2)$ !

# Hyperfine splittings for excited anti-triplets

- Candidates for excited anti-triplets



# Hyperfine splittings for excited sextets



- The mean-field approach (XQSM) predicts **five excited sextet states**.
- The splitting between  $J=1$  and  $J=2$  is twice as large as that between  $J=0$  and  $J=1$ . ( $\Delta_2 = 2\Delta_1$ )

# Scenario I

Assertion: **Five**  $\Omega_c^*$ 's belong to excited sextets.


$J$	$S^P$	$M$ [MeV]	$\kappa'/m_c$ [MeV]	$\Delta_J$ [MeV]
0	$\frac{1}{2}^-$	3000	—	—
1	$\frac{1}{2}^-$	3050	16	61
	$\frac{3}{2}^-$	3066		
2	$\frac{3}{2}^-$	3090	17	47
	$\frac{5}{2}^-$	3119		



# Scenario I

Assertion: Five  $\Omega_c^*$ 's belong to excited sextets.

$J$	$S^P$	$M$ [MeV]	$\kappa'/m_c$ [MeV]	$\Delta_J$ [MeV]
0	$\frac{1}{2}^-$	3000	—	—
1	$\frac{1}{2}^-$	3050	16	61
	$\frac{3}{2}^-$	3066		
2	$\frac{3}{2}^-$	3090	17	47
	$\frac{5}{2}^-$	3119		


$$\frac{\kappa'}{m_c} = 30 \text{ MeV}$$

The HF splittings are **very much deviated** from what we have determined from the excited anti-triplet.

# Scenario I

Assertion: Five  $\Omega_c^*$ 's belong to excited sextets.

$J$	$S^P$	$M$ [MeV]	$\kappa'/m_c$ [MeV]	$\Delta_J$ [MeV]
0	$\frac{1}{2}^-$	3000	—	—
1	$\frac{1}{2}^-$	3050	16	61
	$\frac{3}{2}^-$	3066		
2	$\frac{3}{2}^-$	3090	17	47
	$\frac{5}{2}^-$	3119		

$$\Delta_2 = 2\Delta_1$$

This relation is badly broken.

$$\frac{\kappa'}{m_c} = 30 \text{ MeV}$$

The HF splittings are **very much deviated** from what we have determined from the excited anti-triplet.

# Scenario II

Assertion: **Three**  $\Omega_c^*$ 's belong to excited sextets, whereas **two**  $\Omega_c^*$ 's with smaller widths are the members of the anti-15plet.

$J$	$S^P$	$M$ [MeV]	$\kappa'/m_c$ [MeV]	$\Delta_J$ [MeV]
0	$\frac{1}{2}^-$	3000	—	—
1	$\frac{1}{2}^-$	3066	24	82
	$\frac{3}{2}^-$	3090		
2	$\frac{3}{2}^-$	3222	input	input
	$\frac{5}{2}^-$	3262	24	164

What about other two  $\Omega_c^*$  s?

- We assume that Omega(3050) and Omega(3119) belong to the **third** rotational excitation of the ground states: They will be then **pentaquarks!**

# Scenario II

Assertion: **Three**  $\Omega_c^*$ 's belong to excited sextets, whereas **two**  $\Omega_c^*$ 's with smaller widths are the members of the anti-15plet.

$J$	$S^P$	$M$ [MeV]	$\kappa'/m_c$ [MeV]	$\Delta_J$ [MeV]
0	$\frac{1}{2}^-$	3000	—	—
1	$\frac{1}{2}^-$	3066	24	82
	$\frac{3}{2}^-$	3090		
2	$\frac{3}{2}^-$	3222	input	input
	$\frac{5}{2}^-$	3262	24	164

$\kappa'/m_c \approx 30$  MeV

What about other two  $\Omega_c^*$  s?

- We assume that  $\Omega_c(3050)$  and  $\Omega_c(3119)$  belong to the **third** rotational excitation of the ground states: They will be then **pentaquarks!**

# Scenario II

Assertion: **Three**  $\Omega_c^*$ 's belong to excited sextets, whereas **two**  $\Omega_c^*$ 's with smaller widths are the members of the anti-15plet.

$J$	$S^P$	$M$ [MeV]	$\kappa'/m_c$ [MeV]	$\Delta_J$ [MeV]
0	$\frac{1}{2}^-$	3000	—	—
1	$\frac{1}{2}^-$	3066	24	82
	$\frac{3}{2}^-$	3090		
2	$\frac{3}{2}^-$	3222	input	input
	$\frac{5}{2}^-$	3262	24	164

$$\Delta_2 = 2\Delta_1$$

This relation is satisfied.

$$\kappa'/m_c \approx 30 \text{ MeV}$$

What about other two  $\Omega_c^*$  s?

- We assume that  $\Omega_c(3050)$  and  $\Omega_c(3119)$  belong to the **third** rotational excitation of the ground states: They will be then **pentaquarks!**

# Scenario II

Assertion: **Three**  $\Omega_c^*$ 's belong to excited sextets, whereas **two**  $\Omega_c^*$ 's with smaller widths are the members of the anti-15plet.

$J$	$S^P$	$M$ [MeV]	$\kappa'/m_c$ [MeV]	$\Delta_J$ [MeV]
0	$\frac{1}{2}^-$	3000	—	—
1	$\frac{1}{2}^-$	3066	24	82
	$\frac{3}{2}^-$	3090		
2	$\frac{3}{2}^-$	3222	input	input
	$\frac{5}{2}^-$	3262	24	164

$$\Delta_2 = 2\Delta_1$$

This relation is satisfied.

Bump structure above 3.2 GeV in the data


$$\kappa'/m_c \approx 30 \text{ MeV}$$

What about other two  $\Omega_c^*$  s?

- We assume that  $\Omega_c(3050)$  and  $\Omega_c(3119)$  belong to the **third** rotational excitation of the ground states: They will be then **pentaquarks!**

# Anti-15plet

\* In the heavy-quark sector, we have yet the third representation, i.e. the anti-15plet.



For the anti-15plet

$$T = 1 \rightarrow J = 1 \quad \text{Combined with a charm quark: } 1 \otimes \frac{1}{2} = \frac{1}{2} \oplus \frac{3}{2} \in \overline{15}$$


In the limit of infinitely heavy quark mass,  $1/2$  &  $3/2$  are degenerate, which will be lifted by a hyperfine interaction.

$$\Omega_c(3050)1/2^+ \quad \Omega_c(3119)3/2^+ : \quad M_{\Omega_c(3/2^+)} - M_{\Omega_c(1/2^+)} \simeq 69 \text{ MeV} !$$

$$\frac{\kappa}{m_c} = (68.1 \pm 1.1) \text{ MeV} \quad \text{in excellent agreement with the ground-state value!}$$

# Anti-15plet

- Exotic anti-15plet naturally arises from the XQSM.




- \* All parameters were fixed in the light baryon sector except for the hyperfine interaction.
- \* Considering almost all theoretical uncertainties, we get the following:  $\mathcal{M}_{\Omega_c} = (3140 - 3370) \text{ MeV}$



# Interpretation of the LHCb data

In the present picture




Resonance	Mass (MeV)	$\Gamma$ (MeV)
$\Omega_c(3000)^0$	$3000.4 \pm 0.2 \pm 0.1^{+0.3}_{-0.5}$	$4.5 \pm 0.6 \pm 0.3$
$\Omega_c(3050)^0$	$3050.2 \pm 0.1 \pm 0.1^{+0.3}_{-0.5}$	$0.8 \pm 0.2 \pm 0.1$
		$<1.2$ MeV, 95% C.L.
$\Omega_c(3066)^0$	$3065.6 \pm 0.1 \pm 0.3^{+0.3}_{-0.5}$	$3.5 \pm 0.4 \pm 0.2$
$\Omega_c(3090)^0$	$3090.2 \pm 0.3 \pm 0.5^{+0.3}_{-0.5}$	$8.7 \pm 1.0 \pm 0.8$
$\Omega_c(3119)^0$	$3119.1 \pm 0.3 \pm 0.9^{+0.3}_{-0.5}$	$1.1 \pm 0.8 \pm 0.4$

$J$	$S^P$	$M$ [MeV]	$\kappa'/m_c$ [MeV]	$\Delta_J$ [MeV]
0	$\frac{1}{2}^-$	3000	—	—
1	$\frac{1}{2}^-$	3066	24	82
	$\frac{3}{2}^-$	3090		
2	$\frac{3}{2}^-$	3222	input	input
	$\frac{5}{2}^-$	3262	24	164

# Interpretation of the LHCb data

In the present picture



$\frac{1}{2}^-$      $\frac{1}{2}^-$      $\frac{3}{2}^-$      $\frac{5}{2}^-$      $\in 6'$


Resonance	Mass (MeV)	$\Gamma$ (MeV)
$\Omega_c(3000)^0$	$3000.4 \pm 0.2 \pm 0.1^{+0.3}_{-0.5}$	$4.5 \pm 0.6 \pm 0.3$
$\Omega_c(3050)^0$	$3050.2 \pm 0.1 \pm 0.1^{+0.3}_{-0.5}$	$0.8 \pm 0.2 \pm 0.1$
$\Omega_c(3066)^0$	$3065.6 \pm 0.1 \pm 0.3^{+0.3}_{-0.5}$	$3.5 \pm 0.4 \pm 0.2$
$\Omega_c(3090)^0$	$3090.2 \pm 0.3 \pm 0.5^{+0.3}_{-0.5}$	$8.7 \pm 1.0 \pm 0.8$
$\Omega_c(3119)^0$	$3119.1 \pm 0.3 \pm 0.9^{+0.3}_{-0.5}$	$1.1 \pm 0.8 \pm 0.4$

<1.2 MeV, 95% C.L.

$J$	$S^P$	$M$ [MeV]	$\kappa'/m_c$ [MeV]	$\Delta_J$ [MeV]
0	$\frac{1}{2}^-$	3000	—	—
1	$\frac{1}{2}^-$	3066	24	82
	$\frac{3}{2}^-$	3090		
2	$\frac{3}{2}^-$	3222	input	input
	$\frac{5}{2}^-$	3262	24	164

# Interpretation of the LHCb data

In the present picture



$\frac{1}{2}^-$   $\frac{1}{2}^-$   $\frac{3}{2}^-$   $\frac{3}{2}^-$   $\frac{5}{2}^-$   $\in 6'$

$\frac{1}{2}^+$   $\frac{3}{2}^+$   $\in \overline{15}$   $\Omega_c(3050)$  &  $\Omega_c(3119)$

Resonance	Mass (MeV)	$\Gamma$ (MeV)
$\Omega_c(3000)^0$	$3000.4 \pm 0.2 \pm 0.1^{+0.3}_{-0.5}$	$4.5 \pm 0.6 \pm 0.3$
$\Omega_c(3050)^0$	$3050.2 \pm 0.1 \pm 0.1^{+0.3}_{-0.5}$	$0.8 \pm 0.2 \pm 0.1$
$\Omega_c(3066)^0$	$3065.6 \pm 0.1 \pm 0.3^{+0.3}_{-0.5}$	$3.5 \pm 0.4 \pm 0.2$
$\Omega_c(3090)^0$	$3090.2 \pm 0.3 \pm 0.5^{+0.3}_{-0.5}$	$8.7 \pm 1.0 \pm 0.8$
$\Omega_c(3119)^0$	$3119.1 \pm 0.3 \pm 0.9^{+0.3}_{-0.5}$	$1.1 \pm 0.8 \pm 0.4$

<1.2 MeV, 95% C.L.

$J$	$S^P$	$M$ [MeV]	$\kappa'/m_c$ [MeV]	$\Delta_J$ [MeV]
0	$\frac{1}{2}^-$	3000	—	—
1	$\frac{1}{2}^-$	3066	24	82
	$\frac{3}{2}^-$	3090		
2	$\frac{3}{2}^-$	3222	input	input
	$\frac{5}{2}^-$	3262	24	164

# How can one falsify the present idea?

PRL **118**, 182001 (2017)

PHYSICAL REVIEW LETTERS

week ending  
5 MAY 2017




## Observation of Five New Narrow $\Omega_c^0$ States Decaying to $\Xi_c^+ K^-$

R. Aaij *et al.*\*

(LHCb Collaboration)

(Received 14 March 2017; published 2 May 2017)


- Anti-15plet consists of **three** Omega\_c's (Isovector baryons).
- The same peaks with the same strength can be found not only in the  $\Xi_c^+ K^-$  channel but also in  $\Xi_c^+ K^0$  and  $\Xi_c^0 K^-$ .



$\Omega_c(3050)$  &  $\Omega_c(3119)$

# LHCb Findings: New five Omega\_cs

	$Y$	$T$	$S^P = \frac{1}{2}^+$	$S^P = \frac{3}{2}^+$
$B_c$	$\frac{5}{3}$	$\frac{1}{2}$	2685	2754
$\Sigma_c$	$\frac{2}{3}$	1	2808	2877
$\Lambda_c$	$\frac{2}{3}$	0	2806	2875
$\Xi_c$	$-\frac{1}{3}$	$\frac{1}{2}$	2928	2997
$\Xi_c^{3/2}$	$-\frac{1}{3}$	$\frac{3}{2}$	2931	3000
$\Omega_c$	$-\frac{4}{3}$	1	3050	3119



$B_c$  baryons will decay *weakly*, if they exist.  
So, they should be stable.

# Decay widths of $\Omega_{cS}$

- Collective operator for the strong vertices in SU(3) symmetric case

$$\mathcal{O}_\varphi = \frac{3}{M_1 + M_2} \sum_{i=1,2,3} \left[ G_0 D_{\varphi i}^{(8)} - G_1 d_{ibc} D_{\varphi b}^{(8)} \hat{S}_c - G_2 \frac{1}{\sqrt{3}} D_{\varphi 8}^{(8)} \hat{S}_i \right] p_i$$

- Decay widths

$$\Gamma_{B_1 \rightarrow B_2 + \varphi} = \frac{1}{2\pi} \langle B_2 | \mathcal{O}_\varphi | B_1 \rangle^2 \frac{M_2}{M_1} p$$

$$G_0 = -\frac{M + M'}{6f_\varphi} a_1$$

$$G_{1,2} = \frac{M + M'}{6f_\varphi} a_{2,3}$$

**No additional  
free parameter!**

$$f_\pi = 93 \text{ MeV}, \quad f_K = 1.2 f_\pi$$

$a_1$	$a_2$	$a_3$
$-3.509 \pm 0.011$	$3.437 \pm 0.028$	$0.604 \pm 0.030$

G. Yang and HChK, PRC **92**, 035206 (2015)

- These parameters  $a_i$  have been determined by the data on hyperon semileptonic decays.

# Decay widths of $\Omega_c S$

- Decay widths of the **charm** baryon sextet

**No additional free parameter!**

#	decay	this work	exp.
1	$\Sigma_c^{++}(\mathbf{6}_1, 1/2) \rightarrow \Lambda_c^+(\bar{\mathbf{3}}_0, 1/2) + \pi^+$	1.93	$1.89^{+0.09}_{-0.18}$
2	$\Sigma_c^+(\mathbf{6}_1, 1/2) \rightarrow \Lambda_c^+(\bar{\mathbf{3}}_0, 1/2) + \pi^0$	2.24	$< 4.6$
3	$\Sigma_c^0(\mathbf{6}_1, 1/2) \rightarrow \Lambda_c^+(\bar{\mathbf{3}}_0, 1/2) + \pi^-$	1.90	$1.83^{+0.11}_{-0.19}$
4	$\Sigma_c^{++}(\mathbf{6}_1, 3/2) \rightarrow \Lambda_c^+(\bar{\mathbf{3}}_0, 1/2) + \pi^+$	14.47	$14.78^{+0.30}_{-0.19}$
5	$\Sigma_c^+(\mathbf{6}_1, 3/2) \rightarrow \Lambda_c^+(\bar{\mathbf{3}}_0, 1/2) + \pi^0$	15.02	$< 17$
6	$\Sigma_c^0(\mathbf{6}_1, 3/2) \rightarrow \Lambda_c^+(\bar{\mathbf{3}}_0, 1/2) + \pi^-$	14.49	$15.3^{+0.4}_{-0.5}$
7	$\Xi_c^+(\mathbf{6}_1, 3/2) \rightarrow \Xi_c(\bar{\mathbf{3}}_0, 1/2) + \pi$	2.35	$2.14 \pm 0.19$
8	$\Xi_c^0(\mathbf{6}_1, 3/2) \rightarrow \Xi_c(\bar{\mathbf{3}}_0, 1/2) + \pi$	2.53	$2.35 \pm 0.22$

Experimental data are taken from the PDG Book.

# Decay widths of $\Omega_{cS}$

- Decay widths of the **bottom** baryon sextet

**No additional free parameter!**

#	decay	this work	exp.
1	$\Sigma_b^+(\mathbf{6}_1, 1/2) \rightarrow \Lambda_b^0(\bar{\mathbf{3}}_0, 1/2) + \pi^+$	6.12	$9.7_{-3.0}^{+4.0}$
2	$\Sigma_b^-(\mathbf{6}_1, 1/2) \rightarrow \Lambda_b^0(\bar{\mathbf{3}}_0, 1/2) + \pi^-$	6.12	$4.9_{-2.4}^{+3.3}$
3	$\Xi_b'(\mathbf{6}_1, 1/2) \rightarrow \Xi_c(\bar{\mathbf{3}}_0, 1/2) + \pi$	0.07	$< 0.08$
4	$\Sigma_b^+(\mathbf{6}_1, 3/2) \rightarrow \Lambda_b^0(\bar{\mathbf{3}}_0, 1/2) + \pi^+$	10.96	$11.5 \pm 2.8$
5	$\Sigma_b^-(\mathbf{6}_1, 3/2) \rightarrow \Lambda_c^0(\bar{\mathbf{3}}_0, 1/2) + \pi^-$	11.77	$7.5 \pm 2.3$
6	$\Xi_b^0(\mathbf{6}_1, 3/2) \rightarrow \Xi_b(\bar{\mathbf{3}}_0, 1/2) + \pi$	0.80	$0.90 \pm 0.18$
7	$\Xi_b^-(\mathbf{6}_1, 3/2) \rightarrow \Xi_b(\bar{\mathbf{3}}_0, 1/2) + \pi$	1.28	$1.65 \pm 0.33$

Experimental data are taken from the PDG Book.



# Decay widths of $\Omega_c$ s

- Decay widths of the **charm** baryon antidecapentaplet

**No additional free parameter!**

#	decay	this work	exp.
	$\Omega_c(\overline{\mathbf{15}}_1, 1/2) \rightarrow \Xi_c(\overline{\mathbf{3}}_0, 1/2) + K$	0.339	—
	$\Omega_c(\overline{\mathbf{15}}_1, 1/2) \rightarrow \Omega_c(\mathbf{6}_1, 1/2) + \pi$	0.097	—
	$\Omega_c(\overline{\mathbf{15}}_1, 1/2) \rightarrow \Omega_c(\mathbf{6}_1, 3/2) + \pi$	0.045	—
9	total	0.48	$0.8 \pm 0.2 \pm 0.1$

Experimental data are taken from the LHCb measurement.

- Note that the widths of  $\Omega_c$ 's are rather small!

# Decay widths of $\Omega_c S$

- Decay widths of the **charm** baryon antidecapentaplet

**No additional free parameter!**

#	decay	this work	exp.
	$\Omega_c(\overline{\mathbf{15}}_1, 3/2) \rightarrow \Xi_c(\overline{\mathbf{3}}_0, 1/2) + K$	0.848	—
	$\Omega_c(\overline{\mathbf{15}}_1, 3/2) \rightarrow \Xi_c(\mathbf{6}_1, 1/2) + K$	0.009	—
	$\Omega_c(\overline{\mathbf{15}}_1, 3/2) \rightarrow \Omega_c(\mathbf{6}_1, 1/2) + \pi$	0.169	—
	$\Omega_c(\overline{\mathbf{15}}_1, 3/2) \rightarrow \Omega_c(\mathbf{6}_1, 3/2) + \pi$	0.096	—
10	total	1.12	$1.1 \pm 0.8 \pm 0.4$

Experimental data are taken from the LHCb measurement.

# Summary

- ★ We have aimed in this talk at how to interpret the newly found five  $\Omega_c$ 's by the LHCb within a mean-field approach (Witten).
- ★ The meson mean fields describe well both the lowest-lying singly heavy baryons and the excited anti-triplet.
- ★ We have predicted **Five** excited sextet and **Two** members in the anti-15plet.

## Suggestions to the LHCb & Belle Collaborations

- \* Can you perform the PWAs to determine the quantum numbers of  $\Omega_c$ 's?
- \* Can you scan channels  $\Xi_c^+ K^0$  and  $\Xi_c^0 K^-$  in the range of the invariant masses between 3050 MeV and 3119 MeV to find isovector  $\Omega_c$ 's?
- I am grateful to M.V. Polyakov, M. Praszalowicz, and Gh.-S. Yang for collaboration over decades.

*Though this be madness,  
yet there is method in it.*

Hamlet Act 2, Scene 2

Thank you very much!