



中国科学院高能物理研究所

**On ρ meson
generalized parton distributions
(GPDs)**

Yubing Dong


Institute of High Energy Physics(IHEP)

Chinese Academy of Sciences(CAS)

Collaborator: (Baodong Sun)

PRD 96 (2017) 036019

Chin.Phys. C 42 (2018) 063104



Outline

1, Introduction

2, Spin 1 particle and basic properties

3, Approach: Light-front constituent quark model

4, Impact parameter space

5, Summary

1, Introduction

Electromagnetic probe

- Electric and magnetic proton form factors
- Proton and Neutron charge distributions
- Nucleon spin structure
- Nucleon-Delta transition (other resonances)
- Quark-hadron duality in structure functions
- Generalized parton distributions
- Pion and deuteron form factors

GPDs (*generalized parton distributions*)

GPDs $H_q(x, \xi, Q^2)$ naturally embody the information of both PDFs and FFs, and therefore display the unique properties to present a “3D” description for a system.

GPDs allow for a unified description of a number of hadronic properties; for example:

(1) In the forward limit they reduce to conventional PDFs

$$H_q(x, 0, 0) = q(x),$$

$$\tilde{H}_q(x, 0, 0) = \Delta q(x).$$

(2) When one integrates GPDs over x they reduce to the usual form factors, e.g. the Dirac form factors^a

$$\sum_q e_q \int dx H_q(x, \xi, t) = F_1(t),$$

$$\sum_q e_q \int dx E_q(x, \xi, t) = F_2(t).$$

GPDs (generalized parton distributions)

GPDs for pion,

Broniowski, PLB 574, PRD78; Choi et al., PRD64; Fanelli, EPJC76;

for nucleon (proton and neutron)

Diehl et al., EPJC 73; Kroll, EPJA53; Pire et al., PRD79; Selyugin, PRD91;.....


Light Nuclei: He-3,...

Rinaldi et al., PRC87.....

Deuteron

Cano et al., PRL87, YBD et al., JPG19,.....

Generalized Parton distributions for pion



Covariant amplitude with a reduced photon vertex for pion GPD (left diagram) and its nonvalence $x < \zeta$ part (right diagram).

Broniowski, PLB574, In the limit of $\xi=0$






Fig. 1. The diagram for the evaluation of the generalized parton distribution of the pion in chiral quark models.

GPDs (generalized parton distributions)

Deep virtual Compton Scattering

[Chueng-Ryong Ji '06, Diehl '16]



A GPD factorization formula:

$$A(\xi, \Delta^2, Q^2) = \sum_i \int_{-1}^1 dx C_i(x, \xi; \log(Q/\mu)) H_i(x, \xi, \Delta^2; \mu)$$

DVCS, TCS, meson production

Parton correlation function:

$$H(k, P, \Delta) = (2\pi)^{-4} \int d^4 z e^{izk} \langle p(P + \frac{1}{2}\Delta) | \bar{q}(-\frac{1}{2}z) \Gamma q(\frac{1}{2}z) | p(P - \frac{1}{2}\Delta) \rangle$$

flavor by flavor

Gauge $A^+=0$


It may be measured by
 Deeply virtual Compton scattering
 Or
 Deeply virtual meson electro-productions

The Dirac matrix Γ selects
 the twist and the parton spin
 degrees of freedom.

$$\Gamma^\mu \rightarrow \gamma^\mu$$

Scheme

[Diehl '16]




2, Spin-1 particle and basic properties

Definition of GPDs (spin -1)

• Unpolarized

[PRL: Berger '01 , for the deuteron]

$$\begin{aligned}
 V_{\lambda'\lambda} &= \frac{1}{2} \int \frac{d\omega}{2\pi} e^{ix(Pz)} \langle p', \lambda' | \bar{q}(-\frac{1}{2}z) \not{n} q(\frac{1}{2}z) | p, \lambda \rangle \Big|_{z=\omega n} \\
 &= \sum_i \epsilon'^{\ast\nu} V_{\nu\mu}^{(i)} \epsilon^\mu H_i^q(x, \xi, t)
 \end{aligned}$$



• Polarized

$$\begin{aligned}
 A_{\lambda'\lambda} &= \frac{1}{2} \int \frac{d\omega}{2\pi} e^{ix(Pz)} \langle p', \lambda' | \bar{q}(-\frac{1}{2}z) \not{n} \gamma_5 q(\frac{1}{2}z) | p, \lambda \rangle \Big|_{z=\omega n} \\
 &= \sum_i \epsilon'^{\ast\nu} A_{\nu\mu}^{(i)} \epsilon^\mu \tilde{H}_i^q(x, \xi, t)
 \end{aligned}$$

$$\mathbf{V}_{\mu\nu} : \{ \mathbf{g}_{\mu\nu}, \mathbf{P}_\mu \mathbf{n}_\nu, \mathbf{P}_\nu \mathbf{n}_\mu, \mathbf{P}_\mu \mathbf{P}_\nu, \mathbf{n}_\mu \mathbf{n}_\nu \}$$

$$\mathbf{A}_{\mu\nu} : \text{Levi-civita } \epsilon_{\mu\nu\alpha\beta} (n^\alpha p^\beta, \dots)$$

Symmetry properties:

$$P = \frac{p' + p}{2}, \quad t = \Delta^2 = (p' - p)^2,$$

$$n^2 = 0, \quad (\text{lightlike four-vector})$$

$$\xi = (n \cdot \Delta) / (n \cdot P), \quad \text{skewness parameter},$$

$$\epsilon = \epsilon(p, \lambda), \quad \epsilon' = \epsilon'(p', \lambda'), \quad \text{polarizations},$$

$$H_i(x, \xi, t) = H_i(x, -\xi, t) \quad (I = 1, 2, 3, 5)$$

$$H_4(x, \xi, t) = -H_4(x, -\xi, t)$$

$$\tilde{H}_i(x, \xi, t) = \tilde{H}_i(x, -\xi, t) \quad (I = 1, 2, 4)$$

$$\tilde{H}_3(x, \xi, t) = -\tilde{H}_3(x, -\xi, t)$$


$$\mathbf{G} \quad H_{\rho+}^d(x, \xi, t) = -H_{\rho+}^u(x, -\xi, t)$$

Sum rules

[Frederico '97, Berger '01, Broniowski '08,]

- Form factor decomposition of Local current

$$\begin{aligned}
 I_{\lambda'\lambda}^\mu &= \langle p', \lambda' | \bar{q}(0) \gamma^\mu q(0) | p, \lambda \rangle \\
 &= \epsilon'^{* \beta} \epsilon^\alpha \left[- \left(G_1^q(t) g_{\beta\alpha} + G_3^q(t) \frac{P_\beta P_\alpha}{2M^2} \right) P^\mu + G_2^q(t) \left(g_\alpha^\mu P_\beta + g_\beta^\mu P_\alpha \right) \right]
 \end{aligned}$$

FFs in flavor


- Sum rules

$$\int_{-1}^1 dx H_i^q(x, \xi, t) = G_i^q(t) \quad (i = 1, 2, 3), \quad G_C(t) = G_1(t) + \frac{2}{3} \eta G_Q(t),$$

$$\int_{-1}^1 dx H_i^q(x, \xi, t) = 0 \quad (i = 4, 5).$$

- Conventional Form factors

$$G_M(t) = G_2(t),$$

$$G_Q(t) = G_1(t) - G_2(t) + (1 + \eta) G_3(t),$$

Forward limit

[Hoodbhoy '89, Berger '01 , Cosyn'17]

- GPDs in forward limit

$$H_1(x, 0, 0) = \frac{q^1(x) + q^{-1}(x) + q^0(x)}{3},$$

$$H_5(x, 0, 0) = q^0(x) - \frac{q^1(x) + q^{-1}(x)}{2},$$

for $x > 0$ $\tilde{H}_1(x, 0, 0) = q_{\uparrow}^1(x) - q_{\uparrow}^{-1}(x)$

- DIS structure functions

$$F_1(x) = \frac{1}{2} \sum_q e_q^2 \frac{q^1(x) + q^{-1}(x) + q^0(x)}{3} + \{q \rightarrow \bar{q}\},$$

$$b_1(x) = \frac{1}{2} \sum_q e_q^2 \left[q^0(x) - \frac{q^1(x) + q^{-1}(x)}{2} \right] + \{q \rightarrow \bar{q}\}$$

$$g_1(x) = \frac{1}{2} \sum_q e_q^2 [q_{\uparrow}^1(x) - q_{\uparrow}^{-1}(x)] + \{q \rightarrow \bar{q}\}.$$

- Single-flavor $F_1^{q\uparrow(\downarrow)}$, $b_1^{q\uparrow(\downarrow)}$

Quark densities:

$$q^\lambda(x) = q_{\uparrow}^\lambda(x) + q_{\downarrow}^\lambda(x)$$

$$q_{\uparrow}^\lambda = q_{\downarrow}^{-\lambda}$$

-H1 and -H5 for $x < 0$, antiquark

Callan-Gross
relation

$$F_1^{q\uparrow(\downarrow)}(x) = \frac{1}{2} H_1^u(x, 0, 0)$$

$$b_1^{q\uparrow(\downarrow)}(x) = \frac{1}{2} H_5^u(x, 0, 0)$$

$$W^{\mu\nu} \sim F_1, F_2, g_1, g_2$$

$$b_1, b_2, b_3, b_4$$

3, Approach: Light-front constituent quark model

Isospin combinations


[Berger '01, Frederico '09 , Bronioski'03]

- **Effective Chiral Lagrangian:**

$$\mathcal{L}_{\rho \rightarrow q\bar{q}} = -i(M/f_\rho)\bar{q}S^\mu \tau q \cdot \rho_\mu = -i(M/f_\rho) \left[\bar{u}S^\mu u \rho_\mu^0 + \sqrt{2}\bar{u}S^\mu d \rho_\mu^+ + \sqrt{2}\bar{d}S^\mu u \rho_\mu^- + \bar{d}S^\mu d \rho_\mu^0 \right]$$

- **Quark field doublets:**

$$q(x) = \begin{pmatrix} u(x) \\ d(x) \end{pmatrix}, \quad \tau_3 q(x) = \begin{pmatrix} u(x) \\ -d(x) \end{pmatrix}$$



- **5 un-polarized GPDs: Isospin combinations**

$$\frac{1}{2} \int \frac{d\lambda}{2\pi} e^{ix(Pz)} \langle \rho^b(p', \lambda') | \bar{q}(-\frac{1}{2}z) \not{n} \tau_3 q(\frac{1}{2}z) | \rho^a(p, \lambda) \rangle \Big|_{z=\lambda n} = i\epsilon_{3ab} \left\{ -(\epsilon'^* \cdot \epsilon) H_{1,\rho^b}^{I=1} \right.$$

$$+ \frac{(\epsilon \cdot n)(\epsilon'^* \cdot P) + (\epsilon'^* \cdot n)(\epsilon \cdot P)}{P \cdot n} H_{2,\rho^b}^{I=1} - \frac{2(\epsilon \cdot P)(\epsilon'^* \cdot P)}{m^2} H_{3,\rho^b}^{I=1}$$

$$\left. + \frac{(\epsilon \cdot n)(\epsilon'^* \cdot P) - (\epsilon'^* \cdot n)(\epsilon \cdot P)}{P \cdot n} H_{4,\rho^b}^{I=1} + \left[m^2 \frac{(\epsilon \cdot n)(\epsilon'^* \cdot n)}{(P \cdot n)^2} + \frac{1}{3}(\epsilon'^* \cdot \epsilon) \right] H_{5,\rho^b}^{I=1} \right\}$$

$H_1 \rightarrow F_1$
 $H_5 \rightarrow b_1$
 $\tilde{H}_1 \rightarrow g_1$

Isospin combinations:


$$H_{i,\rho^\pm}^{I=1}(x, \xi, t) = \frac{1}{2} [H_{i,\rho^\pm}^u(x, \xi, t) - H_{i,\rho^\pm}^d(x, \xi, t)]$$

G parity:

$$H_{\rho^+}^d(x, \xi, t) = -H_{\rho^+}^u(x, -\xi, t)$$

Phenomenological vertex ρ meson

[Choi '04, Frederico '09]



$$x' = \frac{-k_s^+}{p_i^+}$$

$$\kappa_\perp = k_{s\perp} - \frac{k_s^+}{p_i^+} p_{i\perp}$$

Phenomenal vertex:

$$S^\mu = \Gamma^\mu \Lambda(k_s, p)$$

Bethe-Salpeter
amplitude(BSA):

$$\Lambda(k_s, p) = \frac{c}{[k_s^2 - m_R^2 + i\epsilon][(p - k_s)^2 - m_R^2 + i\epsilon]}$$

S-wave

Meson vertex:

$$\Gamma^\mu = \gamma^\mu - \frac{(k_q + k_{\bar{q}})^\mu}{M_0 + 2m}$$

Dispersion relation

Kinematic invariant
mass:

$$M_0^2 = \frac{\kappa_\perp^2 + m^2}{1 - x'} + \frac{\kappa_\perp^2 + m^2}{x'}$$

Residuals


[Choi '04, Frederico '09, Miller'09]

• Six pole (Valence)

$$k_{1(2)}^- = P^- + (k - P)_{on(R)}^- - i \frac{\epsilon}{k^+ - P^+},$$

$$k_{3(4)}^- = \frac{\Delta^-}{2} + (k - \frac{\Delta}{2})_{on(R)}^- - i \frac{\epsilon}{k^+ - \frac{\Delta^+}{2}},$$

$$k_{5(6)}^- = -\frac{\Delta^-}{2} + (k + \frac{\Delta}{2})_{on(R)}^- - i \frac{\epsilon}{k^+ + \frac{\Delta^+}{2}}.$$



• Nonvalance kinematic invariant mass

$$M_{0i(v)}^2 = \frac{\kappa_{\perp}^2 + m^2}{1 - x'} + \frac{\kappa_{\perp}^2 + m^2}{x'}$$

$$\rightarrow \frac{\kappa_{\perp}^2 + m^2}{x' - 1} + \frac{\kappa_{\perp}^2 + m^2}{x'} = M_{0i(nv)}^2$$


$$x = \frac{n \cdot k}{n \cdot P} = \frac{k^+}{P^+}$$

$$x' = \frac{1 - x}{1 - |\xi|}$$

$x \rightarrow 0, 1$ **intrinsic momentum go infinite!**

Non-valance

pair production



The struck u quark in the nonvalance regime, yielded by the off-diagonal terms in the Fock space. The black blob represents the non-wave-function vertex. The red line has the negative sign in this regime.


Results: Form factors $G_{C,M,Q}$

- Form factors

$$G_C(t) = G_1(t) + \frac{2}{3}\eta G_Q(t),$$

$$G_M(t) = G_2(t),$$

$$G_Q(t) = G_1(t) - G_2(t) + (1 + \eta)G_3(t),$$



[Melo '97, Gudino '14]

- low-energy observables

$$G_C(0) = 1,$$

$$G_M(0) = 2M\mu,$$

$$G_Q(0) = M^2 Q_\rho,$$

$$\langle r^2 \rangle = \lim_{t \rightarrow 0} \frac{6[G_C(t) - 1]}{t}.$$

	This work	Melo19 97	Exp. [Gudino20 14]
$\langle r^2 \rangle (\text{fm}^2)$	0.52	0.37	--
μ	2.06	2.19	2.1(5)
$Q_2 (\text{fm}^2)$	0.021	0.050	--

m (constituent mass)	mR (regulator mass)
0.403 GeV	1.61 GeV

TABLE I. The comparison of the results for the magnetic moment μ_ρ (in natural magnetons $e/2M_\rho$) in different approaches.







Model	μ_ρ
This work, mIF RHD	2.16 ± 0.03
Cardarelli, LF RHD [1]	2.26
Melo, LF RHD [2]	2.14
Bakker, LF RHD [3]	2.1
Jaus, LF RHD [4]	1.83
Choi, LF RHD [5]	1.92
He, LF, IF RHD [6]	1.5
He, PF RHD [6]	0.9
Biernat, PF RHD [7]	2.20
Sun, LF CQM [8]	2.06
Hawes, Dyson-Schwinger equation (DSE) [9]	2.69
Ivanov, DSE [10]	2.44
Bhagwat, DSE [11]	2.01
Roberts, DSE [12]	2.11
Pitschmann, DSE [13]	2.11
Carrillo-Serrano, Nambu–Jona-Lasinio model (NJL) [14]	2.59
Luan, NJL [15]	2.1
Samsonov, QCD sum rules [16]	2.0 ± 0.3
Aliev, QCD sum rules [17]	2.4 ± 0.4
Melikhov, LF triangle [18]	2.35
Šimonis, bag model [19]	2.06
Bagdasaryan, relativistic CQM [20]	2.3
Badalian, relativistic Hamiltonian [21]	1.96
Djukanovic, effective field theory [22]	2.24
Andersen, lattice [23]	2.25 ± 0.34
Hedditch, lattice [24]	2.02
Lee, lattice [25]	2.39 ± 0.01
Owen, lattice [26]	2.21 ± 0.08
Lushevskaya, lattice [27]	2.11 ± 0.10
Gudinõ, experiment [28]	2.1 ± 0.5





[Krutov, Polezhaev, and Troitsky, PRD97, 033007]

Form factors $G_{1,2,3}$: (non-)valence contributions

$$|\xi| < \frac{1}{\sqrt{1 - 4M^2/t}}$$



Form factors $G_{1,2,3}$: Nonvalence contributions




The nonvalence contributions to FFs $G_{1,2,3}$ at $\xi = -0.2$ (dotted red line), -0.4 (dashed blue line), and -0.6 (dotted-dashed purple line).

Results: unpolarized GPDs $H_{1,2,3}(x, \xi_0, t)$







$$\xi = \begin{cases} 0 \\ -0.4 \end{cases}$$

$$|\xi| < \frac{1}{\sqrt{1-4M^2/t}}$$



Results: unpolarized GPDs $H_{1,2,3}(x, \xi_0, t_0)$

$$|\xi| < \frac{1}{\sqrt{1-4M^2/t}}$$





$\xi = 0$ (solid black line), -0.2 (dotted red line), -0.4 (dashed blue line), -0.6 (dot-dashed purple line)

Forward limit: Single-flavor F_1^q , b_1^q

$$F_1^{q\uparrow(\downarrow)}(x) = \frac{1}{2} H_1^u(x, 0, 0)$$

$$b_1^{q\uparrow(\downarrow)}(x) = \frac{1}{2} H_5^u(x, 0, 0)$$

$$u_{\rho^+}(x) = \bar{d}_{\rho^+}(1-x)$$



4, Impact Parameter Space

[Burkardt '03, Hoodbhoy '89]

- Spin 1/2

$$\begin{aligned}
 q_N(x, \mathbf{b}) &= |\mathcal{N}|^2 \int \frac{d^2 \mathbf{p}_\perp}{(2\pi)^2} \int \frac{d^2 \mathbf{p}'_\perp}{(2\pi)^2} \\
 &\quad \times \langle p^+, \mathbf{p}'_\perp, \lambda | \left[\int \frac{dz^-}{4\pi} \bar{q}\left(-\frac{z^-}{2}, \mathbf{b}_\perp\right) \gamma^+ q\left(\frac{z^-}{2}, \mathbf{b}_\perp\right) e^{-ixp^+ z^-} \right] | p^+, \mathbf{p}_\perp, \lambda \rangle \\
 &= |\mathcal{N}|^2 \int \frac{d^2 \mathbf{p}_\perp}{(2\pi)^2} \int \frac{d^2 \mathbf{p}'_\perp}{(2\pi)^2} H_q(x, \xi = 0, -(\mathbf{p}_\perp - \mathbf{p}'_\perp)^2) e^{i\mathbf{b}_\perp \cdot (\mathbf{p}_\perp - \mathbf{p}'_\perp)} \\
 &= \int \frac{d^2 \Delta_\perp}{(2\pi)^2} H_q(x, 0, -\Delta_\perp^2) e^{-i\mathbf{b}_\perp \cdot \Delta_\perp}, \quad \Rightarrow
 \end{aligned}$$

$$W_{1/2}^{\mu\nu} \sim \mathbf{F}_1, F_2, g_1, g_2$$

Fourier transformation
Density interpretation

- Spin 1

$$\begin{aligned}
 q(x, \mathbf{b}) &= \int \frac{d^2 \Delta_\perp}{(2\pi)^2} H_1(x, 0, -\Delta_\perp^2) e^{-i\mathbf{b}_\perp \cdot \Delta_\perp} \\
 &= \int_0^\infty \frac{\Delta_\perp d\Delta_\perp}{2\pi} J_0(b\Delta_\perp) H_1(x, 0, -\Delta_\perp^2)
 \end{aligned}$$

$$W_1^{\mu\nu} \sim \mathbf{F}_1, F_2, g_1, g_2$$

$$b_1, b_2, b_3, b_4$$

Impact Parameter Distributions & Gaussian Package (Cut off)

$$\int \frac{d^2 \mathbf{p}_\perp dp^+}{(2\pi)^2 p^+} p^+ \delta(p^+ - p_0^+) G(\mathbf{p}_\perp, \frac{1}{\sigma^2}) |p, \lambda\rangle \sim \int \frac{d^2 \mathbf{p}_\perp}{(2\pi)^2} \exp\left(-\frac{\mathbf{p}_\perp^2 \sigma^2}{2}\right) |p^+, \mathbf{p}_\perp, \lambda\rangle$$

[Diehl '02]

$$q_\sigma(x, b) = \int_0^\infty \frac{\Delta_\perp d\Delta_\perp}{2\pi} J_0(b\Delta_\perp) e^{-\Delta_\perp^2 \sigma^2 / 4} H_1(x, 0, -\Delta_\perp^2)$$

$$q_\sigma(b) = \int_0^1 dx q_\sigma(x, b)$$

$$q(x, \mathbf{b}) = \int \frac{d^2 \Delta_\perp}{(2\pi)^2} H_1(x, 0, -\Delta_\perp^2) e^{-i\mathbf{b}_\perp \cdot \Delta_\perp}$$



$$= \int_0^\infty \frac{\Delta_\perp d\Delta_\perp}{2\pi} J_0(b\Delta_\perp) H_1(x, 0, -\Delta_\perp^2)$$

Only limit value of "t" can be measured

$$q(x, \mathbf{b}, \Delta_0) = \int_0^{\Delta_0} \frac{\Delta_\perp d\Delta_\perp}{2\pi} J_0(b\Delta_\perp) H(x, 0, -\Delta_\perp^2)$$

$$q(\mathbf{b}, \Delta_0) = \int_0^1 dx q(x, \mathbf{b}, \Delta_0)$$

Gaussian Package V.S. Cut off



Impact Parameter Distributions & Form factors

$$\int_{-1}^1 dx H_i^q(x, \xi, t) = G_i^q(t) \quad (i = 1, 2, 3),$$

$$\int_{-1}^1 dx H_i^q(x, \xi, t) = 0 \quad (i = 4, 5),$$

Sum rules



$$G_C(t) = G_1(t) + \frac{2}{3}\eta G_Q(t),$$

$$G_M(t) = G_2(t),$$

$$G_Q(t) = G_1(t) - G_2(t) + (1 + \eta)G_3(t),$$

FFs

$$G_C(t) = \int_{-1}^1 dx \left[H_1(x, \xi, t) + \frac{2}{3}\eta [H_1(x, \xi, t) - H_2(x, \xi, t) + (1 + \eta)H_3(x, \xi, t)] \right],$$

$$G_M(t) = \int_{-1}^1 dx H_2(x, \xi, t),$$

$$G_Q(t) = \int_{-1}^1 dx \left[H_1(x, \xi, t) - H_2(x, \xi, t) + (1 + \eta)H_3(x, \xi, t) \right].$$

IPDs of FFs

$$q_\sigma^{C,M,Q}(\mathbf{b}) = \int_0^1 dx q_\sigma^{C,M,Q}(x, \mathbf{b})$$

$$q_\sigma^{QC}(x, \mathbf{b}) \equiv q_\sigma(x, \mathbf{b}) - q_\sigma^C(x, \mathbf{b})$$


$$q_\sigma^C(x, \mathbf{b}) = \frac{1}{G_C(0)} \int \frac{d^2 \Delta_\perp}{(2\pi)^2} e^{-i\mathbf{b}_\perp \cdot \Delta_\perp - \Delta_\perp^2 \sigma^2} \times \left[H_1(x, 0, -\Delta_\perp^2) + \frac{2}{3} \frac{\Delta_\perp^2}{4M^2} \left[H_1(x, 0, -\Delta_\perp^2) - H_2(x, 0, -\Delta_\perp^2) + \left(1 + \frac{\Delta_\perp^2}{4M^2}\right) H_3(x, 0, -\Delta_\perp^2) \right] \right],$$

$$q_\sigma^M(x, \mathbf{b}) = \frac{1}{G_M(0)} \int \frac{d^2 \Delta_\perp}{(2\pi)^2} e^{-i\mathbf{b}_\perp \cdot \Delta_\perp - \Delta_\perp^2 \sigma^2} H_2(x, 0, -\Delta_\perp^2),$$


$$q_\sigma^Q(x, \mathbf{b}) = \frac{1}{G_Q(0)} \int \frac{d^2 \Delta_\perp}{(2\pi)^2} e^{-i\mathbf{b}_\perp \cdot \Delta_\perp - \Delta_\perp^2 \sigma^2} \left[H_1(x, 0, -\Delta_\perp^2) - H_2(x, 0, -\Delta_\perp^2) + \left(1 + \frac{\Delta_\perp^2}{4M^2}\right) H_3(x, 0, -\Delta_\perp^2) \right]$$

Gaussian Package

$q_\sigma(b)$




(a) $q_\sigma(b)$ (fm^{-2}) with packet width $\sigma = 1 \text{ GeV}^{-1}$.




(b) $q_\sigma(b)$ (fm^{-2}) with packet width $\sigma = 2 \text{ GeV}^{-1}$.

Gaussian package,

$q_\sigma(x,b)$







(a) $q_\sigma^M(x,b)$ with $\sigma = 1 \text{ GeV}^{-1}$ and $x = 1/10, 3/10$ and $1/2$.



(b) $q_\sigma^Q(x,b)$ with $\sigma = 1 \text{ GeV}^{-1}$ and $x = 1/10, 3/10$ and $1/2$.

(color online) The impact parameter dependent FPs $q_\sigma^{M/Q/QC}(x,b)$ with $\sigma = 1 \text{ GeV}^{-1}$ and $x = 1/10, 3/10$ and $1/2$.

Impact Parameter Distributions & Form factors



5, Summary

- GPDs for ρ meson (spin-1)
- Phenomenological approach for ρ meson
- ρ meson FFs / GPDs
- Impact parameter Distribution

o GDAs & $\rho\rho$ production

L3 Collaboration



Exp:

PLUTO/ TASSO/ CELLO/ ARGUS

@ DESY, '82-'91

L3 @ LEP, '03-'06

STAR @ RHIC, '07-'09

Babar @ PEP-II, '08

LHCb, '12 (TeV, double charm)



ARGUS Collaboration etc.

[Albrecht '90, '91]




BABAR

$$\sigma(e^+e^- \rightarrow \rho^+\rho^-) = 8.3 \pm 0.7(\text{stat}) \pm 0.8(\text{syst}) \text{ fb}$$

$\gamma^* \gamma \rightarrow \rho\rho$

- Full reaction: [Anikin '04, '05]

$$2e \rightarrow 2e + \rho^0\rho^0 (\rho^+\rho^-)$$



- @LO (twisit-2), $I = 0$
- charged/neutral cross sec. NOT independent (CG coeffs)
- but charged has bremsstrahlung


- Also related to: [Garcia '15, Kfusek-Gawenda '17, Kumano '17, '18]

$$2e \rightarrow 2e + \rho^0 + 2\pi \quad \rightarrow AA + \pi^+\pi^-\pi^+\pi^-$$

$$\quad \rightarrow 4\pi \quad \quad \quad \rightarrow AA + \pi^+\pi^-2\pi^0$$


GDA (Generalized Distribution Amplitude)

[PRL: Diehl '98, '03, Kumano '17]



$$t \leftrightarrow s$$

$$p', p \leftrightarrow p, -p'$$




(I = 0 @ LO)

$$\Phi_q^{\rho\bar{\rho}}(z, \zeta, W^2) = \int \frac{dx^-}{2\pi} e^{-iz\bar{P}^+x^-} \langle \rho(p)\rho(p') | \bar{q}(x^-) \gamma^+ \begin{pmatrix} 1 \\ \gamma_5 \end{pmatrix} q(0) | 0 \rangle = \begin{pmatrix} \mathbf{V}_q^{\rho\bar{\rho}}(z, \zeta, W^2) \\ \mathbf{A}_q^{\rho\bar{\rho}}(z, \zeta, W^2) \end{pmatrix}$$

$$\Phi_q^{\rho\bar{\rho}}(z, \zeta, W^2) \longleftrightarrow H_\rho^h \left(x = \frac{1-2z}{1-2\zeta}, \xi = \frac{1}{1-2\zeta}, t = W^2 \right)$$

[Kawamura '13, Kumano '17, '18]

$$\gamma\gamma^* \rightarrow \mathbf{f}_0(1370; 0(0^{++})) \rightarrow \rho\rho$$



g_{f_0, ρ^0, ρ^0}



Outlook

- Polarized case
- Double parton distributions (DPDs)
- Deuteron

Thanks!

BACKUP

$$\frac{d\sigma_{ee \rightarrow eep^0\rho^0}}{dW^2} \text{ (only GDA)}$$




- GDA: without ρ width ($W_{min} = 2m_\rho$)
- Exp: Phys.Lett. B568 (2003) 11-22




GDA & Hadronic Tensor @ LO

$$T_{\gamma\gamma^* \rightarrow \rho^0 \rho^0}^{\mu\nu} \sim$$



+



$$\begin{aligned} \mathcal{A}_{(+,+)}^{GDA}(\cos \theta, W^2) \\ = \epsilon_{\mu}^{(+)} \epsilon_{\mu}^{(+)} T_{\gamma\gamma^* \rightarrow \rho^0 \rho^0}^{\mu\nu} \end{aligned}$$