



Uniwersytet
Wrocławski

vBag - a bag model extension with non-perturbative corrections

T.Klahn, T.Fischer, M. Hempel



NATIONAL SCIENCE CENTRE
POLAND

2013/09/B/ST2/01560

QCD Phase Diagram

► dense hadronic matter

HIC in collider experiments


Won't cover the whole diagram

Hot and 'rather' symmetric

NS as a 2nd accessible option

Cold and 'rather' asymmetric

Problem is more complex than
It looks at first gaze



QCD Phase Diagram

► dense hadronic matter

HIC in collider experiments

Won't cover the whole diagram


Hot and 'rather' symmetric

NS as a 2nd accessible option

Cold and 'rather' asymmetric

Problem is more complex than


It looks at first gaze



Neutron Stars

- ▶ Variety of scenarios regarding inner structure: with or without QM
- ▶ Question whether/how QCD phase transition occurs is not settled
- ▶ Most honest approach: take both (and more) scenarios into account and compare to available data

Hybrid Star




Inner Crust

- heavy ions
- relativistic electron gas
- superfluid neutrons

Inner Core

- (neutrons, protons)
- electrons, muons
- hyperons
- bosonic condensates
- deconfined quark matter

Neutron Star



Outer Crust


- ions
- electron gas

Core

- neutrons, protons
- electrons, muons
- superconducting protons


- strange quark matter

Strange Star



Neutron Stars = Quark Cores?

Hybrid Star




Inner Crust

- heavy ions
- relativistic electron gas
- superfluid neutrons

Inner Core

- (neutrons, protons)
- electrons, muons
- hyperons
- bosonic condensates
- deconfined quark matter

Neutron Star



Outer Crust


- ions
- electron gas

Core

- neutrons, protons
- electrons, muons
- superconducting protons

- strange quark matter


Strange Star



Neutron Stars = Quark Cores?



Hybrid Star




Inner Crust

- heavy ions
- relativistic electron gas
- superfluid neutrons

Inner Core

- (neutrons, protons)
- electrons, muons
- hyperons
- bosonic condensates
- deconfined quark matter

Neutron Star




Outer Crust

- ions
- electron gas


Core

- neutrons, protons
- electrons, muons
- superconducting protons
- strange quark matter

Strange Star



Neutron Stars = Quark Cores?



Neutron Stars = Quark Cores?




Hybrid Star



Neutron Star



Strange Star




Inner Crust

- heavy ions
- relativistic electron gas
- superfluid neutrons

Inner Core

- (neutrons, protons)
- electrons, muons
- hyperons
- bosonic condensates
- deconfined quark matter




Outer Crust

- ions
- electron gas

Core

- neutrons, protons
- electrons, muons
- superconducting protons
- strange quark matter




1st order phase transition observable in neutrino signal

High mass NSs do not rule out QM cores

They are no evidence neither.

General problem:
Which observable would convince that QCD phase transition happens in nature?

Sagert, Fischer et al. →
(PRL 2010)



1st neutrino burst shortly after core bounce, **deleptonization burst**, standard feature in all supernova models

Hard to detect because it comes in ν_e

2nd burst due to 2nd-shock propagation across neutrinospheres, dominated by:

$$\bar{\nu}_e \quad (\nu_{\mu/\tau}, \bar{\nu}_{\mu/\tau})$$

Neutrinos are emitted locally and come from low densities (hadronic phase)

2nd burst last only few milliseconds

Accompanied by significant rise of average neutrino energies

Observable for currently operating neutrino-detector facilities

Problem: Violation of current constraints from astrophysics

High mass NSs do not rule out QM cores

Demorest et al. (2010), Nature 09466, **J1614-2230**

Antoniadis et al. (2013), Science 340, 448, **J1614-2230**

They are no evidence neither. Steiner et al. (2010), ApJ 722 (Bayesian analysis of few selected low-mass X-ray binary systems)


General problem:

Which observable would convince that QCD phase transition happens in nature?

Sagert, Fischer et al. →
(PRL 2010)

$$M_{\max} = (1.97 - 2.01) \pm 0.04 M_{\odot}$$

$$R|_{M=1.4 M_{\odot}} = 12 \pm 1 \text{ km}$$



All quark-bag hybrid EOS tested are ruled out !

Quark Matter

What is so special about quarks?

Confinement: No isolated quark has ever been observed
Quarks are confined in baryons and mesons

Dynamical Mass Generation:
Proton 940 MeV, 3 constituent quarks with each 5 MeV
→ 98.4% from somewhere?

and then this:
eff. quark mass in proton: $940 \text{ MeV}/3 \approx 313 \text{ MeV}$
eff. quark mass in pion : $140 \text{ MeV}/2 = 70 \text{ MeV}$

quark masses generated by interactions only
,out of nothing'
interaction in QCD through (self interacting) gluons
dynamical chiral symmetry breaking (DCSB)
is a distinct nonperturbative feature!

Confinement and DCSB are connected. Not trivially seen from QCD Lagrangian.
Investigating quark-hadron phase transition requires nonperturbative approach.

Quark Matter

Confinement and DCSB are features of QCD.

It would be too nice to account for these phenomena
when describing QM in Compact Stars...

Current approaches mainly used to describe dense, deconfined QM:

Bag-Model :

While Bag-models certainly account for confinement (constructed to do exactly this) Chodos, Jaffe et al: Baryon Structure (1974)
they do not exhibit DCSB (quark masses are fixed - bare quark masses). Farhi, Jaffe: Strange Matter (1984)

NJL-Model :

While NJL-type models certainly account for DCSB (applied, because they do)
they do not (trivially) exhibit confinement.

Nambu, Jona-Lasinio (1961)

Modifications to address confinement exist (e.g. PNJL) but are not entirely satisfying

Both models: **Inspired by, but not originally based on QCD.**

Lattice QCD still fails at T=0 and finite μ


Dyson-Schwinger Approach

Derive gap equations from QCD-Action. Self consistent self energies.

Successfully applied to describe meson and baryon properties

Extension from vacuum to finite densities desirable

→ EoS within QCD framework



Quark Matter

Confinement and DCSB are features of QCD.

It would be too nice to account for these phenomena
when describing QM in Compact Stars...

Current approaches mainly used to describe dense, deconfined QM:

Bag-Model :

While Bag-models certainly account for confinement (constructed to do exactly this) Chodos, Jaffe et al: Baryon Structure (1974)
they do not exhibit DCSB (quark masses are fixed - bare quark masses). Farhi, Jaffe: Strange Matter (1984)

NJL-Model :

While NJL-type models certainly account for DCSB (applied, because they do)
they do not (trivially) exhibit confinement.

Nambu, Jona-Lasinio (1961)

Modifications to address confinement exist (e.g. PNJL) but are not entirely satisfying

Both models: **Inspired by, but not originally based on QCD.**

Lattice QCD still fails at T=0 and finite μ

Dyson-Schwinger Approach


Derive gap equations from QCD-Action. Self consistent self energies.

Successfully applied to describe meson and baryon properties


Extension from vacuum to finite densities desirable

→ EoS within QCD framework

→ **THIS TALK: Bag and NJL model as simple limits within DS approach**



DSE : dynamical, momentum dependent mass generation



momentum dep. (here @ $T=\mu=0$)
LQCD as benchmark

Neither NJL nor BAG have this

How do momentum dependent
gap solutions affect

- EoS of deconfined quark matter?
- EoS of confined quark matter?
- transport properties in medium?

Roberts (2011)
Bhagwat et al. (2003,2006,2007)
P. O. Bowman et al. (2005)

Bag model: bare quark mass at all momenta and densities

NJL model: dressed quark mass at all momenta, changing dynamically with chemical potential


Dyson-Schwinger Perspective

One particle gap equation(s)

$$S^{-1}(p; \mu) = i\vec{\gamma}\vec{p} + i\gamma_4(p_4 + i\mu) + m + \Sigma(p; \mu)$$

Self energy -> entry point for simplifications

$$\Sigma(p; \mu) = \int_{\Lambda} \frac{d^4 q}{(2\pi)^4} g^2 D_{\rho\sigma}(p - q) \gamma_{\rho} \frac{\lambda^a}{2} S(q) \Gamma_{\sigma}^a(p; q)$$



General (in-medium) gap solutions

$$S^{-1}(p; \mu) = i\vec{\gamma}\vec{p}A(p; \mu) + i\gamma_4(p_4 + i\mu)C(p; \mu) + B(p; \mu)$$

Effective gluon propagator

$$S(p;\mu)^{-1} = Z_2(i\vec{\gamma}\vec{p} + i\gamma_4(p_4 + i\mu) + m_{\text{bm}}) + \Sigma(p;\mu)$$

$$\Sigma(p;\mu) = Z_1 \int_q^\Lambda g^2(\mu) D_{\rho\sigma}(p-q;\mu) \frac{\lambda^a}{2} \gamma_\rho S(q;\mu) \Gamma_\sigma^a(q,p;\mu)$$

Ansatz for self energy (rainbow approximation, effective gluon propagator(s))

$$Z_1 \int_q^\Lambda g^2 D_{\mu\nu}(p-q) \frac{\lambda^a}{2} \gamma_\mu S(q) \Gamma_\nu^a(q,p) \rightarrow \int_q^\Lambda \mathcal{G}((p-q)^2) D_{\mu\nu}^{\text{free}}(p-q) \frac{\lambda^a}{2} \gamma_\mu S(q) \frac{\lambda^a}{2} \gamma_\nu$$

Specify behaviour of $\mathcal{G}(k^2)$

$$\frac{\mathcal{G}(k^2)}{k^2} = 8\pi^4 D \delta^4(k) + \frac{4\pi^2}{\omega^6} D k^2 e^{-k^2/\omega^2} + 4\pi \frac{\gamma_m \pi}{\frac{1}{2} \ln \left[\tau + \left(1 + k^2/\Lambda_{\text{QCD}}^2 \right)^2 \right]} \mathcal{F}(k^2)$$

Infrared strength running coupling for large k (zero width + finite width contribution)

EoS (finite densities):

1st term (Munczek/Nemirowsky (1983))

2nd term

NJL model:

$$g^2 D_{\rho\sigma}(p-q) = \frac{1}{m_G^2} \delta_{\rho\sigma}$$

delta function in momentum space → Klähn et al. (2010)

→ Chen et al.(2008,2011)

delta function in configuration space = const. In mom. space

Munczek/Nemirovsky -> NJL's complement

Wigner Phase $\frac{\mathcal{G}(k^2)}{k^2} = 8\pi^4 D \delta^4(k) + \frac{4\pi^2}{\omega^6} D k^2 e^{-k^2/\omega^2} + 4\pi \frac{\gamma_m \pi}{\frac{1}{2} \ln \left[\tau + \left(1 + k^2/\Lambda_{\text{QCD}}^2 \right)^2 \right]} \mathcal{F}(k^2)$

$B_W = 0, A_W = C_W$:

$$C_W(p, \mu) = \frac{1}{2} \left(1 + \sqrt{1 + \frac{2\eta^2}{p_3^2 + (p_4 + i\mu)^2}} \right)$$

Nambu Phase

$A_N = C_N$.
 $\Re(\tilde{p}^2) < \frac{\eta^2}{4}$:

$$B_N(p, \mu) = \sqrt{\eta^2 - 4(p_3^2 + (p_4 + i\mu)^2)}$$

$$C_N(p, \mu) = 2$$

$\Re(\tilde{p}^2) > \frac{\eta^2}{4}$:

$$A_N = A_W, B_N = B_W, C_N = C_W.$$



MN antithetic to NJL


NJL: contact interaction in x

MN: contact interaction in p
(background field in x)


Munczek/Nemirovsky

$$f_1(|\vec{p}|; \mu) = \frac{1}{4\pi} \int_{-\infty}^{\infty} dp_4 \text{tr}_D(-\gamma_4) S(p; \mu)$$

Wigner Phase



$$P(\mu < \eta) = P_0 + \int_0^\mu d\mu' n(\mu') \propto P_0 + \text{const} \times \mu^5$$



$\mu^2 \geq 2\eta^2$ to obtain

$f_1(\vec{p}^2 = 0) = 1$ model is scale invariant regarding μ/η
 $P(\mu) \propto \mu^5$ well satisfied up to $\mu/\eta \approx 1$

($\eta = 1.09 \text{ GeV}$)


'small' chem. Potential: $f_1(\vec{p}^2 = 0, \mu < \eta) \propto \mu \leftarrow$

$$n(\mu < \eta) = \frac{2N_c N_f}{2\pi^2} \int d^3 \vec{p} f_1(|\vec{p}|) \propto \mu^4$$

DSE – simple effective gluon coupling

$$\frac{\mathcal{G}(k^2)}{k^2} = 8\pi^4 D \delta^4(k) + \frac{4\pi^2}{\omega^6} D k^2 e^{-k^2/\omega^2} + 4\pi \frac{\gamma_m \pi}{\frac{1}{2} \ln \left[\tau + \left(1 + k^2/\Lambda_{\text{QCD}}^2 \right)^2 \right]} \mathcal{F}(k^2)$$

Wigner Phase Less extreme, but again, 1 particle number density distribution different from free Fermi gas (quasi particle) distribution




DSE -> NJL model

$$g^2 D_{\rho\sigma}(p - q) = \frac{1}{m_G^2} \delta_{\rho\sigma},$$

Gluon contact interaction in configuration space (other models exist)

$$\Gamma_\rho^a(p; q) = \frac{\lambda^a}{2} \gamma_\rho.$$

Rainbow approximation



$$A = 1$$

$$\vec{p}^2 A_p = \vec{p}^2 + \frac{8N_c}{9m_G^2} \int_{\Lambda} \frac{d^4 q}{(2\pi)^4} \frac{\vec{p} \vec{q} A_q}{\vec{q}^2 A_q^2 + \tilde{q}_4^2 C_q^2 + B_q^2},$$

$$B_p = m + \frac{16N_c}{9m_G^2} \int_{\Lambda} \frac{d^4 q}{(2\pi)^4} \frac{B_q}{\vec{q}^2 A_q^2 + \tilde{q}_4^2 C_q^2 + B_q^2},$$

$$B_\mu = m + \frac{4N_c}{9m_G^2} n_s(T, \mu^*, B),$$

$$\tilde{p}_4^2 C_p = \tilde{p}_4^2 + \frac{8N_c}{9m_G^2} \int_{\Lambda} \frac{d^4 q}{(2\pi)^4} \frac{\tilde{p}_4 \tilde{q}_4 C_q}{\vec{q}^2 A_q^2 + \tilde{q}_4^2 C_q^2 + B_q^2},$$

$$\mu = \mu^* - \frac{2N_c}{9m_G^2} n_v(T, \mu^*, B),$$

$$\tilde{p}_4 C = p_4 + i(\mu + \omega_\mu) \equiv \hat{p}_4$$

Thermodynamical Potential

DS: steepest descent

$$P[S] = \text{Tr} \ln[S^{-1}] - \frac{1}{2} \text{Tr}[\Sigma S].$$

$$P_{FG} = \text{Tr} \ln S^{-1} = 2N_c \int_{\Lambda} \frac{d^4 p}{(2\pi)^4} \ln(\vec{p}^2 + \hat{p}_4^2 + B_{\mu}^2)$$

$$P_I = -\frac{1}{2} \text{Tr} \Sigma S = \frac{3}{4} m_G^2 \omega_{\mu}^2 - \frac{3}{8} m_G^2 \phi_{\mu}^2$$

Compare to NJL type model with following Lagrangian (interaction part only):

$$\mathcal{L}_I = \mathcal{L}_S + \mathcal{L}_V = G_s \sum_{a=0}^8 (\bar{q} \tau_a q)^2 + G_v (\bar{q} i \gamma_0 q)^2.$$

$$\Omega_q = \Omega_q^0 + \frac{\phi^2}{4G_s} - \frac{\omega^2}{2G_v} - \Omega_q(T = \mu = 0)$$

$$\phi_{\mu} = 2G_s N_c n_s(T, m_f^*, \mu_f^*)$$

$$\omega_{\mu} = -2G_s N_c n_v(T, m_f^*, \mu_f^*)$$

$$\frac{\partial \Omega_q}{\partial \phi_{\mu}} = \frac{\partial \Omega_q}{\partial \omega_{\mu}} = 0.$$

Thermodynamical Potential

DS: steepest descent

$$P[S] = \text{Tr} \ln[S^{-1}] - \frac{1}{2} \text{Tr}[\Sigma S].$$

NJL model is easily understood
as a particular approximation
of QCD's DS gap equations

$$P_{FG} = \text{Tr} \ln S^{-1} = 2N_c \int_{\Lambda} \frac{d^4 p}{(2\pi)^4} \ln(\vec{p}^2 + \hat{p}_4^2 + B_{\mu}^2)$$

$$P_I = -\frac{1}{2} \text{Tr} \Sigma S = \frac{3}{4} m_G^2 \omega_{\mu}^2 - \frac{3}{8} m_G^2 \phi_{\mu}^2$$

Compare to NJL type model with following Lagrangian (interaction part only):

$$\mathcal{L}_I = \mathcal{L}_S + \mathcal{L}_V = G_s \sum_{a=0}^8 (\bar{q} \tau_a q)^2 + G_v (\bar{q} i \gamma_0 q)^2.$$

$$\Omega_q = \Omega_q^0 + \frac{\phi^2}{4G_s} - \frac{\omega^2}{2G_v} - \Omega_q(T = \mu = 0)$$

$$\phi_{\mu} = 2G_s N_c n_s(T, m_f^*, \mu_f^*)$$

$$\omega_{\mu} = -2G_s N_c n_v(T, m_f^*, \mu_f^*)$$

$$\frac{\partial \Omega_q}{\partial \phi_{\mu}} = \frac{\partial \Omega_q}{\partial \omega_{\mu}} = 0.$$

ADJUSTING THE QUARK MATTER EOS



Effective Lagrangian

$$\begin{aligned}\mathcal{L}_{int} = & G_S \eta_D \sum_{a,b=2,5,7} (\bar{q} i \gamma_5 \tau_a \lambda_b C \bar{q}^T) (q^T C i \gamma_5 \tau_a \lambda_a q) \\ & + G_S \sum_{a=0}^8 [(\bar{q} \tau_a q)^2 + \eta_V (\bar{q} i \gamma_0 q)^2]\end{aligned}$$

Thermodynamical potential

$$\begin{aligned}\Omega(T, \mu) = & \frac{\phi_u^2 + \phi_d^2 + \phi_s^2}{8G_S} + \frac{\Delta_{ud}^2 + \Delta_{us}^2 + \Delta_{ds}^2}{4G_D} \\ & - \int \frac{d^3 p}{(2\pi)^3} \sum_{n=1}^{18} \left[E_n + 2T \ln \left(1 + e^{-E_n/T} \right) \right] + \Omega_{Iep} - \Omega_0\end{aligned}$$

ADJUSTING THE QUARK MATTER EOS





- » Large Mass ($\sim 2 M_{\odot}$) and radius ($R \geq 12 \text{ km}$) \Rightarrow stiff quark matter EoS;
Note: DU problem of DBHF removed by deconfinement! **and:** CFL core Hybrids unstable!
- » Flow in Heavy-Ion Collisions \Rightarrow not too stiff EoS !
Note: Quark matter removes violation by DBHF at high densities

T. Klähn et al., PLB 654, 170 (2007); arxiv:nucl-th/0609067

A phase transition softens the equation of state! **VERY GOOD!!! Solves some problems.**

ADJUSTING THE QUARK MATTER EOS




Doesn't look very systematic

- » Large Mass ($\sim 2 M_{\odot}$) and radius ($R \geq 12$ km) \Rightarrow stiff quark matter EoS;
Note: DU problem of DBHF removed by deconfinement! **and:** CFL core Hybrids unstable!
- » Flow in Heavy-Ion Collisions \Rightarrow not too stiff EoS !
Note: Quark matter removes violation by DBHF at high densities

T. Klähn et al., PLB 654, 170 (2007); arxiv:nucl-th/0609067


A phase transition softens the equation of state! **VERY GOOD!!!** Solves some problems.

ADJUSTING THE QUARKYONIC ISLAND



ADJUSTING THE QUARK RADIUS

no stable hybrids
PT at too large n




ADJUSTING THE QUARK ONSET MASS

no stable hybrids

PT at too large n

Quark Stars?

PT at too small n



ADJUSTING THE QUARK ONSET

no stable hybrids


PT at too large n

Quark Stars?

PT at too small n

PSR J1614

$$M_{\max} = (1.97 \pm 0.04) M_{\text{sun}}$$



ADJUSTING THE QUARK ONSET

no stable hybrids


PT at too large n

Quark Stars?

PT at too small n

PSR J1614

$$M_{\max} = (1.97 \pm 0.04) M_{\text{sun}}$$



ADJUSTING THE QUARK CORE SIZE

no stable hybrids


PT at too large n

Quark Stars?

PT at too small n

PSR J1614

$$M_{\max} = (1.97 \pm 0.04) M_{\text{sun}}$$



ADJUSTING THE QUARK ONSET

no stable hybrids

PT at too large n

Quark Stars?

PT at too small n

PSR J1614

$$M_{\max} = (1.97 \pm 0.04) M_{\text{sun}}$$

Comparison with
sym. matter: $n_{\text{onset}}^{\text{sym}}$


QM in PSR J1614?

Yes $\rightarrow n_{\text{onset}}^{\text{sym}} \leq 4n_{\text{sat}}$

No $\rightarrow n_{\text{onset}}^{\text{sym}} > 4n_{\text{sat}}$



It is remarkable that
this critical value is
rather constant!

Still: For this model...



NJL MODEL STUDY FOR NS

(TK, R.ŁASTOWIECKI, D.BLASCHKE, PRD **88**, 085001 (2013))



Conclusion: NS may or may not support a significant QM core.
additional interaction channels won't change this if coupling strengths are not precisely known.


1st order phase transition observable in neutrino signal

High mass NSs do not rule out QM cores

They are no evidence neither.

General problem:
Which observable would convince that QCD phase transition happens in nature?

Sagert, Fischer et al. →
(PRL 2010)



Standard feature: **deleptonization burst** form core bounce

Signal from strong 1st order phase transition at high densities

1st neutrino burst shortly after core bounce, **deleptonization burst**, standard feature in all supernova models

Hard to detect because it comes in ν_e

2nd burst due to 2nd-shock propagation across neutrinospheres, dominated by:

$$\bar{\nu}_e \quad (\nu_{\mu/\tau}, \bar{\nu}_{\mu/\tau})$$

Neutrinos are emitted locally and come from low densities (hadronic phase)

2nd burst last only few milliseconds

Accompanied by significant rise of average neutrino energies

Observable for currently operating neutrino-detector facilities

Problem: Violation of current constraints from astrophysics

High mass NSs do not rule out QM cores

Demorest et al. (2010), Nature 09466, **J1614-2230**

Antoniadis et al. (2013), Science 340, 448, **J1614-2230**

They are no evidence neither. Steiner et al. (2010), ApJ 722 (Bayesian analysis of few selected low-mass X-ray binary systems)


General problem:

Which observable would convince that QCD phase transition happens in nature?

Sagert, Fischer et al. →
(PRL 2010)

$$M_{\max} = (1.97 - 2.01) \pm 0.04 M_{\odot}$$

$$R|_{M=1.4 M_{\odot}} = 12 \pm 1 \text{ km}$$



All quark-bag hybrid EOS tested are ruled out !

Thermodynamical Potential

DS: steepest descent

$$P[S] = \text{Tr} \ln[S^{-1}] - \frac{1}{2} \text{Tr}[\Sigma S].$$

NJL model is easily understood
as a particular approximation
of QCD's DS gap equations

$$P_{FG} = \text{Tr} \ln S^{-1} = 2N_c \int_{\Lambda} \frac{d^4 p}{(2\pi)^4} \ln(\vec{p}^2 + \hat{p}_4^2 + B_{\mu}^2)$$

$$P_I = -\frac{1}{2} \text{Tr} \Sigma S = \frac{3}{4} m_G^2 \omega_{\mu}^2 - \frac{3}{8} m_G^2 \phi_{\mu}^2$$

Compare to NJL type model with following Lagrangian (interaction part only):

$$\mathcal{L}_I = \mathcal{L}_S + \mathcal{L}_V = G_s \sum_{a=0}^8 (\bar{q} \tau_a q)^2 + G_v (\bar{q} i \gamma_0 q)^2.$$

$$\Omega_q = \Omega_q^0 + \frac{\phi^2}{4G_s} - \frac{\omega^2}{2G_v} - \Omega_q(T = \mu = 0)$$

$$\phi_{\mu} = 2G_s N_c n_s(T, m_f^*, \mu_f^*)$$

$$\omega_{\mu} = -2G_s N_c n_v(T, m_f^*, \mu_f^*)$$


$$\frac{\partial \Omega_q}{\partial \phi_{\mu}} = \frac{\partial \Omega_q}{\partial \omega_{\mu}} = 0.$$

Bag Model from NJL perspective

(TK, T.Fischer, ApJ , 2015)

obvious differences between NJL and Bag:

- $D\chi SB$
- confinement
- vector interaction




Bag Model from NJL perspective

(TK, T.Fischer, ApJ , 2015)

obvious differences between NJL and Bag:


- $D\chi SB$
- confinement
- vector interaction



Bag Model from NJL perspective

obvious differences between NJL and Bag:

- $D\chi SB$
- **confinement**
- vector interaction



confinement

Pressure Quark NJL/Bag


Pressure Nuclear Matter

Pressure not zero at χ transition

Bag Model from NJL perspective

obvious differences between NJL and Bag:

- $D\chi_{SB}$
- **confinement**
- vector interaction



confinement

Pressure Quark NJL/Bag

Pressure Nuclear Matter

Pressure not zero at χ transition
Reduce χ bag pressure to match
to nuclear EoS



(Pagliara&Schaffner-Bielich PRD, 2008)

Bag Model from NJL perspective

obvious differences between NJL and Bag:

- D χ SB
- confinement
- **vector interaction**

$$B_\mu = m + \frac{4N_c}{9m_G^2} n_s(T, \mu^*, B),$$
$$\mu = \mu^* - \frac{2N_c}{9m_G^2} n_v(T, \mu^*, B),$$



vBag: vector interaction enhanced bag model

Chiral + Vector:

$$P_{BM}^i(\mu_i) = P_{kin}(\mu_i^*) + \frac{K_v}{2} n_v^2(\mu_i^*) - P_{BAG}^i$$

DCSB

$$\varepsilon_{BM}^i(\mu_i) = \varepsilon_{kin}(\mu_i^*) + \frac{K_v}{2} n_v^2(\mu_i^*) + P_{BAG}^i$$

Vector Interaction

$$\mu_i = \mu_i^* + K_v n_v(T, \mu_i^*)$$

'Confinement':


$$P = \sum_f P_f^{kin} - B_{eff} \text{ with } B_{eff} = \sum_f B_\chi^f - B_{dc}$$

Confinement
(needs hadron EoS!)


And, of course, chiral+vector+'confinement'

(Klahn & Fischer [arXiv:1503.07442](https://arxiv.org/abs/1503.07442) ApJ 2015)

Neutron Stars with QM core – vBAG vs BAG



Neutron Stars with QM core – vBAG vs BAG



Conclusions Part I

Vector enhanced bag like model can be motivated from NJL - which can be obtained from DS gap equations

Bag model character: bare quark masses
effective bag pressure

Difference: chiral bag pressure as consequence of D χ SB, flavor dependence
confining bag pressure with opposite sign (binding energy)
accounts for vector interaction -> stiff EoS, promising for astrophysical applications

What NJL couldn't: reduced chiral bag pressure due to confinement -> by hand, no harm to td consistence

Advantage of the model: extremely simple to use, no regularization required, Fermi gas expressions, bare masses
no (obvious) gap equation

$$P_{BM}^i(\mu_i) = P_{kin}(\mu_i^*) + \frac{K_v}{2} n_v^2(\mu_i^*) - P_{BAG}^i \quad P = \sum_f P_f^{kin} - B_{eff} \text{ with } B_{eff} = \sum_f B_X^f - B_{dc}$$
$$\varepsilon_{BM}^i(\mu_i) = \varepsilon_{kin}(\mu_i^*) + \frac{K_v}{2} n_v^2(\mu_i^*) + P_{BAG}^i$$

$$\mu_i = \mu_i^* + K_v n_v(T, \mu_i^*)$$

Absolutely Stable Strange Matter?

(very) brief review:

Three essential papers:

1

PHYSICAL REVIEW D

VOLUME 9, NUMBER 12

15 JUNE 1974

New extended model of hadrons*

A. Chodos, R. L. Jaffe, K. Johnson, C. B. Thorn, and V. F. Weisskopf
*Laboratory for Nuclear Science and Department of Physics,
Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*
(Received 25 March 1974)

We propose that a strongly interacting particle is a finite region of space to which fields are confined. The confinement is accomplished in a Lorentz-invariant way by endowing the finite region with a constant energy per unit volume, B . We call this finite region a "bag." The contained fields may be either fermions or bosons and may have any spin; they may or may not be coupled to one another. Equations of motion and boundary conditions are obtained from a variational principle. The confining region has no dynamical freedom but constrains the fields inside: There are no excitations of the coordinates determining the confining region. The model possesses many desirable features of hadron dynamics: (i) a parton




FIG. 1. A color-singlet bag attempting to fission into two bags which are not color singlets. The flux lines of the colored gluon field are shown explicitly.

Key assumptions: Bag is a given, massless colored quark and gluon fields, boundary conditions ensure confinement

Absolutely Stable Strange Matter?

(very) brief review:

Three essential papers:

2

PHYSICAL REVIEW D

VOLUME 30, NUMBER 2

15 JULY 1984

Cosmic separation of phases

Edward Witten*

Institute for Advanced Study, Princeton, New Jersey 08540

(Received 9 April 1984)

A first-order QCD phase transition that occurred reversibly in the early universe would lead to a surprisingly rich cosmological scenario. Although observable consequences would not necessarily survive, it is at least conceivable that the phase transition would concentrate most of the quark excess in dense, invisible quark nuggets, providing an explanation for the dark matter in terms of QCD effects only. This possibility is viable only if quark matter has energy per baryon less than 938 MeV. Two related issues are considered in appendices: the possibility that neutron stars generate a quark-matter component of cosmic rays, and the possibility that the QCD phase transition may have produced a detectable gravitational signal.

The average quark kinetic energy is proportional to μ , so (with a common pressure in the two cases) it is smaller in the three-flavor case by a factor

$$\tilde{\mu}/(\frac{1}{3}\mu + \frac{2}{3}2^{1/3}\mu) = [3/(1+2^{4/3})]^{3/4} \simeq 0.89 .$$

strange-quark mass will reduce this effect, but it is still plausible that strange quarks lower the energy per baryon of quark matter by 50–70 MeV per baryon. This is

Absolutely Stable Strange Matter?

(very) brief review:

Three essential papers:

3

PHYSICAL REVIEW D

VOLUME 30, NUMBER 11

1 DECEMBER 1984

Strange matter

Edward Farhi and R. L. Jaffe

*Center for Theoretical Physics, Laboratory for Nuclear Science and Department of Physics,
Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*

(Received 9 May 1984)

We explore the properties of quark matter in equilibrium with the weak interactions, containing comparable numbers of up, down, and strange quarks. Witten has recently conjectured that this “strange matter” may be absolutely stable. Using a Fermi-gas model including $O(\alpha_c)$ corrections we establish the conditions under which strange matter in bulk is stable and describe its characteristics. Augmenting our model with surface-tension and Coulomb effects we study strange matter with intermediate baryon number, $10^2 \lesssim A \lesssim 10^7$. For low baryon numbers $A \lesssim 10^2$, we replace the Fermi gas by the bag model and study shell effects and the approach to the bulk limit. Finally, we discuss the phenomenology of strange matter in all its forms.

Absolutely Stable Strange Matter?

(very) brief review:

Three essential papers:

3

PHYSICAL REVIEW D

VOLUME 30, NUMBER 11

1 DECEMBER 1984

Strange matter


In Sec. II, we investigate the properties of stable strange matter in bulk. Our study rests on several plausible assumptions. The first, as we have already mentioned, is that the system is well approximated by a Fermi gas separated from the vacuum by a phase boundary. We further assume that the effects of dynamical chiral-symmetry breakdown (e.g., dynamical quark masses, Goldstone pions) can be ignored in the quark gas so quarks are characterized by their current-algebra masses. Finally, we assume that the properties of the quark Fermi gas can be computed using renormalization-group-improved QCD perturbation theory. Unfortunately, at the momentum scale typical of the problem at hand (roughly $M_N/3$) α_c is not small. Other methods (e.g., lattice Monte Carlo simulations of QCD) may eventually yield information about quark matter; at present, perturbative QCD is the only tool available. Our study of strange matter in bulk is

Three important statements:

1. Limiting case of original (MIT) bag model
(bag is filled with relativistic Fermi gas)
-> thermodynamic bag model
 2. Chiral symmetry is restored
bare quark masses
 3. Perturbation theory applicable
(more or less)
2. and 3. are related.


Absolutely Stable Strange Matter?

EDWARD FARHI AND R. L. JAFFE




Absolutely Stable Strange Matter?

Table 2
Single Flavor, Effective Two-flavor Chiral Bag Constants, and $\mu_{\chi/dc}$ for the Parameterizations of Table 1




Absolutely Stable Strange Matter?

Table 2
Single Flavor, Effective Two-flavor Chiral Bag Constants, and $\mu_{\chi/dc}$ for the Parameterizations of Table 1



Absolutely Stable Strange Matter?



prediction of absolutely stable strange quark matter crucially relies on neglecting dynamical chiral symmetry breaking for light quarks

Difficult to confirm even if one assumes no DCSB for strange quarks at all

Conclusions Part II



vBAG: ■

- vector interaction resolves the problem of too soft bag model EoS w/o perturbative corrections
- No problem at all to obtain stable hybrid neutron star configurations
- Standard BAG models bag constant is understood to mimic confinement, D χ SB is absent
- vBAG introduces effective bag constant with similar values to original BAG


$$\bar{B_{eff}} = \sum_f \bar{B_\chi^f} - \bar{B_{dc}}$$

- However, positive value due to chiral symmetry breaking, (de)confinement reduces B
- Absolutely stable strange matter hypothesis is not trivial to hold up accounting for D χ SB
- NJL and partially Bag model result from particular approximation within Dyson-Schwinger approach
rainbow approximation (quark-gluon vertex) + contact interaction (gluon propagator)
- Consequence: both models lack momentum dependent gap solutions

Finite Temperature



Medium Corrections



Coherent picture:


(de)confinement bag constant reduces with temperature

- > nuclear and chiral quark matter become similar
- > indicates cross-over behaviour

Careful:

Model is not able to actually describe crossover
1st order phase transition is ‘hardwired’ :
NM and QM EoS are modeled independently
NM EoS doesn’t know about quarks

Phase Diagram



Location of transition line


$v\text{Bag}$:

defined by chiral transition
does not depend on hadronic EoS
'low' μ

NJL(+Maxwell) :

changes with NM EoS
'high' μ

Phase Diagram



Location of transition line

Onset of coexistence domain:
Depends on NM EoS for both

Onset of pure quark phase:
vBag:
defined by chiral transition
does not depend on hadronic EoS


NJL(+Maxwell):
changes with NM EoS
'high' μ

Medium Corrections

$$\begin{aligned}
& \frac{\partial B_{dc}}{\partial \mu_C} =: n_{C,dc}(T, \mu_C) \\
&= \frac{\partial}{\partial \mu_C} \{ P^H(T, \mu_C, \mu_B = \mu_{B,\chi}(T, \mu_C)) \} \\
&= \frac{\partial P^H}{\partial \mu_C}(T, \mu_C, \mu_{B,\chi}) + \frac{\partial \mu_{B,\chi}}{\partial \mu_C} \frac{\partial P^H}{\partial \mu_B}(T, \mu_C, \mu_{B,\chi}) \\
&= n_C^H(T, \mu_C, \mu_{B,\chi}) - \tilde{n}_C^Q(T, \mu_C, \mu_{B,\chi}) \frac{n_B^H(T, \mu_C, \mu_{B,\chi})}{n_B^Q(T, \mu_C, \mu_{B,\chi})} \\
& \frac{\partial B_{dc}}{\partial T} =: s_{dc}(T, \mu_C) \\
&= \frac{\partial}{\partial T} \{ P^H(T, \mu_C, \mu_B = \mu_{B,\chi}(T, \mu_C)) \} \\
&= \frac{\partial P^H}{\partial T}(T, \mu_C, \mu_{B,\chi}) + \frac{\partial \mu_{B,\chi}}{\partial T} \frac{\partial P^H}{\partial \mu_B}(T, \mu_C, \mu_{B,\chi}) \\
&= s^H(T, \mu_C, \mu_{B,\chi}) - \tilde{s}^Q(T, \mu_C, \mu_{B,\chi}) \frac{n_B^H(T, \mu_C, \mu_{B,\chi})}{n_B^Q(T, \mu_C, \mu_{B,\chi})}
\end{aligned}$$

$$s^Q(T, \mu_C, \mu_B) = \tilde{s}^Q(T, \mu_C, \mu_B) + \frac{\partial B_{\text{dc}}}{\partial T}$$


TK, T.Fischer, M.Hempel [arXiv:1603.03679](https://arxiv.org/abs/1603.03679), ApJ (subm)




Medium Corrections

$$R_\varepsilon = \varepsilon_{\text{dc}} / \varepsilon^Q$$

$$\begin{aligned} \varepsilon^Q(T, \mu_C, \mu_B) = & \tilde{\varepsilon}^Q(T, \mu_C, \mu_B) - B_{\text{dc}}(T, \mu_C) \\ & + T s_{\text{dc}}(T, \mu_C) + \mu_C n_{C,\text{dc}}(T, \mu_C) \end{aligned}$$




Proto Neutron Star Configurations



Conclusions

QCD in medium (near critical line):

- Task is difficult
- Not addressable by LQCD
- Not addressable by pQCD
- DSE are promising tool to tackle non-perturbative in-medium QCD
- Qualitatively very different results depending on effective gluon coupling
- Bag model mostly a simple limiting case of NJL model
- NJL model a simple contact interaction model in the gluon sector
- vBag connects them, other models exist



CompStar online supernovae EoS

compose.obspm.fr

- △ free-access website (compose.obspm.fr)
- △ hosted at LUTH, l'Observatoire de Paris
- △ **database of EoS tables**
- △ thermodynamic properties, chemical composition, microscopic quantities
- △ tabulation in temperature, baryon density and hadronic charge fraction
- △ flexible data format
- △ cold neutron star EoS direct input in Nrotstar code of LORENE library
- △ **software**
- △ extraction and interpolation of data
- △ calculation of additional quantities
- △ **manual [arXiv:1307.5715](https://arxiv.org/abs/1307.5715) (75 pp.)**
- △ **links to related projects**

CompOSE

Home Manual EoS

EoS Overview EoS Details 1 parameter EoS 2 parameter EOS 3 Parameter EOS

Software (Interpolation, etc) About CompOSE Newsletter Contact

CompStar Online
Supernovæ Equations of State



Cold neutron star EoS

► Relativistic mean field models

EoS	Particles	Tmin	Tmax	pts	nmin	nmax	pts	Ymin	Ymax	pts	details
DDHdelta	npe	(0)	(0)	(1)	7.92e-15	1.00e+00	426	xx	xx	(1)	details
GM1	npe	(0)	(0)	(1)	7.92e-15	1.00e+00	247	xx	xx	(1)	details

► Relativistic mean field models with hyperons

EoS	Particles	Tmin	Tmax	pts	nmin	nmax	pts	Ymin	Ymax	pts	details
GM1 Y4	npeBs	(0)	(0)	(1)	7.92e-15	1.19e+00	301	xx	xx	(1)	details
GM1 Y5	npeBs	(0)	(0)	(1)	7.92e-15	1.31e+00	345	xx	xx	(1)	details
GM1 Y6	npeBs	(0)	(0)	(1)	7.92e-15	1.12e+00	336	xx	xx	(1)	details
DDHdelta	npeBs	(0)	(0)	(1)	7.92e-15	0.99e+00	305	xx	xx	(1)	details
Y4											

► Microscopic calculations

EoS	Particles	Tmin	Tmax	pts	nmin	nmax	pts	Ymin	Ymax	pts	details
APR	npem	(0)	(0)	(1)	7.92e-15	1.34e+00	172	xx	xx	(1)	details
BHF-BBB2	npe	(0)	(0)	(1)	7.92e-15	1.50e+00	84	xx	xx	(1)	details

core team : S. Typel, M. Oertel, T.K.

- S: DCSB
- V: renormalizes μ
- D: diquarks \rightarrow 2SC, CFL
- TD Potential minimized
in mean-field approximation
- Effective model by its nature;
can be motivated (1g-exchange)
doesn't have to though and can
be extended (KMT, PNJL)
- possible to describe hadrons


Effective Lagrangian

$$\begin{aligned}\mathcal{L}_{int} = & G_S \eta_D \sum_{a,b=2,5,7} (\bar{q} i \gamma_5 \tau_a \lambda_b C \bar{q}^T) (q^T C i \gamma_5 \tau_a \lambda_a q) \\ & + G_S \sum_{a=0}^8 [(\bar{q} \tau_a q)^2 + \eta_V (\bar{q} i \gamma_0 q)^2]\end{aligned}$$


Thermodynamical potential

$$\begin{aligned}\Omega(T, \mu) = & \frac{\phi_u^2 + \phi_d^2 + \phi_s^2}{8G_S} - \frac{\omega_u^2 + \omega_d^2 + \omega_s^2}{8G_V} + \frac{\Delta_{ud}^2 + \Delta_{us}^2 + \Delta_{ds}^2}{4G_D} \\ & - \int \frac{d^3 p}{(2\pi)^3} \sum_{n=1}^{18} \left[E_n + 2T \ln \left(1 + e^{-E_n/T} \right) \right] + \Omega_l - \Omega_0.\end{aligned}$$

DCSB and Confinement



Qin et al., PRL (2011)



Fischer et al., Phys.Lett. B (2011)