

A Global Analysis of Resonance-enhanced Light Scalar Dark Matter

Tobias Binder, Sreemanti Chakraborti, Shigeki Matsumoto and Yu Watanabe arXiv:2205.10149


1. Introduction

- Thermal dark matter ($\mathcal{O}(1)\text{MeV} \lesssim m_{\text{DM}} \lesssim \mathcal{O}(100)\text{TeV}$) is an attractive DM candidate.
- Among them, WIMP ($\mathcal{O}(1)\text{GeV} \lesssim m_{\text{DM}} \lesssim \mathcal{O}(1)\text{TeV}$) has intensively searched for, however not been founded.
→ **Light thermal DM ($m_{\text{DM}} \lesssim \mathcal{O}(1)\text{GeV}$) is getting a more attention.**
- σ of them at recombination (freeze-out) is required to be less than (about) 1pb by CMB (relic abundance), so $\langle\sigma v\rangle$ should be **velocity-dependent** e.g. s-channel resonance. Furthermore such candidate may **solve the core-cusp problem**.
- As an example of those models we studied the model including the **scalar singlet DM and mediator**.

3. Favored parameter regions

Self-Scattering


- N-body simulations (astrophysical observations) prefer DM density profile with a cusp (core) near the galactic center (GC). This mismatch may be **solved by self-scattering of DM**, which thermalize DM near the GC.



- Self-scattering cross sections needed to solve core-cusp problem.
- Velocity-dependent self-scattering cross sections seem to fit the data well.

CMB


- Decoupling of thermal BSM particles after that of neutrino is forbidden due to the asymmetrical entropy injection to the primordial plasma.



- $\langle\sigma v\rangle$ at the recombination era ($v \ll 10$ km/s) should be small in order not to modify the anisotropy of CMB.
- S-channel resonance make $\langle\sigma v\rangle$ **enhanced (suppressed) at freeze-out (recombination)**.

Relic abundance

- We assume DM is produced by freeze-out mechanism.
- Since DM annihilations are enhanced through s-channel resonance, $\sin\theta$ becomes small. → **Early Kinetic Decoupling** occurs.
- This **reduces the abundance**, and makes $\sin\theta$ **6 times smaller**.



- The parameter region favored by above 3 conditions at 1σ (red) 2σ (blue) with the yellow crossing mark being the best fit parameter set.

- The correlation among v_R , γ_ϕ and σ_0 , $\sin\theta$ and range of m_ϕ are decided by self-scattering, relic abundance and CMB, respectively.

2. Model

- The Lagrangian is

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \frac{1}{2}(\partial_\mu \chi)^2 - \frac{\mu_\chi^2}{2}\chi^2 - \frac{\lambda_{H\chi}}{2}|H|^2\chi^2 - \frac{\lambda_\chi}{4!}\chi^4 + \frac{1}{2}(\partial_\mu \Phi)^2 - \frac{\mu_{\Phi\chi}}{2}\Phi\chi^2 - \frac{\lambda_{\Phi\chi}}{4}\Phi^2\chi^2 - V(\Phi, H),$$

$$V(\Phi, H) = \mu_{\Phi H}\Phi|H|^2 + \frac{\lambda_{\Phi H}}{2}\Phi^2|H|^2 + \mu_1^3\Phi + \frac{\mu_\Phi^2}{2}\Phi^2 + \frac{\mu_3}{3!}\Phi^3 + \frac{\lambda_\Phi}{4!}\Phi^4,$$

- We assume $m_\phi \simeq 2m_\chi$ (**s-channel resonance**). →
- Phenomenologically important parameters are
 - m_ϕ ... mass of mediator
 - v_R ... place of resonance ($\equiv 2(m_\phi/m_\chi - 2)^{1/2}$)
 - $\sin\theta$... mixing angle between mediator and higgs boson
 - γ_ϕ ... invisible decay rate of mediator ($\Gamma(\phi \rightarrow \chi\chi) = \gamma_\phi m_\phi v_{\text{DM}}$) ↓
 - σ_0 ... velocity-independent part of the self-scattering cross section


4. Constraints from experiments

Collider

- Since in the favored parameter space the **mediator decays invisibly**, **invisible decays of SM particles** e.g. Higgs boson, K and B meson gives the severe constraint.

Direct Detection


- The **scattering cross section between DM and a nucleon** is constrained. **Migdal effect** plays an important role.



- The constraint from collider (gray dots), direct detection at present (gray-shaded region) and that in the near future (green line).
- Since $\sin\theta$ is **suppressed**, the constraint is **weak**.

Indirect Detection

- Voyager can observe e^\pm produced by annihilation of DM.
- COMPTEL gives the most stringent constraint to the γ -ray produced by annihilation of DM.
- There are **large uncertainties** e.g. DM profile, hadronic fragmentation functions of sub-GeV DM.



- The constraint from Voyager (gray dots), COMPTEL (gray-shaded region) and γ -ray observations in the near future (green line).
- Several parameter sets **survives at present** and **almost all of them will be constrained**.

5. Summary

- Light thermal DM with velocity-dependent $\langle\sigma v\rangle$** is an attractive DM candidate.
- As an example of those models we studied the model including **the scalar singlet DM and mediator**.
- A part of **attractive regions** in which DM can solve core-cusp problem, explain the relic density and overcome the constraint from CMB is still **surviving** from constraints at present concerning the uncertainties.
- Almost all of these will be **constrained** by **near future MeV γ -ray** observations e.g. GECCO and COSI.