

Smart and Human

常翔学園

摂南大学



Complex Langevin analysis of the spontaneous breaking of 10D rotational symmetry in the Euclidean IKKT matrix model (arXiv:2002.07410)

Takehiro Azuma (Setsunan Univ.)

Asia-Pacific Symposium for Lattice Field Theory

Aug. 5th 2020, 14:00-14:20

with Konstantinos N. Anagnostopoulos (NTUA), Yuta Ito (KEK), Jun Nishimura (KEK, SOKENDAI), Toshiyuki Okubo (Meijo Univ.) and Stratos Kovalkov Papadoudis(NTUA)

Difficulties in simulating complex partition functions.

$$Z = \int dA \exp(-S_0 + i\Gamma), \quad Z_0 = \int dA e^{-S_0}$$

Sign problem:

The reweighting $\langle \mathcal{O} \rangle = \frac{\langle \mathcal{O} e^{i\Gamma} \rangle_0}{\langle e^{i\Gamma} \rangle_0}$ requires configs. $\exp[O(N^2)]$

$\langle^* \rangle_0 = (\text{V.E.V. for the phase-quenched partition function } Z_0)$

2. Euclidean type IIB matrix model

3



Promising candidate for nonperturbative string theory

[N. Ishibashi, H. Kawai, Y. Kitazawa and A. Tsuchiya, hep-th/9612115] (a.k.a. "IKKT model")

$$Z = \int dA d\psi e^{-(S_b + S_f)}$$
$$S_b = -\frac{N}{4} \text{tr}[A_\mu, A_\nu]^2, \quad S_f = N \text{tr} \bar{\psi}_\alpha (\Gamma^\mu)_{\alpha\beta} [A_\mu, \psi_\beta]$$

-Euclidean case after Wick rotation $A_0 \rightarrow iA_{10}, \Gamma^0 \rightarrow -i\Gamma_{10}$.
 \Rightarrow Path integral is finite without cutoff.

[W. Krauth, H. Nicolai and M. Staudacher, hep-th/9803117, P. Austing and J.F. Wheeler, hep-th/0103059]

• $A_\mu, \Psi_\alpha \Rightarrow N \times N$ Hermitian traceless matrices.

($\mu=1,2,\dots,10, \alpha,\beta=1,2,\dots,16$)

• Eigenvalues of A_μ : spacetime coordinate $\Rightarrow \mathcal{N}=2$ SUSY

2. Euclidean type IIB matrix model

4



$$Z = \int dA d e^{-S_b} \underbrace{\left(\int d\psi e^{-S_f} \right)}_{= \text{Pf } \mathcal{M} = |\text{Pf } \mathcal{M}| e^{i\Gamma} = e^{-S_{f,\text{eff}}}} = \int dA \underbrace{e^{-S}}_{e^{-(S_b + S_{f,\text{eff}})}}$$

- Pf \mathcal{M} 's *complex phase* Γ contributes to the **Spontaneous Symmetry Breaking (SSB)** of SO(10).
- Result of Gaussian Expansion Method (GEM)

[T.Aoyama, J.Nishimura, and T.Okubo, arXiv:1007.0883, J.Nishimura, T.Okubo and F.Sugino, arXiv:1108.1293]

SSB SO(10) \rightarrow SO(3).

Dynamical compactification to 3-dim spacetime.

Complex Langevin Method (CLM)

⇒ Solve the complex version of the Langevin equation.

[Parisi, Phys.Lett. 131B (1983) 393, Klauder, Phys.Rev. A29 (1984) 2036]

$$\frac{d(A_\mu)_{ij}}{dt} = - \underbrace{\frac{\partial S}{\partial (A_\mu)_{ji}}}_{\text{drift term}} + \eta_{\mu,ij}(t) = - \left\{ \frac{\partial S_b}{\partial (A_\mu)_{ji}} - \underbrace{\frac{1}{2} \text{Tr} \left(\frac{\partial \mathcal{M}}{\partial (A_\mu)_{ji}} \mathcal{M}^{-1} \right)}_{=\partial \log \text{Pf.} \mathcal{M} / \partial (A_\mu)_{ji}} \right\} + \eta_{\mu,ij}(t)$$

▪ A_μ : **Hermitian** → **general complex** traceless matrices.

▪ η_μ : Hermitian-matrix white noise obeying the probability distribution $\exp \left(-\frac{1}{4} \int \text{tr} \eta^2(t) dt \right)$

3. Complex Langevin Method

6



CLM does not work when it encounters these problems:

(1) Excursion problem: A_μ is too far from Hermitian
 \Rightarrow **Gauge Cooling** minimizes the **Hermitian norm**

$$\mathcal{N} = \frac{-1}{10N} \sum_{\mu=1}^{10} \text{tr}[(A_\mu - (A_\mu)^\dagger)^2]$$

(2) Singular drift problem:

The drift term $dS/d(A_\mu)_{ji}$ diverges due to \mathcal{M} 's **near-zero** eigenvalues.

We trust CLM when the distribution $p(u)$ of the **drift norm**

$$u = \sqrt{\frac{1}{10N^3} \sum_{\mu=1}^{10} \sum_{i,j=1}^N \left| \frac{\partial S}{\partial (A_\mu)_{ji}} \right|^2} \quad \text{falls exponentially as } p(u) \propto e^{-au}.$$

[K. Nagata, J. Nishimura and S. Shimasaki, arXiv:1606.07627]

Look at the **drift term** \Rightarrow Get the drift of CLM!!

3. Complex Langevin Method

Mass deformation [Y. Ito and J. Nishimura, arXiv:1609.04501]

• SO(D) symmetry breaking term $\Delta S_b = N \frac{\epsilon}{2} \sum_{\mu=1}^{10} m_{\mu} \text{tr}(A_{\mu})^2$

Order parameters for SSB of SO(10): $\lambda_{\mu} = \text{Re} \left\{ \frac{1}{N} \text{tr}(A_{\mu})^2 \right\}$

• Fermionic mass term: $\Delta S_f = N m_f \text{tr} \left(\bar{\psi}_{\alpha} (i \Gamma_8 \Gamma_9^{\dagger} \Gamma_{10})_{\alpha\beta} \psi_{\beta} \right)$



Avoids the singular eigenvalue distribution of \mathcal{M} .

Extrapolation (i) $N \rightarrow \infty \Rightarrow$ (ii) $\epsilon_m \rightarrow 0 \Rightarrow$ (iii) $m_f \rightarrow 0$.

4. Result

$$\Delta S_b = N \frac{\epsilon}{2} \sum_{\mu=1}^{10} m_{\mu} \text{tr}(A_{\mu})^2 \quad \Delta S_f = N m_f \text{tr} \left(\bar{\psi}_{\alpha} (i \Gamma_8 \Gamma_9^{\dagger} \Gamma_{10})_{\alpha\beta} \psi_{\beta} \right) \quad \rho_{\mu}(\epsilon, m_f) = \frac{\langle \lambda_{\mu} \rangle_{\epsilon, m_f}}{\sum_{\nu=1}^{10} \langle \lambda_{\nu} \rangle_{\epsilon, m_f}}$$

$$m_{\mu} = (0.5, 0.5, 1, 2, 4, 8, 8, 8, 8, 8)$$







4. Result

$$\Delta S_b = N \frac{\epsilon}{2} \sum_{\mu=1}^{10} m_{\mu} \text{tr}(A_{\mu})^2 \quad \Delta S_f = N m_f \text{tr} \left(\bar{\psi}_{\alpha} (i \Gamma_8 \Gamma_9 \Gamma_{10})_{\alpha\beta} \psi_{\beta} \right) \quad \rho_{\mu}(\epsilon, m_f) = \frac{\langle \lambda_{\mu} \rangle_{\epsilon, m_f}}{\sum_{\nu=1}^{10} \langle \lambda_{\nu} \rangle_{\epsilon, m_f}}$$

(ii) $\epsilon \rightarrow 0$ after $N \rightarrow \infty$

$$m_{\mu} = \begin{cases} (0.5, 0.5, 0.5, 1, 2, 4, 8, 8, 8, 8) & (m_f = 3.0) \\ (0.5, 0.5, 1, 2, 4, 8, 8, 8, 8, 8) & (m_f < 3.0) \end{cases}$$







4. Result

$$\Delta S_f = Nm_f \text{tr} \left(\bar{\psi}_\alpha (i\Gamma_8 \Gamma_9^\dagger \Gamma_{10})_{\alpha\beta} \psi_\beta \right) \quad (\text{no } \Delta S_b = N \frac{\epsilon}{2} \sum_{\mu=1}^{10} m_\mu \text{tr}(A_\mu)^2 \text{ term})$$

GEM result for $m_f > 0$ at 3 loop solutions of SO(6), SO(7) ansatz

$$\rho_\mu(\epsilon, m_f) = \frac{\langle \lambda_\mu \rangle_{\epsilon, m_f}}{\sum_{\nu=1}^{10} \langle \lambda_\nu \rangle_{\epsilon, m_f}}$$



Dynamical compactification of the spacetime in the Euclidean IKKT model.

"Complex Langevin Method"

⇒ trend of SSB $SO(10) \rightarrow SO(3)$.

Future works

Application of CLM to the Lorentzian version of the type IIB matrix model

[K.N. Anagnostopoulos, T. Azuma, [K. Hatakeyama, M. Hirasawa](#), Y. Ito, J. Nishimura, S.K. Papadoudis and A. Tsuchiya, work in progress] [their talks](#)