

Perturbative dynamics of fuzzy spheres at large N (hep-th/0410263)

Takehiro Azuma

High-energy Accelerator Research Organization (KEK)

JPS meeting 2005 at Tokyo University of Science

Mar. 27th 2005, 11:30 ~ 11:45

Contents

1	Introduction	2
2	The model and its classical solution	3
3	All order calculation from one-loop effective action	6
4	Conclusion	10

1 Introduction

Matrix models on a homogeneous space

Motivations of fuzzy manifold studies:

- Relation between the non-commutative field theory and the superstring.
- Novel regularization scheme alternative to lattice regularization.
- Prototype of the curved-space background in the large- N reduced models.

Matrix models on a homogeneous space G/H :

G = (a Lie group), H = (a closed subgroup of G).

$$S^2 = \text{SU}(2)/\text{U}(1), S^2 \times S^2, S^4 = \text{SO}(5)/\text{U}(2), \text{CP}^2 = \text{SU}(3)/\text{U}(2), \dots$$

These fuzzy manifolds are compact, and thus realized by finite matrices.

The Chern-Simons term is added to accommodate the classical solution of the fuzzy manifolds.

2 The model and its classical solution

3d Yang-Mills-Chern-Simons (YMCS) model

⇒ a toy model with fuzzy sphere solutions:

S. Iso, Y. Kimura, K. Tanaka and K. Wakatsuki, hep-th/0101102.

$$S[A] = N \text{tr} \left(-\frac{1}{4} [A_\mu, A_\nu]^2 + \frac{2i\alpha}{3} \epsilon_{\mu\nu\rho} A_\mu A_\nu A_\rho \right).$$

- Defined in the 3-dimensional Euclidean space ($\mu, \nu, \rho = 1, 2, 3$).
- Convergence of the path integral P. Austing and J. F. Wheeler, hep-th/0310170.
- Classical equation of motion: $[A_\nu, [A_\mu, A_\nu]] - i\alpha \epsilon_{\mu\nu\rho} [A_\nu, A_\rho] = 0$.
- fuzzy S^2 classical solutions: $A_\mu = X_\mu = \alpha L_\mu$, (where $[L_\mu, L_\nu] = i\epsilon_{\mu\nu\rho} L_\rho$).
 $L_\mu = (N \times N \text{ representation of the } \text{SU}(2) \text{ Lie algebra})$.
 Casimir operator: $Q = A_1^2 + A_2^2 + A_3^2 = R^2 1_N$.
 $R = (\text{radius of the fuzzy sphere}) = \frac{\alpha}{2} \sqrt{N^2 - 1}$.

First-order phase transition

Monte Carlo simulation launched from fuzzy sphere classical solution:

Critical point at $\alpha_{\text{cr}} \simeq \frac{2.1}{\sqrt{N}}$.

- $\alpha < \alpha_{\text{cr}}$: Yang-Mills phase

Strong quantum effects.

behavior like the $\alpha = 0$ case.


T. Hotta, J. Nishimura and A. Tsuchiya, hep-th/9811220,

$$\left\langle \frac{S}{N^2} \right\rangle \simeq O(1), \quad \left\langle \frac{1}{N} \text{tr} A_\mu^2 \right\rangle \simeq O(1).$$


- $\alpha > \alpha_{\text{cr}}$: fuzzy sphere phase.

Fuzzy sphere configuration is stable.

Yang-Mills phase



Fuzzy sphere phase



Phase transition from the one-loop effective action

The effective action Γ is saturated at the **one-loop** level at large N .

T. Imai, Y. Kitazawa, Y. Takayama and D. Tomino, hep-th/0307007.

Effective action at one-loop around


$A_\mu = tX_\mu$ (where $\tilde{\alpha} = \alpha\sqrt{N}$).

$$\frac{\Gamma_{1\text{-loop}}}{N^2} \simeq \tilde{\alpha}^4 \left(\frac{t^4}{8} - \frac{t^3}{6} \right) + \log t.$$

The local minimum disappears at

$$\tilde{\alpha} < \tilde{\alpha}_{\text{cr}} = \left(\frac{8}{3}\right)^{\frac{3}{4}} \simeq 2.086 \dots$$

Consistent with the Monte Carlo simulation.



3 All order calculation from one-loop effective action

The free energy W can be obtained by the extremum of the effective action.

Expansion around $A_\mu = \beta L_\mu$: ($\tilde{\beta} = \beta\sqrt{N}$)

$$\lim_{N \rightarrow \infty} \frac{1}{N^2} \Gamma(\tilde{\beta}) = \left(\frac{\tilde{\beta}^4}{8} - \frac{1}{6} \tilde{\alpha} \tilde{\beta}^3 \right) + \log \tilde{\beta}.$$

Local minimum for $\tilde{\alpha} > \tilde{\alpha}_{\text{cr}} = \sqrt[3]{\frac{512}{27}}$:

$$\begin{aligned} \tilde{\beta} &= f(\tilde{\alpha}) = \frac{\tilde{\alpha}}{4} \left(1 + \sqrt{1 + \delta} + \sqrt{2 - \delta + \frac{2}{\sqrt{1 + \delta}}} \right) \\ &= \tilde{\alpha} \left(1 - \frac{2}{\tilde{\alpha}^4} - \frac{12}{\tilde{\alpha}^8} - \frac{120}{\tilde{\alpha}^{12}} - \frac{1456}{\tilde{\alpha}^{16}} - \dots \right), \text{ where} \\ \delta &= 4\tilde{\alpha}^{-\frac{4}{3}} \left[\left(1 + \sqrt{1 - \frac{512}{27\tilde{\alpha}^4}} \right)^{\frac{1}{3}} + \left(1 - \sqrt{1 - \frac{512}{27\tilde{\alpha}^4}} \right)^{\frac{1}{3}} \right]. \end{aligned}$$

Free energy and observables:

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N^2} W &= \left(\frac{1}{8} f(\tilde{\alpha})^4 - \frac{1}{6} \tilde{\alpha} f(\tilde{\alpha})^3 \right) + \log f(\tilde{\alpha}) \\ &= -\frac{\tilde{\alpha}^4}{24} + \log \tilde{\alpha} - \frac{1}{\tilde{\alpha}^4} - \frac{14}{3\tilde{\alpha}^8} - \frac{110}{3\tilde{\alpha}^{12}} - \frac{364}{\tilde{\alpha}^{16}} - \dots, \\ \lim_{N \rightarrow \infty} \frac{1}{N^2} \langle S \rangle &= \frac{3}{4} - \frac{1}{24} \tilde{\alpha} f(\tilde{\alpha})^3 \\ &= -\frac{\tilde{\alpha}^4}{24} + 1 + \underbrace{\frac{1}{\tilde{\alpha}^4}} + \frac{28}{3\tilde{\alpha}^8} + \frac{110}{\tilde{\alpha}^{12}} + \frac{1456}{\tilde{\alpha}^{16}} + \dots \end{aligned}$$

agrees with two-loop calculation!

All order calculation of generic observables \mathcal{O}

Consider the action $S_\epsilon = S + \epsilon\mathcal{O}$.

Corresponding free energy:

$$\begin{aligned} W_\epsilon &= -\log \left(\int d\tilde{A} e^{-(S+\epsilon\mathcal{O})} \right) = -\log \left(\int d\tilde{A} e^{-S} \right) + \epsilon \frac{\int d\tilde{A} \mathcal{O} e^{-S}}{\int d\tilde{A} e^{-S}} + \mathcal{O}(\epsilon^2) \\ &= W + \epsilon \langle \mathcal{O} \rangle + \mathcal{O}(\epsilon^2). \end{aligned}$$

One-loop effective action (take only 1PI diagrams into account)

$$\Gamma_\epsilon(\tilde{\beta}) = \Gamma(\tilde{\beta}) + \epsilon\Gamma_1(\tilde{\beta}) + \mathcal{O}(\epsilon^2).$$

Its saddle point:

$$\frac{\partial}{\partial \tilde{\beta}} \Gamma_\epsilon(\tilde{\beta}) = 0, \quad \Rightarrow \tilde{\beta} = f(\tilde{\alpha}) + \epsilon g(\tilde{\alpha}) + \mathcal{O}(\epsilon^2).$$

Plugging this solution, we obtain the free energy as

$$W_\epsilon = \Gamma_\epsilon(f(\tilde{\alpha}) + \epsilon g(\tilde{\alpha}) + \dots) = \Gamma(f(\tilde{\alpha})) + \epsilon \left(\Gamma_1(f(\tilde{\alpha})) + g(\tilde{\alpha}) \underbrace{\left(\frac{\partial \Gamma}{\partial \tilde{\beta}} \right) \Big|_{\tilde{\beta}=f(\tilde{\alpha})}}_{=0} \right) + \mathcal{O}(\epsilon^2).$$

We thus obtain $\langle \mathcal{O} \rangle = \Gamma_1(f(\tilde{\alpha}))$.

All order calculation of the spacetime content:





$$\lim_{N \rightarrow \infty} \frac{1}{N} \left\langle \frac{1}{N} \text{tr} A_{\mu}^2 \right\rangle = \frac{\tilde{\alpha}^2}{4} \underbrace{- \frac{1}{\tilde{\alpha}^2}}_{\text{one-loop}} \cdot$$

The one-loop effect comes from **tadpole diagrams**.

$$\frac{1}{N} \left\langle \frac{1}{N} \text{tr} A^2 \right\rangle = \frac{1}{4} f(\tilde{\alpha})^2 = \frac{1}{4} \tilde{\alpha}^2 - \frac{1}{\tilde{\alpha}^2} - \frac{5}{\tilde{\alpha}^6} - \frac{48}{\tilde{\alpha}^{10}} - \frac{572}{\tilde{\alpha}^{14}} - \dots$$

Other observables:

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{\sqrt{N}} \langle M \rangle &= -\frac{1}{6} f(\tilde{\alpha})^3 = -\frac{1}{6} \tilde{\alpha}^3 + \frac{1}{\tilde{\alpha}} + \frac{4}{\tilde{\alpha}^5} + \frac{112}{3\tilde{\alpha}^9} + \frac{440}{\tilde{\alpha}^{13}} + \dots, \\ \lim_{N \rightarrow \infty} \left\langle \frac{1}{N} \text{tr} (F_{\mu\nu})^2 \right\rangle &= 3 + \frac{1}{2} \tilde{\alpha} f(\tilde{\alpha})^3 = \frac{1}{2} \tilde{\alpha}^4 - \frac{12}{\tilde{\alpha}^4} - \frac{112}{\tilde{\alpha}^8} - \frac{1320}{\tilde{\alpha}^{12}} - \dots \end{aligned}$$



4 Conclusion

- In this talk, we have scrutinized the perturbative dynamics of the **3d YMCS model**.
- We have obtained the **all order results** for generic observables at large N .

Future direction

- Extension of this technique to the **4-dimensional fuzzy manifolds**:
fuzzy CP^2 (hep-th/0405277), fuzzy $S^2 \times S^2$ (hep-th/0503*).**