

Compositeness of baryon resonances in chiral dynamics



Tetsuo Hyodo^a,

Daisuke Jido^b, and Atsushi Hosaka^c


Tokyo Institute of Technology^a YITP, Kyoto^b RCNP, Osaka^c

supported by Global Center of Excellence Program
“Nanoscience and Quantum Physics”

2011, Jan. 27th 1


Structure of hadron resonances

Example) baryon excited state



Excited states
= resonances in hadron scattering

Exotic structure **near threshold?**
c.f. ^{12}C Hoyle state



Study of the internal structure

How to investigate the internal structure?




- Comparison of **model calculation** with **experiments**
(mass, width, decay properties, etc.)

- : Any model can describe data with appropriate corrections
- : Model-dependent result


- **Extrapolation** to the ideal world, change the environment
(large N_c , symmetry restoration, etc.)

- : Structure may change during the extrapolation
- : Qualitative discussion only

--> **model-independent** and **quantitative** study?

-  Introduction
-  Chiral SU(3) dynamics
-  Origin of resonances in chiral dynamics
 - Natural renormalization condition

[T. Hyodo, D. Jido, A. Hosaka, Phys. Rev. C78, 025203 \(2008\)](#)


-  Compositeness of bound states
 - Field renormalization constant Z

[T. Hyodo, D. Jido, A. Hosaka, arXiv:1009.5754 \[nucl-th\]; in preparation](#)

-  Summary

s-wave low energy interaction

Low energy NG boson (Ad) + target hadron (T) scattering

$$\alpha \left[\begin{array}{c} \text{Ad}(q) \\ T(p) \end{array} \right] = \frac{1}{f^2} \frac{p \cdot q}{2M_T} \langle \mathbf{F}_T \cdot \mathbf{F}_{\text{Ad}} \rangle_\alpha + \mathcal{O}\left(\left(\frac{m}{M_T}\right)^2\right)$$


Projection onto s-wave: Weinberg-Tomozawa (WT) term

Y. Tomozawa, *Nuovo Cim.* 46A, 707 (1966); S. Weinberg, *Phys. Rev. Lett.* 17, 616 (1966)

$$V_{ij} = -\frac{C_{ij}}{4f^2} (\omega_i + \omega_j) \quad \text{energy dependence (derivative coupling)}$$

decay constant of π ($g_V=1$)

$$C_{ij} = \sum_{\alpha} C_{\alpha,T} \left(\begin{array}{cc|c} 8 & T & \alpha \\ I_{M_i}, Y_{M_i} & I_{T_i}, Y_{T_i} & I, Y \end{array} \right) \left(\begin{array}{cc|c} 8 & T & \alpha \\ I_{M_j}, Y_{M_j} & I_{T_j}, Y_{T_j} & I, Y \end{array} \right)$$

$$C_{\alpha,T} = \langle 2\mathbf{F}_T \cdot \mathbf{F}_{\text{Ad}} \rangle_\alpha = C_2(T) - C_2(\alpha) + 3$$

Group theoretical structure and flavor **SU(3) symmetry** determines **the sign and the strength** of the interaction

Low energy theorem: leading order term in ChPT

Scattering amplitude and unitarity

Unitarity of S-matrix: Optical theorem

$$\text{Im}[T^{-1}(s)] = \frac{\rho(s)}{2} \quad \text{phase space of two-body state}$$

General amplitude by dispersion relation

$$T^{-1}(\sqrt{s}) = \sum_i \frac{R_i}{\sqrt{s} - W_i} + \tilde{a}(s_0) + \frac{s - s_0}{2\pi} \int_{s_+}^{\infty} ds' \frac{\rho(s')}{(s' - s)(s' - s_0)}$$

R_i, W_i, a : to be determined by chiral interaction

Identify dispersion integral = loop function G , the rest = V^{-1}

$$T(\sqrt{s}) = \frac{1}{V^{-1}(\sqrt{s}) - G(\sqrt{s}; a)}$$

Scattering amplitude

V ? chiral expansion of T , (conceptual) matching with ChPT

$$T^{(1)} = V^{(1)}, \quad T^{(2)} = V^{(2)}, \quad T^{(3)} = V^{(3)} - V^{(1)}GV^{(1)}, \quad \dots$$

Amplitude T : consistent with chiral symmetry + unitarity

Chiral unitary approach

Meson-baryon scattering amplitude

- Interaction \leftarrow chiral symmetry

Y. Tomozawa, *Nuovo Cim.* 46A, 707 (1966); S. Weinberg, *Phys. Rev. Lett.* 17, 616 (1966)

- Amplitude \leftarrow unitarity in **coupled channels**

R.H. Dalitz, T.C. Wong, G. Rajasekaran, *Phys. Rev.* 153, 1617 (1967)

$$T = \frac{1}{1 - VG} V$$

chiral
cutoff

N. Kaiser, P. B. Siegel, W. Weise, *Nucl. Phys.* A594, 325 (1995),

E. Oset, A. Ramos, *Nucl. Phys.* A635, 99 (1998),

J. A. Oller, U. G. Meissner, *Phys. Lett.* B500, 263 (2001),

M.F.M. Lutz, E. E. Kolomeitsev, *Nucl. Phys.* A700, 193 (2002), ... many others

It works successfully, also in S=0 sector, meson-meson scattering sectors, systems including heavy quarks, ...

Hadron excited states

Resonances are “dynamically generated”

light baryon	$J^P = 1/2^-$	Λ(1405)	Λ(1670)	Σ(1670)	
		N(1535)	Ξ(1620)	Ξ(1690)	
	$J^P = 3/2^-$	Λ(1520)	Ξ(1820)	Σ(1670)	
heavy		Λ _c (2880)	Λ _c (2593)		D _s (2317)
light meson	$J^P = 1^+$	b ₁ (1235)	h ₁ (1170)	h ₁ (1380)	a ₁ (1260)
		f ₁ (1285)	K ₁ (1270)	K ₁ (1440)	
	$J^P = 0^+$	σ(600)	κ(900)	f ₀ (980)	a ₀ (980)

No states with **exotic** quantum number

- No attraction in exotic channel

T. Hyodo, D. Jido, A. Hosaka, Phys. Rev. Lett. 97, 192002 (2006);

Phys. Rev. D75, 034002 (2007)


--> **Structure** of these resonances?

Classification of resonances

Resonances in two-body scattering

- Knowledge of interaction (potential)
- Experimental data (cross section, ...)


Dynamical state: composite particle, two-body molecule, ...



e.g.) Deuteron in NN, positronium in e^+e^- , ...

CDD pole: elementary particle, preformed state, ...


L. Castillejo, R.H. Dalitz, F.J. Dyson, *Phys. Rev.* 101, 453 (1956)



e.g.) J/ψ in e^+e^- , ...

(Known) CDD poles in chiral unitary approach


Explicit resonance field in V (interaction): $\Delta(1232)$, $\Sigma(1385)$,...



U.G. Meissner, J.A. Oller, Nucl. Phys. A673, 311 (2000)


D. Jido, E. Oset, A. Ramos, Phys. Rev. C66, 055203 (2002)

Contracted resonance propagator in higher order V



G. Ecker, J. Gasser, A. Pich, E. de Rafael, Nucl. Phys. B321, 311 (1989)

V. Bernard, N. Kaiser, U.G. Meissner, Nucl. Phys. A615, 483 (1997)



J.A. Oller, E. Oset, J.R. Pelaez, Phys. Rev. D59, 074001 (1999)

Is that all? **subtraction constant?**

CDD pole in subtraction constant?

Phenomenological (standard) scheme

--> V is given, “ a ” is determined by data

$$T = \frac{1}{(V^{(1)})^{-1} - \underline{G(a)}} \quad \text{leading order}$$

$$T = \frac{1}{(V^{(1)} + V^{(2)})^{-1} - \underline{G(a')}} \quad \text{next to leading order}$$



“ a ” represents the effect which is not included in V .
CDD pole contribution in G ?

Natural renormalization scheme

--> fix “ a ” first, then determine V

to exclude CDD pole contribution from G ,
based on theoretical argument.

Natural renormalization condition

Conditions for the subtraction constant

1) Loop function G should be **negative** below threshold.

\leftrightarrow no states below threshold

$$G(\sqrt{s}) \sim \sum_n \frac{|\langle \dots \rangle|^2}{\sqrt{s} - E_n} \leq 0 \quad \text{for} \quad \sqrt{s} \leq E_0 \quad \rightarrow \text{upper limit for "a"}$$

2) T matches with the chiral interaction V at **low energy**.

$$T(\mu_m; a) = V(\mu_m) \quad \text{for} \quad M_T \leq \mu_m \leq M_T + m \quad \rightarrow \text{lower limit for "a"}$$

To satisfy 1) and 2), "a" is uniquely determined as

$$G(\sqrt{s} = M_T) = 0 \quad \Leftrightarrow \quad T(M_T) = V(M_T)$$

- subtraction constant: a_{natural}

We regard this condition as the **exclusion of the CDD pole contribution from G** .

Pole in the effective interaction: single channel

Leading order V: Weinberg-Tomozawa term

$$V_{\text{WT}} = -\frac{C}{2f^2}(\sqrt{s} - M_T) \quad G(\sqrt{s}; a) = \frac{2M_T}{(4\pi)^2} \left\{ a + \dots \right.$$

$$T^{-1} = V_{\text{WT}}^{-1} - G(a_{\text{pheno}}) = V_{\text{natural}}^{-1} - G(a_{\text{natural}})$$

↑ ChPT

↑ data fit

↑ given

Effective interaction in natural scheme

$$V_{\text{natural}} = -\frac{C}{2f^2}(\sqrt{s} - M_T) + \boxed{\frac{C}{2f^2} \frac{(\sqrt{s} - M_T)^2}{\sqrt{s} - M_{\text{eff}}}} \quad \text{pole!}$$

a seed of resonance?

$$M_{\text{eff}} = M_T - \frac{16\pi^2 f^2}{CM_T \Delta a}, \quad \Delta a = a_{\text{pheno}} - a_{\text{natural}}$$

There is always a pole for $a_{\text{pheno}} \neq a_{\text{natural}}$

- small deviation \Leftrightarrow pole at irrelevant energy scale
- **large deviation \Leftrightarrow pole at relevant energy scale**

Pole in the effective interaction



Pole in the effective interaction (M_{eff}) : pure **CDD pole**

$$T^{-1} = V_{\text{WT}}^{-1} - G(a_{\text{pheno}}) = \boxed{V_{\text{natural}}^{-1}} - G(a_{\text{natural}})$$

For $\Lambda(1405)$: $z_{\text{eff}}^{\Lambda^*} \sim 7.9 \text{ GeV}$ **irrelevant!**

For $N(1535)$: $z_{\text{eff}}^{N^*} = 1693 \pm 37i \text{ MeV}$ **relevant?**

Difference of interactions $\Delta V \equiv V_{\text{natural}} - V_{\text{WT}}$




==> Important CDD pole contribution in N(1535)

Next question: **quantitative** measure for compositeness?

Structure of the deuteron

Deuteron: elementary or NN bound state?

S. Weinberg, Phys. Rev. 137 B672-B678 (1965)



model independent result for a weakly bound state:

$$\boxed{a_s} = \left[\frac{2(1-Z)}{2-Z} \right] \boxed{R} + \mathcal{O}(m_\pi^{-1}), \quad \boxed{r_e} = \left[\frac{-Z}{1-Z} \right] \boxed{R} + \mathcal{O}(m_\pi^{-1})$$

a_s : scattering length

r_e : effective range

←-- Experiments

R : deuteron radius (binding energy)

$$a_s = +5.41 \text{ [fm]}, \quad r_e = +1.75 \text{ [fm]}, \quad R \equiv (2\mu B)^{-1/2} = 4.31 \text{ [fm]}$$

$\Rightarrow Z \lesssim 0.2$ --> **deuteron is almost composite!**

Definition of the compositeness 1-Z

Hamiltonian of two-body system: **free** + interaction V

$$\mathcal{H} = \mathcal{H}_0 + V$$

Complete set for **free** Hamiltonian: bare $|B_0\rangle$ + continuum

$$1 = |B_0\rangle\langle B_0| + \int d\mathbf{k} |\mathbf{k}\rangle\langle \mathbf{k}|$$

$$\mathcal{H}_0|B_0\rangle = E_0|B_0\rangle, \quad \mathcal{H}_0|\mathbf{k}\rangle = E(\mathbf{k})|\mathbf{k}\rangle$$

Physical bound state $|B\rangle$: eigenstate of **full** Hamiltonian

$$(\mathcal{H}_0 + V)|B\rangle = -B|B\rangle$$


B: binding energy

Define **Z** as the **overlap of B and B_0**
: probability of finding the physical bound state in the bare state $|B_0\rangle$

$$Z \equiv |\langle B_0 | B \rangle|^2$$

1 - Z : **Compositeness** of the bound state

They are assumed to be elementary



Model-independent but approximated method

With the Schrödinger equation, we obtain

$$1 - Z = \int d\mathbf{k} \frac{|\langle \mathbf{k} | V | B \rangle|^2}{[E(\mathbf{k}) + B]^2} \quad \langle \mathbf{k} | V | B \rangle : \quad B \rightleftharpoons \begin{array}{c} \nearrow V \\ \bullet \\ \searrow \end{array} \left. \vphantom{\begin{array}{c} \nearrow \\ \bullet \\ \searrow \end{array}} \right\} \mathbf{k}$$

$$= 4\pi \sqrt{2\mu^3} \int_0^\infty dE \frac{\sqrt{E} |G_W(E)|^2}{(E + B)^2} \quad \langle \mathbf{k} | V | B \rangle \equiv G_W[E(\mathbf{k})] \quad \text{for s-wave}$$

Approximation: For small binding energy $B \ll 1$, the vertex $G_W(E)$ can be regarded as a constant: $G_W(E) \sim g_W$

Then the integration can be done analytically, leading to

$$1 - Z = 2\pi^2 \sqrt{2\mu^3} \frac{g_W^2}{\sqrt{B}}$$

Compositeness \leftarrow coupling g and binding energy B

S. Weinberg, Phys. Rev. 137 B672-B678 (1965)

- **Model-independent:** no information of V
- **Approximated:** valid only for small B

Exact but model-dependent method

Formal solution of the Lippmann-Schwinger equation

$$T(E) = V + V \frac{1}{E - \mathcal{H}} V$$

Insert complete set for **full** Hamiltonian (Low's equation)

$$1 = |B\rangle\langle B| + \int dk |\mathbf{k}, \text{in}\rangle\langle \mathbf{k}, \text{in}| \quad V|\mathbf{k}, \text{in}\rangle = T|\mathbf{k}\rangle$$

$$t(E) = v(E) + \frac{|G_W(E)|^2}{E + B} + 4\pi\sqrt{2\mu^3} \int_0^\infty dE' \frac{\sqrt{E'} |t(E')|^2}{E - E' + i\epsilon} \quad (\text{for s-wave})$$

--> integrand of the formula for 1-Z !

Exact expression of the compositeness 1-Z

$$\begin{aligned} 1 - Z &= 4\pi\sqrt{2\mu^3} \int_0^\infty dE \frac{\sqrt{E} |G_W(E)|^2}{(E + B)^2} \\ &= 4\pi\sqrt{2\mu^3} \int_0^\infty dE \frac{\sqrt{E}}{E + B} \left[t(E) - v(E) - 4\pi\sqrt{2\mu^3} \int_0^\infty dE' \frac{\sqrt{E'} |t(E')|^2}{E - E' + i\epsilon} \right] \end{aligned}$$

Short summary

We have defined the compositeness of the bound state 1-Z.

$$1 - Z = 1 - |\langle B_0 | B \rangle|^2 = \int d\mathbf{k} \frac{|\langle \mathbf{k} | V | B \rangle|^2}{[E(\mathbf{k}) + B]^2}$$

Method 1: model independent but approximated

$$1 - Z = 2\pi^2 \sqrt{2\mu^3} \frac{g_W^2}{\sqrt{B}}$$

Method 2: exact (valid for any B) but model dependent

$$1 - Z = 4\pi \sqrt{2\mu^3} \int_0^\infty dE \frac{\sqrt{E}}{E + B} \left[t(E) - v(E) - 4\pi \sqrt{2\mu^3} \int_0^\infty dE' \frac{\sqrt{E'} |t(E')|^2}{E - E' + i\epsilon} \right]$$

- **Model dependent**: interaction V has to be specified (c.f. potential + wave function \rightarrow observable)
- Imaginary part **vanishes** by the optical theorem
- RHS can be calculated by model (chiral unitary approach)
- **Completeness of the full Hamiltonian \leftrightarrow energy dep.?**

Single-channel chiral unitary approach

To apply the argument on Z , we study the **bound state** with mass M_B in the **single channel** chiral unitary approach.

- particle masses: M and m , bound state M_B
- Weinberg-Tomozawa interaction
- parameters: coupling strength $C/2f^2$, subtraction “ a ”

Natural renormalization scheme formulae

$$T^{-1} = V_{\text{WT}}^{-1} - G(a_{\text{pheno}}) = V_{\text{natural}}^{-1} - G(a_{\text{natural}})$$

$$V_{\text{natural}} = -\frac{C}{2f^2}(\sqrt{s} - M_T) + \frac{C}{2f^2} \frac{(\sqrt{s} - M_T)^2}{\sqrt{s} - M_{\text{eff}}}$$

$$M_{\text{eff}} = M_T - \frac{16\pi^2 f^2}{CM_T \Delta a}, \quad \Delta a = a_{\text{pheno}} - a_{\text{natural}}$$

If a_{natural} corresponds to the purely composite case, then M_{eff} for a_{pheno} corresponds to the bare mass M_{B0}
 <-- to be checked in the followings

Single-channel chiral unitary approach

We use the **model-independent** formula

$$1 - Z = 2\pi^2 \sqrt{2\mu^3} \frac{g_W^2}{\sqrt{B}}$$

We need to calculate the coupling g and binding energy B

- **condition for the bound state: $M_B = M + m - B$**

$$1 - \frac{C}{2f^2} (M_B - M) G(M_B; a) = 0$$

--> **parameter of the system: (M_B, a) or $(M_B, M_{\text{eff}} = M_{B0})$**

- **coupling constant: residue of the pole at M_B**

$$[g(M_B; a)]^2 = \lim_{W \rightarrow M_B} (W - M_B) T(W) = - \frac{M_B - M}{G(M_B; a) + (M_B - M) G'(M_B)}$$


- **normalization of the amplitude (a kinematical factor)**

$$1 - Z = \frac{M |\bar{q}(M_B)|}{8\pi M_B (M + m - M_B)} [g(M_B; a)]^2 \quad (\text{for small } B = M + m - M_B)$$

Numerical analysis

Compositeness of the bound state in chiral unitary approach

1) B dependence with $M_{B0} \rightarrow \infty$ ($a = a_{\text{natural}}$)



- $M_{B0} \rightarrow \infty$: $Z \sim 0$


- $Z = 0$ at $B = 0$

- large B behavior is not justified by the approximation

Numerical analysis

Compositeness of the bound state in chiral unitary approach


2) M_{B0} dependence with $B = 10$ MeV




- $M_{B0} \rightarrow M_B : Z \rightarrow 1$
- Mass difference of M_{B0} and M_B : self-energy of bare state
 --> large if the composite component is large


Summary

Structure of resonances/bound states

 Natural renormalization scheme
exclude CDD pole contribution from
the loop function to generate **purely
molecule resonance**

[T. Hyodo, D. Jido, A. Hosaka, Phys. Rev. C78, 025203 \(2008\)](#)

 Field renormalization constant Z :
quantitative measure of **compositeness**

 Natural scheme corresponds to **$Z \sim 0$**
--> generated bound state: composite

[T. Hyodo, D. Jido, A. Hosaka, arXiv:1009.5754 \[nucl-th\]](#)