


Reverse construction of initial conditions: from supernovae to progenitors

Yudai Suwa

Center for Gravitational Physics, Yukawa Institute for Theoretical Physics,
Kyoto University



Key observables characterizing supernovae

* **Explosion energy:** $\sim 10^{51}$ erg

* **Ni mass:** $\sim 0.1 M_{\odot}$

measured by fitting
SN light curves

* **Ejecta mass:** $\sim M_{\odot}$

related

* **NS mass:** $\sim 1 - 2 M_{\odot}$

measured by
binary systems

final goal of first-principle (*ab initio*) simulations

Supernova simulation is an initial value problem

stellar evolutionary calculations

$$\rho(r), T(r), Y_e(r), v_r(r)$$

Time integration

Hydrodynamic equations

$$\begin{aligned} \frac{d\rho}{dt} + \rho \nabla \cdot \mathbf{v} &= 0, \\ \rho \frac{d\mathbf{v}}{dt} &= -\nabla P - \rho \nabla \Phi, \\ \frac{de^*}{dt} + \nabla \cdot [(e^* + P) \mathbf{v}] &= -\rho \mathbf{v} \cdot \nabla \Phi + Q_E, \\ \frac{dY_c}{dt} &= Q_N, \\ \Delta \Phi &= 4\pi G \rho, \end{aligned}$$

Solve simultaneously


Neutrino Boltzmann equation

$$\begin{aligned} \frac{df}{cdt} + \mu \frac{\partial f}{\partial r} + \left[\mu \left(\frac{d \ln \rho}{cdt} + \frac{3v}{cr} \right) + \frac{1}{r} \right] (1 - \mu^2) \frac{\partial f}{\partial \mu} \\ + \left[\mu^2 \left(\frac{d \ln \rho}{cdt} + \frac{3v}{cr} \right) - \frac{v}{cr} \right] E \frac{\partial f}{\partial E} \\ = j(1 - f) - \chi f + \frac{E^2}{c(hc)^3} \\ \times \left[(1 - f) \int R f' d\mu' - f \int R (1 - f') d\mu' \right]. \end{aligned}$$

supernova explosions

Uncertainties in stellar evolutionary calculations


Suwa+, ApJ (2016)



Different codes lead to different structure

Even with the *same* code, different (time or space) resolutions lead to different structure

Sukhbold & Woosley (2014)




$$\xi_M = \frac{M/M_\odot}{r_M/1000 \text{ km}}$$


“Compactness parameter”
O’Connor & Ott (2011)

Asteroseismology

Constantino+ 2015




Constantino+ 2016



core helium burning (CHeB) stars

A possibility



Problem reduction

traditional way

supernova explosion

s11.2-R0.0-3D 110ms



Takiwaki+ 2016



stellar evolution

time

new approach

supernova explosion

Q1. *what is the better initial condition for explosion?*


stellar structure

Q2. *Is it possible to produce such structure?*

stellar evolution

Parametric initial conditions

[Suwa & E. Müller, MNRAS, 460, 2664 (2016)]



- M_1 : the edge of the final convection in the radiative core
- M_2 : the inner edge of the convection zone in the iron core
- M_3 : the NSE core
- M_4 : the iron core mass
- M_5 : the base of the silicon/oxygen shell

Parametric initial conditions

[Suwa & E. Müller, MNRAS, 460, 2664 (2016)]


$s(M_r)$ $Y_e(M_r)$

$P(\rho, s, Y_e)$

$$\frac{dP}{dM_r} = -\frac{GM_r}{4\pi r^4}$$


+

$$\frac{dM_r}{dr} = 4\pi r^2 \rho$$



Parametric initial conditions

[Suwa & E. Müller, MNRAS, 460, 2664 (2016)]






Hydrodynamics simulations

[Suwa & E. Müller, MNRAS, 460, 2664 (2016)]

Agile-IDSA: 1D/GR/neutrino-radiation hydro code, publicly available

<https://physik.unibas.ch/~liebend/download/>





Parameter regime beyond evolution models

[Suwa & E. Müller, MNRAS, 460, 2664 (2016)]

Model	S_c	S_1 [k_B /baryon]	S_2	S_5	Y_{ec}	Y_{e3}	ρ_c [10^{10} g cm $^{-3}$]
BC01	0.5	0.63	1.6	4.0	0.415	0.46	2.0
BC02	0.4	0.63	1.6	4.0	0.415	0.46	2.0
BC03	0.6	0.63	1.6	4.0	0.415	0.46	2.0
BC04	0.5	0.53	1.6	4.0	0.415	0.46	2.0
BC05	0.5	0.73	1.6	4.0	0.415	0.46	2.0
BC06	0.5	0.63	1.5	4.0	0.415	0.46	2.0
BC07	0.5	0.63	1.7	4.0	0.415	0.46	2.0
BC08	0.5	0.63	1.6	3.0	0.415	0.46	2.0
BC09	0.5	0.63	1.6	6.0	0.415	0.46	2.0
BC10	0.5	0.63	1.6	4.0	0.411	0.46	2.0
BC11	0.5	0.63	1.6	4.0	0.425	0.46	2.0
BC12	0.5	0.63	1.6	4.0	0.415	0.452	2.0
BC13	0.5	0.63	1.6	4.0	0.415	0.47	2.0
BC14	0.5	0.63	1.6	4.0	0.415	0.46	1.0
BC15	0.5	0.63	1.6	4.0	0.415	0.46	3.0
BC16	0.4	0.73	1.6	4.0	0.415	0.46	2.0
BC17	0.4	0.63	1.7	4.0	0.415	0.46	2.0
BC18	0.4	0.63	1.6	6.0	0.415	0.46	2.0
BC19	0.4	0.63	1.6	4.0	0.425	0.46	2.0
BC20	0.4	0.63	1.6	4.0	0.415	0.47	2.0
BC21	0.4	0.63	1.6	4.0	0.415	0.46	1.0
BC22	0.4	0.63	1.6	4.0	0.415	0.46	3.0

Explosions in 1D

[Suwa & E. Müller, MNRAS, 460, 2664 (2016)]



Summary

Question:

How can we produce strong ($E_{\text{exp}} \sim 10^{51}$ erg) explosion?

Possible Answer:

Change initial conditions. By starting from specific initial conditions, strong explosions are obtained *without* any change of simulation codes.

Next Question:

Which kind of stellar evolutionary calculations can produce these *perforable* presupernova structure?