

非可換ソリトンと無限個の保存量

浜中 真志

(名古屋大学 多元数理)

Ref: MH, “Commuting Flows and Conservation Laws for NC Lax Hierarchies,”[[hep-th/0311206](#)]

URL: <http://www2.yukawa.kyoto-u.ac.jp/~hamanaka>

1. Introduction

Successful points on NC theories

- Appearance of new physical objects
- Description of real physics
- Various successful applications to D-brane dynamics etc.

NC Solitons play important roles
(Integrable!)

Final goal: NC extension of all soliton theories

Ward's observation:

Almost all integrable equations are
reductions of the ASDYM eqs.

R.Ward, Phil.Trans.Roy.Soc.Lond.A315(85)451

Anti-Self-Dual Yang-Mills eq.

↓ Reductions

KP eq. BCS eq.
KdV eq. Boussinesq eq.
NLS eq. mKdV eq.
sine-Gordon eq. Burgers eq. ...
(Almost all !)

e.g. [Mason&Woodhouse]

NC Ward's observation: Almost all
NC integrable equations are
reductions of the NC ASDYM eqs.

MH&K.Toda, PLA316(03)77[hep-th/0211148]

NC ASDYM eq.

Successful

↓ Reductions

Reductions

NC KP eq. NC BCS eq.

NC KdV eq. NC Boussinesq eq.

NC NLS eq. NC mKdV eq.

NC sine-Gordon eq. NC Burgers eq. ...

(Almost all !?)

Successful?

A general framework is needed

2. NC Sato's Theory

- Sato's Theory : one of the most beautiful theory of solitons
 - Based on the existence of hierarchies and tau-functions
- Sato's theory reveals essential aspects of solitons:
 - Construction of exact solutions
 - Structure of solution spaces
 - Infinite conserved quantities
 - Hidden infinite-dim. symmetry

Let's discuss NC extension of Sato's theory

Derivation of soliton equations

- Give a Lax operator which is a pseudo-differential operator

$$L := \partial_x + u_2 \partial_x^{-1} + u_3 \partial_x^{-2} + u_4 \partial_x^{-3} + \dots$$

$$u_k = u_k(\underbrace{x^1, x^2, x^3, \dots}_{\text{Noncommutativity is introduced here:}})$$

- Define a differential operator

$$B_m := (L * \dots * L)_{\geq 0}$$

m times

Noncommutativity
is introduced here:
 $[x^i, x^j] = i\theta^{ij}$

- Introduce NC (KP) hierarchy equation:

$$\frac{\partial L}{\partial x^m} = [B_m, L]_*$$

$$\begin{aligned} &\partial_m u_2 \partial_x^{-1} + \\ &\partial_m u_3 \partial_x^{-2} + \\ &\partial_m u_4 \partial_x^{-3} + \dots \end{aligned}$$

$$\begin{aligned} &F_{m2}(u) \partial_x^{-1} + \\ &F_{m3}(u) \partial_x^{-2} + \\ &F_{m4}(u) \partial_x^{-3} + \dots \end{aligned}$$



Star product:

$$f(x) * g(x) := f(x) \exp\left(\frac{i}{2} \theta^{ij} \vec{\partial}_i \vec{\partial}_j\right) g(x)$$

which realizes the
noncommutativity

Each coefficient yields
a differential equation.

Closer look at NC (KP) hierarchy

For m=2

$$\partial_x^{-1}) \quad \partial_2 u_2 = \underline{2u_3'} + u_2''$$

$$\partial_x^{-2}) \quad \partial_2 u_3 = \underline{2u_4'} + u_3'' + 2u_2 * u_2' + 2[u_2, u_3]_*$$

$$\partial_x^{-3}) \quad \partial_2 u_4 = \underline{2u_5'} + u_4'' + 4u_3 * u_2' - 2u_2 * u_2'' + 2[u_2, u_4]_*$$

⋮

Infinite kind of fields are represented
in terms of one kind of field $u_2 \equiv u$

MH&K.Toda, [hep-th/0309265]

$$u_x := \frac{\partial u}{\partial x}$$

$$\partial_x^{-1} := \int^x dx'$$

For m=3

$$\partial_x^{-1}) \quad \partial_3 u_2 = u_2''' + 3u_3'' + 3u_4'' + 3u_2' * u_2 + 3u_2 * u_2'$$

⋮



$$u_t = \frac{1}{4}u_{xxx} + \frac{3}{4}(u_x * u + u * u_x) + \frac{3}{4}\partial_x^{-1}u_{yy} + \frac{3}{4}[u, \partial_x^{-1}u_{yy}]_*$$

(2+1)-dim.
NC KP equation

and other NC equations
(NC hierarchy equations)

$$u = u(x^1, x^2, x^3, \dots)$$

$\updownarrow \quad \updownarrow \quad \updownarrow$
 $x \quad y \quad t$

etc.

(KP hierarchy) ^{reductions} \rightarrow (various hierarchies.)

- (Ex.) KdV hierarchy

Reduction condition

$$L^2 = B_2(=:\partial_x^2 + u) \quad : \text{2-reduction}$$

gives rise to NC KdV hierarchy

which includes (1+1)-dim. NC KdV eq.:

$$u_t = \frac{1}{4}u_{xxx} + \frac{3}{4}(u_x * u + u * u_x)$$

Note $\frac{\partial u}{\partial x_{2N}} = 0$: dimensional reduction in x_{2N} directions

KP :	$u(x^1, x^2, x^3, x^4, x^5, \dots)$	
	$x \quad y \quad t$: (2+1)-dim.
		\downarrow
KdV :	$u(x^1, \quad x^3, \quad x^5, \dots)$	
	$x \quad t$: (1+1)-dim.

/-reduction of NC KP hierarchy yields wide class of other NC hierarchies

- No-reduction \rightarrow NC KP $(x, y, t) = (x^1, x^2, x^3)$
- 2-reduction \rightarrow NC KdV $(x, t) = (x^1, x^3)$
- 3-reduction \rightarrow NC Boussinesq $(x, t) = (x^1, x^2)$
- 4-reduction \rightarrow NC Coupled KdV ...
- 5-reduction \rightarrow ...
- 3-reduction of BKP \rightarrow NC Sawada-Kotera
- 2-reduction of mKP \rightarrow NC mKdV
- Special 1-reduction of mKP \rightarrow NC Burgers
- ...



Noncommutativity should be introduced into space-time coords

4. Conservation Laws

- Conservation laws: $\overset{\text{time}}{\partial_t} \sigma = \overset{\text{space}}{\partial_i} J^i$ σ : Conserved density

Then $Q := \int_{\text{space}} dx \sigma$ is a conserved quantity.

$$\because \partial_t Q = \int_{\text{space}} dx \partial_t \sigma = \int_{\text{spatial infinity}} dS_i J^i = 0$$

Conservation laws for the hierarchies

$$\overset{\text{time}}{\partial_m} \text{res}_{-1} L^n = \overset{\text{space}}{\partial_x} J + \underbrace{\theta^{ij} \partial_j \Xi_i}$$

I have found the explicit form !

$\text{res}_{-r} L^n$: coefficient
of ∂_x^{-r} in L^n

Noncommutativity should be introduced
in space-time directions only. \rightarrow

$$t \equiv x^m$$

∂_j should be space or time derivative
 \rightarrow ordinary conservation laws !

Hot Results

Infinite conserved densities for NC hierarchy eqs. ($n=1,2,\dots$)

$$\sigma = \text{res}_{-1} L^n + \theta^{mi} \sum_{k=1}^m \sum_{l=n+1}^{n+1+m-k} (-1)^{m+n-k-l} \binom{m-k}{l-n-1} a_{n-l} \diamond \partial_i b_{m-k}^{(m+n-k-l+1)}$$

$$t \equiv x^m \quad L^n = \partial_x^n + \sum_{l=1}^{\infty} a_{n-l} \partial_x^{n-l}, \quad B_m = \partial_x^m + \sum_{k=1}^m b_{m-k} \partial_x^{m-k}$$

\diamond : Strachan's product

$$f(x) \diamond g(x) := f(x) \left(\sum_{s=0}^{\infty} \frac{(-1)^s}{(2s+1)!} \left(\frac{1}{2} \theta^{ij} \tilde{\partial}_i \tilde{\partial}_j \right)^{2s} \right) g(x)$$

We can calculate the explicit forms of conserved densities for wide class of NC soliton equations.

- Space-Space noncommutativity:

NC deformation is slight: $\sigma = \text{res}_{-1} L^n$

- Space-time noncommutativity

NC deformation is drastical:

– Example: NC KP and KdV equations $([t, x] = i\theta)$

$$\sigma = \text{res}_{-1} L^n - 3\theta((\text{res}_{-1} L^n) \diamond u'_3 + (\text{res}_{-2} L^n) \diamond u'_2)$$

meaningful ?

4. Conclusion and Discussion

- We proved the existence of infinite conserved quantities for wide class of NC Lax hierarchies.
- We gave the infinite conserved densities explicitly from the viewpoint of Sato's theory, which suggests that infinite-dim. symmetry would be hidden in the NC (soliton) equations.
- The results show that NC soliton eqs. have very special properties though they include non-linear terms and infinite (time-!) derivatives.
- Of course, there are still many things to be seen in order to confirm the integrability.

Further directions

- Completion of NC Sato's theory
 - Hirota's bilinearization and tau-functions $u = \partial_x^2 \log \tau$
→ hidden symmetry (deformed affine Lie algebras?)
Cf. Dimakis&Mueller-Hoissen, Wang&Wadati...
 - Geometrical descriptions from NC extension of the theories of Krichever, Mulase and Segal-Wilson and so on.
- Confirmation of NC Ward's conjecture
 - NC twistor theory
e.g. Kapustin&Kuznetsov&Orlov, Hannabuss, Hannover group,...
 - D-brane interpretations → physical meanings
- Foundation of Hamiltonian formalism with space-time noncommutativity
 - Initial value problems, Liouville's theorem, Noether's thm,...