

The many contributions of mean-field theory to the understanding of super-heavy element stability.

Peter Möller

Current affiliations: *emeriti Lund University, Consultant MSU*

PRESENTATION KYOTO,

JUNE 16, 2022

Collaborators on this and other projects, see coauthors on papers posted on URL below and publications listed at end.

More details about masses, other projects (beta-decay, fission), associated ASCII data files and figures are at

<http://t2.lanl.gov/nis/molleretal/>

and at list of relevant publications after conclusions.

Stanislav Ulam has remarked:

It is remarkable how a few characters scribbled on a blackboard can change the course of world history.

Feynman:

- I do not care how smart you are
- or how complicated your model is
- If it does not agree with experimental measurements it is wrong!

Letters to the Editor

[The Editor does not hold himself responsible for opinions expressed by his correspondents. Neither can he undertake to return, nor to correspond with the writers of, rejected manuscripts intended for this or any other part of NATURE. No notice is taken of anonymous communications.]

Possible Existence of a Neutron

It has been shown by Bothe and others that beryllium when bombarded by α -particles of polonium emits a radiation of great penetrating power, which has an absorption coefficient in lead of about 0.3 (cm.)^{-1} . Recently Mme. Curie-Joliot and M. Joliot found, when measuring the ionisation produced by this beryllium radiation in a vessel with a thin window, that the ionisation increased when matter containing hydrogen was placed in front of the window. The effect appeared to be due to the ejection of protons with velocities up to a maximum of nearly $3 \times 10^9 \text{ cm. per sec.}$ They suggested that the transference of energy to the proton was by a process similar to the Compton effect, and estimated that the beryllium radiation had a quantum energy of 50×10^6 electron volts.

I have made some experiments using the valve counter to examine the properties of this radiation emitted in beryllium. The valve counter consists of a small ionisation chamber connected to an amplifier, and the sudden production of ions by the entry of a particle, such as a proton or α -particle, is recorded by the deflexion of an oscillograph. These experiments have shown that the radiation ejects particles from hydrogen, helium, lithium, beryllium, carbon, air, and argon. The particles ejected from hydrogen behave, as regards range and ionising power, like protons with speeds up to about $3.2 \times 10^9 \text{ cm. per sec.}$ The particles from the other elements have a large ionising power, and appear to be in each case recoil atoms of the elements.

If we ascribe the ejection of the proton to a Compton recoil from a quantum of 52×10^6 electron volts, then the nitrogen recoil atom arising by a similar process should have an energy not greater than about 400,000 volts, should produce not more than about 10,000 ions, and have a range in air at N.T.P. of about 1.3 mm. Actually, some of the recoil atoms in nitrogen produce at least 30,000 ions. In collaboration with Dr. Feather, I have observed the recoil atoms in an expansion chamber, and their range, estimated visually, was sometimes as much as 3 mm. at N.T.P.

These results, and others I have obtained in the course of the work, are very difficult to explain on the assumption that the radiation from beryllium is a quantum radiation, if energy and momentum are to be conserved in the collisions. The difficulties disappear, however, if it be assumed that the radiation consists of particles of mass 1 and charge 0, or neutrons. The capture of the α -particle by the Be^9 nucleus may be supposed to result in the formation of a C^{12} nucleus and the emission of the neutron. From the energy relations of this process the velocity of the neutron emitted in the forward direction may well be about $3 \times 10^9 \text{ cm. per sec.}$ The collisions of this neutron with the atoms through which it passes give rise to the recoil atoms, and the observed energies of the recoil atoms are in fair agreement with this view. Moreover, I have observed that the protons ejected from hydrogen by the radiation emitted in the opposite direction to that of the exciting α -particle appear to have a much smaller range than those ejected by the forward radiation.

This again receives a simple explanation on the neutron hypothesis.

If it be supposed that the radiation consists of quanta, then the capture of the α -particle by the Be^9 nucleus will form a C^{13} nucleus. The mass defect of C^{13} is known with sufficient accuracy to show that the energy of the quantum emitted in this process cannot be greater than about 14×10^6 volts. It is difficult to make such a quantum responsible for the effects observed.

It is to be expected that many of the effects of a neutron in passing through matter should resemble those of a quantum of high energy, and it is not easy to reach the final decision between the two hypotheses. Up to the present, all the evidence is in favour of the neutron, while the quantum hypothesis can only be upheld if the conservation of energy and momentum be relinquished at some point.

J. CHADWICK.

Cavendish Laboratory,
Cambridge, Feb. 17.

The Oldoway Human Skeleton

A LETTER appeared in NATURE of Oct. 24, 1931, signed by Messrs. Leakey, Hopwood, and Reck, in which, among other conclusions, it is stated that "there is no possible doubt that the human skeleton came from Bed No. 2 and not from Bed No. 4". This must be taken to mean that the skeleton is to be considered as a natural deposit in Bed No. 2, which is overlaid by the later beds Nos. 3 and 4, and that all consideration of human interment is ruled out.

If this be true, it is a most unusual occurrence. The skeleton, which is of modern type, with filed teeth, was found completely articulated down even to the phalanges, and in a position of extraordinary contraction. Complete mammalian skeletons of any age are, as field palaeontologists know, of great rarity. When they occur, their perfection can usually be explained as the result of sudden death and immediate covering by volcanic dust. Many of the more or less perfect skeletons which may be seen in museums have been rearticulated from bones found somewhat scattered as the result of death from floods, or in the neighbourhood of drying water-holes. We know of no case of a perfect articulated skeleton being found in company with such broken and scattered remains as appear to be abundant at Oldoway. Either the skeletons are all complete, as in the *Stenomylus* quarry at Sioux City, Nebraska, or are all scattered and broken in various degrees, as in ordinary bone beds. The probability, therefore, that the Oldoway skeleton represents an artificial burial is thus one that will occur to palaeontologists.

The skeleton was exhumed in 1913, and published photographs show that the excavation made for its disinterment was extensive. It is, therefore, very difficult to believe that in 1931 there can be reliable evidence left at the site as to the conditions under which it was deposited. If naturally deposited in Bed No. 2, the skeleton is of the highest possible importance, because it would be of pre-Mousterian age, and would be in the company of *Pithecanthropus* and the Piltdown, Heidelberg, and Peking men, all of whose remains are fragmentary to the last degree. Of the few other human remains for which such antiquity is claimed, the Galley Hill skeleton and the Ipswich skeleton are, or apparently were, complete. The first of these was never seen *in situ* by any trained observer, and the latter has, we believe, been withdrawn by its discoverer. The other fragments, found long ago, are entirely without satisfactory evidence as to their mode of occurrence.

Nuclear **BINDING ENERGY**

Bethe-Bacher (-Weizäcker) (1936)

$$B(N, Z) =$$

$$+a_V A \quad (\text{Volume energy})$$

$$-a_S A^{2/3} \quad (\text{Surface energy})$$

$$-a_C \frac{Z^2}{A^{1/3}} \quad (\text{Coulomb energy})$$

$$-a_I \frac{(N - Z)^2}{A} \quad (\text{Symmetry energy})$$

$$-\delta(A) \quad (\text{Pairing energy})$$

Bethe and Bacher, Revs. Mod. Phys. **8** (1936) 82

Bethe and Bacher,
Revs. Mod. Phys. **8** (1936) 82 (in §33):

“There remains thus the nucleus containing 8 neutrons and 8 protons, i.e. , ^{16}O , to test the shell structure” hypothesis by means of nuclear energies. It seems in fact that there is ample evidence for a particular stability of ^{16}O , and thus for the individual-particle approximation.”

So, already in 1936, shell-structure, single-particle models, and how they might modify a macroscopic model were in mainstream discussions.

Hahn and Strassman conclusively identified barium in the products after bombarding uranium with neutrons

(Naturwiss. **27** (1939) 11)

Meitner and Frisch proposed that observations of barium in the reaction products were due to nucleus deforming like a drop

(Nature **143** (1939) 239)

Frisch measured (the predicted) fragment high kinetic energies

(Nature **143** (1939) 239)

Bohr and Wheeler Calculated

(Phys. Rev. **56** (1939) 426):

**Nuclear POTENTIAL ENERGY
versus deformation**

$$B(N, Z) =$$

$$+a_V A \quad (\text{Volume energy})$$

$$-a_S A^{2/3} B_S(\beta) \quad (\text{Surface energy})$$

$$-a_C \frac{Z^2}{A^{1/3}} B_C(\beta) \quad (\text{Coulomb energy})$$

$$-a_I \frac{(N - Z)^2}{A} \quad (\text{Symmetry energy})$$

$$-\delta(A) \quad (\text{Pairing energy})$$

Nuclear Deformation Energy

Let the nuclear surface be described by

$$r(\theta, \phi) = R_0 [1 + \alpha_2 P_2(\cos \theta)]$$

The surface energy lowest order Taylor expansion:

$$E_s = E_s^0 \left(1 + \frac{2}{5} \alpha_2^2\right)$$

The Coulomb energy lowest order Taylor expansion

$$E_C = E_C^0 \left(1 - \frac{1}{5} \alpha_2^2\right)$$

The energy at deformation α_2 relative to spherical shape

$$E_{\text{def}}(\alpha_2) = E_C(\alpha_2) + E_s(\alpha_2) - (E_C^0 + E_s^0)$$

If E_{def} is negative then the system has no barrier wrt fission

$$E_{\text{def}}(\alpha_2) = \frac{2}{5} \alpha_2^2 E_s^0 - \frac{1}{5} \alpha_2^2 E_C^0 < 0$$

$$1 < \frac{E_C^0}{2E_s^0} = x$$

The surface energy for a sphere

$$E_s^0 = 17.80A^{2/3}$$

The Coulomb energy for a sphere

$$E_C^0 = 0.7103 \frac{Z^2}{A^{1/3}}$$

The fissility parameter x :

$$x = \frac{Z^2}{50.13A}$$

Z	A	x
50	124	0.402
82	208	0.645
92	138	0.709
100	252	0.792
114	298	0.870
125	328	0.950
130	335	1.006

since a rigid or liquid nucleus as a whole would have no orbital momentum in its lowest state.

The scheme proposed by Mayer follows exactly the order in a potential well. It achieves the breaks at the correct places by the assumption of a very strong spin-orbit coupling at high angular momentum values.

A summary of the three schemes is given in Table I. All three schemes give, of course, the empirical shell numbers and a statistical correlation with observed spins and moments. A decision between the schemes may be hoped for through discussion of new data which may tend to tip the scales in a definite direction, or by more theoretical work. Among the latter would be a refined calculation of the effects of the Coulomb forces on the density distribution in a nucleus, improved treatment of the many body problem, and better understanding of the spin-orbit coupling in nuclei.

It should be emphasized that the existence and the characteristics of nuclear shell structure have become now much more clearly established than formerly in spite of the ambiguities in their interpretation. Particularly there is a definite correlation between spin and shell structure. This does not mean necessarily that the individual particle model is better than hitherto assumed. The shell structure in nuclei, is, however, so pronounced an effect that one may hope to obtain an interpretation even on basis of such a crude approximation as the individual particle model.

* This letter has been written on request by the editor of the *Physica Review*, who received the papers, reference 1 and 2, by the same mail.

¹ Eugene Feenberg and Kenyon C. Hammack, *Phys. Rev.* **75**, 1877 (1949).

² L. W. Nordheim, *Phys. Rev.* **75**, 1894 (1949).

³ Maria G. Mayer, *Phys. Rev.* **75**, 1969 (1949).

On Closed Shells in Nuclei. II

MARIA GOEPPERT MAYER

Argonne National Laboratory and Department of Physics,
University of Chicago, Chicago, Illinois

February 4, 1949

THE spins and magnetic moments of the even-odd nuclei have been used by Feenberg^{1,2} and Nordheim³ to determine the angular momentum of the eigenfunction of the odd particle. The tabulations given by them indicate that spin orbit coupling favors the state of higher total angular momentum. If strong spin-orbit coupling, increasing with angular momentum, is assumed, a level assignment different from either Feenberg or Nordheim is obtained. This assignment encounters a very few contradictions with experimental facts and requires no major crossing of the levels from those of a square well potential. The magic numbers 50, 82, and 126 occur at the place of the spin-orbit splitting of levels of high angular momentum.

Table I contains in column two, in order of decreasing binding energy, the levels of the square well potential. The quantum number gives the number of radial nodes. Two levels of the same quantum number cannot cross for any type of potential well, except due to spin-orbit splitting. No evidence of any crossing is found. Column three contains the usual spectroscopic designation of the levels, as used by Nordheim and Feenberg. Column one groups together those levels which are degenerate for a three-dimensional isotropic oscillator potential. A well with rounded corners will have a behavior in between these two potentials. The shell grouping is given in column five, with the numbers of particles per shell and the total number of particles up to and including each shell in column six and seven, respectively.

Within each shell the levels may be expected to be close in energy, and not necessarily in the order of the table, although the order of levels of the same orbital angular momentum and different spin should be maintained. Two exceptions, ${}_{11}\text{Na}^{23}$

with spin 3/2 in stead of the expected $d_{5/2}$, and ${}_{25}\text{Mn}^{55}$ with 5/2 instead of the expected $f_{7/2}$, are the only violations.

Table II lists the known spins and orbital assignments from magnetic moments⁴ when these are known and unambiguous, for the even-odd nuclei up to 83. Beyond 83 the data is limited and no exceptions to the assignment appear.

Up to Z or $N=20$, the assignment is the same as that of Feenberg and Nordheim. At the beginning of the next shell, $f_{7/2}$ levels occur at 21 and 23, as they should. At 28 the $f_{7/2}$ levels should be filled, and no spins of 7/2 are encountered any more in this shell. This subshell may contribute to the stability of Ca^{48} . If the $g_{3/2}$ level did not cross the $p_{1/2}$ or $f_{5/2}$

TABLE I.

Osc. No.	Square well	Spect. term	Spin term	No. of states	Shells	Total No.		
0	1s	1s	1s _{1/2}	2	2	2		
1	1p	2p	1p _{1/2}	4	6	8		
			1p _{3/2}	2				
2	1d	3d	1d _{5/2}	6	12	20		
			1d _{3/2}	4				
			2s	2s			2s _{1/2}	2
3	1f	4f	1f _{7/2}	8	22	28?		
			1f _{5/2}	6				
			2p _{3/2}	4				
			2p _{1/2}	2				
4	2p	3p	1g _{9/2}	10	32	50		
			1g _{7/2}	8				
			2d _{5/2}	6				
			2d _{3/2}	4				
			3s	3s			3s _{1/2}	2
5	1h	6h	1h _{11/2}	12	44	82		
			1h _{9/2}	10				
			2f _{7/2}	8				
			2f _{5/2}	6				
			3p _{3/2}	4				
			3p _{1/2}	2				
6	1i	7i	1i _{13/2}	14	126	126		
			1i _{11/2}	12				
			2g	6g			6g	6
			3d	5d			4s	4s

levels, the first spin of 9/2 should occur at 41, which is indeed the case. Three nuclei with N or $Z=49$ have $g_{9/2}$ orbits. No s or d levels should occur in this shell and there is no evidence for any.

The only exception to the proposed assignment in this shell is the spin 5/2 instead of 7/2 for Mn^{55} , and the fact that the magnetic moment of ${}_{27}\text{Co}^{59}$ indicates a $g_{7/2}$ orbit instead of the expected $f_{7/2}$.

In the next shell two exceptions to the assignment occur. The spin of 1/2 for Mo^{95} with 53 would be a violation, but is experimentally doubtful. The magnetic moment of Eu^{153} indicates $f_{5/2}$ instead of the predicted $d_{5/2}$. No $h_{11/2}$ levels appear. It seems that these levels are filled in pairs only,

Swiatecki (and others) observed that experimental actinide spontaneous-fission half-lives differed substantially from what could be explained from smoothly varying (with neutron number and proton number) liquid drop barriers.

He correlated the differences with differences between liquid-drop ground state masses and measured masses and found that such ground-state “shell structure” could account for the observed behavior of actinide spontaneous fission half-lives.

trend is consistent with a straight line, defining $(Z^2/A)_e = 40.2 \pm 0.7$. The equation of the line leads to the semiempirical formula,

$$M_2 - M_1 = 0.090(40.2 \pm 0.7 - Z^2/A)^{1/2}A. \quad (4)$$

One may combine Eq. (4) with the relation:

$$M_2 + M_1 = A - \nu$$

(ν = number of neutrons emitted in fission),

to predict the positions of the peaks in the yield curves of elements that have not yet been investigated. If an average value $\bar{\nu} = 2.8$ is used, one finds

$$M_2 = \frac{1}{2}A - 1.4 + 0.045(40.2 \pm 0.7 - Z^2/A)^{1/2}A, \quad (5)$$

$$M_1 = \frac{1}{2}A - 1.4 - 0.045(40.2 \pm 0.7 - Z^2/A)^{1/2}A. \quad (6)$$

The present analysis provides a reason for the empirical observation that in the fission of different elements the position of the heavy peak remains

TABLE I. Positions of the peaks in the fission yield curves.

Compound nucleus	Position of peaks				Remarks	Reference
	Observed ^a	M_1	Formulas (5), (6)			
	M_2	M_1	M_2	M_1		
Th ²³³	140	91	139.1	91.1	Low-energy neutron fission	b
U ²³⁹	140	98	141.1	95.1		c
U ²³⁶	138.5	95	138.2	95.0		d
U ²³⁴	137	93	136.2	95.0		b, e
Pu ²⁴⁰	138	99	137.9	99.3		c
U ²³⁸	140	96	140.2	95.0	Spontaneous fission	f, g
Cm ²⁴²	136	103	134.7	104.5		g
Cf ²⁵²	139	108	140.2	109.0		h

^a The uncertainty in the observed values of M_2 and M_1 is of the order of ± 1 or ± 2 mass units. (It is more in the cases of U²³⁹ and U²³⁸.) No systematic attempt has been made to adjust $A - M_2 - M_1$ to agree with available information on the number of emitted neutrons.

^b A. Turkevich and J. B. Niday, Phys. Rev. **84**, 52 (1951).
^c E. B. Steinberg and M. S. Freedman, *Radiochemical Studies: The Fission Products* (McGraw-Hill Book Company, Inc., New York, 1951), Paper No. 219, National Nuclear Energy Series, Plutonium Project Record, Vol. 9, Div. IV, Part V.

^d Glendenin, Steinberg, Ingraham, and Hess, Phys. Rev. **84**, 860 (1951).

^e Steinberg, Glendenin, Ingraham, and Hayden, Phys. Rev. **95**, 867 (1954).

^f G. W. Wetherill, Phys. Rev. **92**, 907 (1953).

^g E. P. Steinberg and L. E. Glendenin, Phys. Rev. **95**, 431 (1954).

^h E. P. Steinberg and L. E. Glendenin, J. Inorg. Nuc. Chem. **1**, 45 (1955).

approximately constant. If the degree of asymmetry remained unchanged from nucleus to nucleus, both peaks would move towards higher masses with increasing A . In fact, there is superimposed on this shift a coming together of the peaks with increasing Z^2/A . Since the over-all trend of Z^2/A is to increase with A , the result is that for the light peak the two shifts add up whereas for the heavy peak they partly cancel. This is illustrated in Table I, where M_2 and M_1 , calculated according to (5) and (6), are compared with the observed values.

Further measurements of fission asymmetries would be interesting, especially in the region of Z^2/A close to the critical value, where the present considerations suggest a rapid decrease of $M_2 - M_1$.

It is a pleasure to acknowledge stimulating discussions with Professor S. G. Thompson, Dr. A. C. Pappas, and Dr. T. Maris.

¹ N. Bohr and J. A. Wheeler, Phys. Rev. **56**, 426 (1939).

² A. E. S. Green, Phys. Rev. **95**, 1006 (1954).

³ W. J. Swiatecki (to be published).

⁴ D. L. Hill and J. A. Wheeler, Phys. Rev. **89**, 1102 (1953).

Systematics of Spontaneous Fission Half-Lives

W. J. SWIATECKI

Institute for Mechanics and Mathematical Physics and The Gustaf Werner Institute for Nuclear Chemistry, Uppsala, Sweden

(Received July 18, 1955)

SEVERAL authors have noted the over-all trend of spontaneous fission half-lives to decrease with increasing Z^2/A as well as the considerable deviations (by several powers of 10) from any smooth dependence on this parameter.¹ We should like to discuss the close correlation which seems to exist between the half-lives and the finer details in the systematics of the ground-state masses of nuclei.²

A simple way of exhibiting this correlation is to plot the deviation $\delta\tau$ from a straight line in a plot of τ [$\tau = \log_{10}(\text{half-life})$] vs Z^2/A , against deviations (δM) of the masses M of the nuclei from a smooth reference surface $M_{\text{ref}}(A, Z)$. We made such a plot, with M_{ref} taken to be the semiempirical mass surface of Green³ (based on the liquid drop model):

$$\begin{aligned} \delta M &= M - M_{\text{ref}}, \\ M_{\text{ref}} &= 1000A - 8.3557A + 19.120A^{\frac{2}{3}} \\ &\quad + 0.76278Z^2/A^{\frac{1}{3}} + 25.444(N - Z)^2/A \\ &\quad + 0.420(N - Z) \text{ millimass units.} \quad (1) \end{aligned}$$

The experimental masses M were taken from Glass *et al.*⁴

In the case of even-even nuclei the plot of $\delta\tau$ vs δM suggested a series of straight lines, one for each Z , indicating that for the isotopes of one element special stability of a nucleus (small δM) is invariably associated with a longer lifetime (large $\delta\tau$). The lines had approximately the same slope, thus defining a spontaneous-fission hindrance factor which corresponds to about 10^6 times longer lifetime for each millimass unit of extra stability. This suggested that if the observed lifetimes were corrected for the variations in stability of the ground states, a more regular dependence of τ on Z^2/A might be discernible.

Figure 1 shows the effect on the plot of τ vs Z^2/A of adding to the observed τ_{exp} an empirical correction $k\delta M$ ($k \sim 5$ if δM in mMU). For even-even nuclei the values of $\tau_{\text{exp}} + k\delta M$ define a fairly smooth curve, with indications of a similar curve for odd- A nuclei. [In a

preliminary plot the hindrance factor k was taken to be 5. A small but significant further smoothing of the points resulted from making k vary with Z^2/A according to $k=5-(Z^2/A-37.5)$. This is the case shown in Fig. 1.]

The result can be stated in the form of an empirical formula for half-lives; e.g., for even-even nuclei,

$$\tau_{ee} = f(Z^2/A) - k\delta M, \quad (2)$$

where f is the curve defined by the even-even points in Fig. 1. The relation of the points for odd- A nuclei to the curve obtained from (2) by a shift upwards of 6.6 units is also shown in Fig. 1. The lifetime of the odd-odd nucleus E^{254} (einsteinium, $Z=99$) is consistent with a further shift of 4.9 units. The curve $f(Z^2/A)$ can be




FIG. 1. Plot of spontaneous fission half-lives against Z^2/A . The observed lifetimes τ_{exp} occupy the bottom left-hand part of the figure; the "corrected" values $\tau_{\text{exp}} + k\delta M$ group themselves around the three curves. Experimental points for even-even nuclei are joined by straight lines. Odd- A nuclei are designated by special symbols which, reading from left to right along the odd- A curve, refer to U^{235} , Pu^{239} , Bk^{249} , Cf^{249} , E^{253} (einsteinium, $Z=99$), and Fm^{255} (fermium, $Z=100$). The odd-odd nucleus E^{254} is marked by a square.

represented for example by a cubic, which leads to the following formulas for the lifetimes:

$$\left. \begin{array}{l} \tau_{ee} = 18.2 \\ \tau_{\text{odd } A} = 24.8 \\ \tau_{oo} = 29.7 \end{array} \right\} - 7.8\theta + 0.35\theta^2 + 0.073\theta^3 - (5-\theta)\delta M, \quad (3)$$

where $\theta = (Z^2/A) - 37.5$, and δM is the deviation in mMU of the experimental mass from the surface (1). Table I compares the observed half-lives with the values calculated by means of (3). The remarkable

TABLE I. Values of $\log_{10}(\text{half-life})$.

Nucleus	Experi- mental ^a	Formula (3)	Nucleus	Experi- mental ^a	Formula (3)
Even-even nuclei			Even-even nuclei		
Th 230	≥ 7.18	19.39	Cf 246	3.32	3.27
232	18.15	18.84	248	3.85	3.92
U 232	13.90	13.56	250	4.18	4.24
234	16.30	15.98	252	1.82	1.60
236	16.30	15.21	254	-0.70	-1.02
238	15.90	15.52	Fm 254	-0.30	-0.85
Pu 236	9.54	9.66	256	-3.52	-3.02
238	10.69	11.57	Odd- A nuclei		
240	11.08	11.09	U 235	17.26?	18.02
242	10.86	11.22	Pu 239	15.74	15.42
244	10.40	10.13	Bk 249	8.78	8.67
Cm 244	6.28	6.27	Cf 249	9.18	8.65
242	6.86	7.27	E 253	5.48	4.38
244	7.15	7.09	Fm 255	1.30	2.79
246	7.48	7.88	Odd-odd nuclei		
			E 254	5.18	5.17

^a The experimental values are from a summary by A. Ghiorso, kindly lent to me by Professor S. G. Thompson.

degree of smoothing achieved by means of the unsophisticated correction $k\delta M$ is illustrated by the fact that the deviations from (3) rarely exceed 0.5. (Note that a shift in τ of this amount would be produced by an error of 0.1 mMU in δM .)

The importance of shell structure in the fission process is suggested by the fact that, according to the present considerations, the oscillations of the masses (associated with individual particle structure) in the range $\delta M = 1-3$ mMU shorten the lifetimes by factors of 10^5 to 10^{15} . On the other hand the irregularities in the original plot of τ_{exp} against Z^2/A are seen to be largely due to irregularities in the ground-state masses, associated with *shell structure in the ground-state configuration*. The smoothness of the points $\tau_{\text{exp}} + k\delta M$ suggests that, after correcting for shell structure in the ground-state configuration, the description of the fission process in terms of a model in which single-particle features are treated in an average way may be useful. Qualitative reasons for the greater validity of such an averaged description for the more strongly deformed nuclear shapes occurring in fission may be found in the disappearance for such shapes of degeneracies in the energy spectrum associated with the proximity to a spherically symmetric configuration.

It is a pleasure to acknowledge discussions with Professor S. G. Thompson and Dr. A. C. Pappas and stimulating contacts with Dr. Aage Bohr and Dr. B. R. Mottelson and members of the C.E.R.N. Theoretical Study Group in Copenhagen.

¹ See for example J. R. Huizenga, Phys. Rev. **94**, 158 (1954).

² The existence of correlations between nuclear masses, fission thresholds, and half-lives has been considered by Professor D. Frisch, to whom I am greatly indebted for stimulating discussions.

³ A. E. S. Green, Phys. Rev. **95**, 1006 (1954).

⁴ Glass, Thompson, and Seaborg, J. Inorg. Nuc. Chem. **1**, 3 (1955).

CERN 55 - 30

THEORETICAL STUDY DIVISION

Det Kongelige Danske Videnskabernes Selskab

Matematisk-fysiske Meddelelser, bind 29, nr 16

Dan Mat Fys Medd 29, no 16 (1955)

BINDING STATES
OF INDIVIDUAL NUCLEONS IN
STRONGLY DEFORMED NUCLEI

BY


SVEN GÖSTA NILSSON



København 1955

i kommission hos Ejnar Munksgaard

There is a scale for both of the deformation variables η and δ . To first order δ equals the deformation parameter β used in the papers of A. Bohr and B. Mottelson. The energy is plotted in the A - and δ -dependent energy unit $\pi h \omega_0(\delta)$, where $h \omega_0$ varies with A as $A^{-1/3}$. Further, for $A \approx 100$ one expects $h \omega_0 \approx 8.8$ MeV. The constant π is chosen equal to 0.05 in the diagram. The levels are labelled by the J^π -number and the parity. The numbers to the left of the curves refer to Table I.



Are the heaviest actinides ending the periodic system of observable elements? Scharff-Goldhaber mentions in *Nucleonica* already in 1957 that *There may be for instance, another region of relative stability at the doubly-magic nucleus ${}_{126}X^{310}$ (the closing of the j neutron shell)*”

Since she mentions it so casually this possibility was probably always recognized by the community. Exactly which nuclei might be stable was overlooked in calculations for a further 10–25 years, see below.

nuclear physics

By GERTRUDE SCHARFF-GOLDHABER

					13	Al 26.98 #13
					12	Mg 24.32 #12
					11	Na 22.991 #11
					10	Ne 20.183 #10
						Ne 18 16 s #*3.2 E 4.2
						Ne 19 18.5 s #*2.2 E 3.2
					9	F 19.00 #9
						F 17 66 s #*1.75 E 2.77
						F 18 1.87 h #*65 E 1.87
					8	O 16.000 #8
						O 14 72 s #*1.83 #2.30 E 3.5
						O 15 2.1 m #*1.7 E 2.7
						O 16 99759 #*4.0003 E 0.00005
						O 17 0.037 #*4 E 0.00483
						N 12 0.012 s #*16.7 (3e-4) E 17.7
						N 13 10.0 m #*120 E 2.22
						N 14 99.63 #*1.7 E 0.00752
						N 15 0.37 #*0.0002 E 0.00488
						N 16 7.4 s #*4.04 #6.37022 E 10.4
						C 10 19 s #*13 #72,103 E 3.8
						C 11 20.5 m #*98 #1 E 1.90
						C 12 98.89 #*0.0022 E 0.00090
						C 13 1.11 #*1008 E 0.00746
						C 14 5600 y #*158 #7 #0.10 E 1.58
						C 15 2.3 s #*43.99 #53 E 9.8
						B 9
						B 10 1.8
						B 11
						B 12 1.025 s

EVEN A SUPERFICIAL GLANCE backward will teach us that it is impossible to predict in detail the future of a fast-moving science like nuclear physics. It will remind us that often entirely unexpected events changed the direction of endeavor in this field. These events were either of an experimental nature, as, for instance, the finding that beta rays have continuous energy spectra, or they consisted in the formation of new concepts—as, for example, of the liquid-drop model of the nucleus.

It is, of course, similarly impossible to predict what extraneous happenings may in the future affect scientific progress as profoundly as two world wars and political persecution have affected it in the past. During the last twelve years the great importance attached to atomic energy has induced an unprecedented increase in the tempo of research, and new nuclear physics centers have sprung up all over the world. This development is viewed by many with delight while others are afraid that it may have a negative effect on nuclear physics as a pure science.

In spite of all the uncertainties mentioned, it is useful to interrupt from time to time one's preoccupation with the problem at hand to investigate the trends that current research seems to follow, both in experiment and theory, and to try to recognize how far these may serve to bring us closer to the solution of outstanding problems.

The central problem is to understand the nucleus in the same sense in which one might have said in 1926 that the atom was understood: one knew then not only that the forces between the nucleus and the atomic electrons were pure Coulomb forces, but also that the excited states in which the system could exist were governed by the laws of quantum mechanics, including the Pauli exclusion principle. (It is true, only the states of the simplest atom, hydrogen, could be exactly calculated, while already the helium atom presented such overwhelming mathematical difficulties that it took about another quarter century and development of computers to reach the same stage.)

If we want to study events happening inside the nucleus, its finite extension and the distribution of charge and current inside it, we have a task of a higher degree of difficulty than in the atomic case: (a) We do not know the exact nature of the force between two nucleons, nor do we know whether a potential exists for this force. (b) One cannot consider, in first approximation, the interaction of two nucleons only, because all nucleons are close to each other. (c) We do not know whether the laws of quantum mechanics are sufficient to describe a nucleus completely.

Quantum mechanics may need to be modified, e.g., by the introduction of the concept of a fundamental length, before nuclear phenomena can be explained

On the other hand, we have two important clues on the nature of nuclear forces:

1. Apart from the lightest ones, all nuclei have the same density $\rho = 1.7 \times 10^{28}$ nucleons/cm³, at least in their central part, so that it is reasonable to speak of "nuclear matter." Hence, if it were not for the Coulomb repulsion between the protons, nuclei of arbitrarily large size would exist. We therefore speak of the "saturation" of nuclear forces, which prevents nuclear matter from collapsing to a density less than ρ and from flying apart.

2. Experiments have shown that the forces between two protons, corrected for the effect of Coulomb repulsion, are the same as between two neutrons, i.e., charge symmetry prevails, and probably the forces are also the same between a proton and a neutron (charge independence).

In recent years a number of theoretical physicists, under the leadership of K. Brueckner, have tried to understand the nuclear phenomena by treating the nucleus as a many-body system assuming that nuclear forces can be derived from a two-body potential. The simplified case of infinite

	#138-7.2 E14	#3.24 E4.26	#23 28.99008	#2.87 E4.65	
	Mg 23 12 s #73.0 E4.0	Mg 24 78.8 #03 23.97254	Mg 25 10.1 #27 24.99375	Mg 26 11.1 #03 25.99080	Mg 27 9.5 m #175.157 #84.102.18 E2.59
Na 21 25 s #130 E1.92	Na 22 2.6 y #54.4 E2.84	Na 23 100 #53 22.99705	Na 24 15.0 h #138.4 #2753.568 E5.51	Na 25 60 s #42.2.6.3.4 #98.58.38 E4.0	
Ne 20 40.8 #977	Ne 21 0.26 Z100060	Ne 22 8.9 Z189835	Ne 23 40 s #42.3.6. #44.165. E4.2	Ne 24 3.4 m #190.1.4.3 #470.88. E2.42	
F 19 190 #91 #3446	F 20 11 s #342 #163 E7.05	F 21 5 s # E5.7	14		
O 18 0.204 #3688 #369486	O 19 29 s #32.4.4 #370.37.110 E4.75		By permission of General Electric Co.		
N 17 4.14 s #17 #101 E4.8			12		

nuclear matter is considered first. A few months ago P. S. Signell and R. E. Marshak and independently J. Gammel and R. Thaler showed that the phase shifts derived from nucleon-nucleon scattering experiments up to 150 Mev can be fitted remarkably well by a charge-independent potential including a spin-orbit term of the same sign as in shell theory. This potential has already been introduced in a Brueckner-type theory by de Dominicis, and reasonable answers for the binding energy of infinite nuclear matter have been obtained.

One would hope to apply similar methods to the more complicated case of the finite nucleus, thereby demonstrating the validity of the shell model for such a system. The next decade may see an understanding of complex nuclei based on interactions derived from nucleon-nucleon scattering experiments.

While the solution of these very difficult fundamental problems progresses slowly, experimental nuclear physicists are guided in their research by a number of more phenomenological nuclear models, each one of which is limited in its application. Usually these models are not born in a finished form but have to be modified continuously to fit the facts. Sometimes, they even seem to contradict fundamental principles, as when the basic assumption of the shell model that a nucleon may be considered to move in a central potential seemed at first to violate the idea that the mean free path of a nucleon is of the order of its diameter. J. H. D. Jensen, in a talk given in 1956 at the International Congress on Theoretical Physics in Seattle, recounted how Maria Mayer and independently Haxel, Jensen and Suess revived and modified—by the introduction of strong spin-orbit coupling—the early shell-model ideas (which in turn were conceived in analogy to the ideas (which in turn were conceived in analogy to the atomic case) merely as a working hypothesis. A possibility of removing the apparent contradiction was first

pointed out by Weisskopf, who emphasized the importance of the role of the exclusion principle in simplifying the particle motions in the presence of strong forces. Later quantitative studies of Brueckner, Eden and Bethe have shown that this conjecture is correct and that the shell model can be used to give a useful first approximation to the nucleus up to ~ 10 Mev.

Two other models which have played an important role in recent years are the collective model developed by A. Bohr and B. Mottelson, which is likewise valid at low energies only, and the optical model, which successfully describes nuclear absorption and scattering processes, e.g., the so-called mountain resonances for neutrons and protons.

Let us now consider some of the aspects of these models which are likely to lead to further progress in nuclear research: The shell model will, of course, continue to be useful for the construction of level schemes and in giving the characters (spins and parities) of levels not only of stable nuclei, but also of radioactive nuclei, in providing a means for the classification of beta decays and for the explanation of the occurrence of nuclear isomers, etc. Of more far-reaching importance are the possibilities for a quantitative analysis of the energies of nuclear levels. In the near future such an analysis will be restricted to the immediate neighborhood of doubly magic nuclei, e.g., O^{16} and Pb^{208} .

The successful analysis by Elliott and Flowers, by D. Kurath and by B. French and others not only of the level energies of some light nuclei, but also of the transition rates of a number of beta transitions, and of several electromagnetic transitions—mainly of dipole character—justifies the expectation of further progress in this field.

As an example of a field where much progress can be expected in the near future, and in which the author has been particularly interested, a few words may be said about those even-even nuclei which have vibrational level schemes and which lie between the "magic" nuclei and the rotational region. The vibrational level schemes have been tentatively interpreted on the basis of the Bohr-Mottelson model in the region of weak to moderate coupling. Such a model predicts a triplet, of characters 0^+ , 2^+ , 4^+ , at about twice the energy of the first 2^+ state. Since states differing in spin by 4 would escape detection with most of the usual methods of determining level schemes, it is not



G. Scharff-Goldhaber

Physics Department, Brookhaven National Laboratory; has worked on nuclear energy levels, neutron and electron diffraction, spontaneous fission, and ferromagnetism.

surprising that only one such close-lying triplet has been found so far, namely, in Cd¹¹⁴. The method used here was the analysis of conversion electrons accompanying neutron capture in Cd¹¹³ (H. Motz). A systematic search for triplets by means of suitably chosen experiments should throw light on the question of the nature of the vibrational even-even nuclei and, in turn, of those odd-*A* nuclei of which they form the core.

As is well known, the collective model has been enormously successful in describing not only the level characters and energy ratios, but also the transition probabilities of electromagnetic transitions, the $\log ft$ values, etc., in strongly deformed nuclei.

Recent efforts to interpret the level scheme of F¹⁹ have led to an interesting discovery: the results of one group of physicists who applied a shell-model analysis agreed surprisingly well with those of another group who applied a collective-model interpretation. Ironically, the theoretical values agreed even more closely among each other than with the experimental values. The reason for the good agreement is by no means obvious and is now being studied by a number of theoreticians. It allows one to conclude, however, that it will be possible to set up a unified model of the nucleus, in which the individual particle motions and the collective motions are self-consistent as in molecules. Important attempts in this direction have been made by Peierls and Yoccoz and by Wheeler and Griffin.

Now a few words about higher-energy nuclear reactions: As was mentioned above, the optical model is successful in describing the energy dependence of the absorption and scattering cross sections of the particles impinging on a nucleus for a considerable energy range. However, a theory has still to be evolved which will give the probabilities and angular correlations for the decomposition of the system target nucleus plus bombarding particle into the various energetically allowed end products. It may be added that the optical model is based on the assumption that the energy spectrum of the incoming particles overlaps many resonance levels in the target nucleus. It is likely that with increasing energy definition this approach will give way to a renewed interest in the fine structure, which is of particular importance in the fission process.

We can look forward to definitive progress toward understanding the bewildering variety of phenomena observed in fission

A few thoughts on beta-decay theory: the recent revolution in thought brought about by the discovery that parity conservation and the conservation of charge conjugation do not hold in weak interactions has attracted great interest to this field. Although at present the nature of the interactions for the nucleon-electron-neutrino system is not known, it is very probable that by the end of the coming decade it will be quite well understood. This will be brought about by studies of polarization of the electrons emitted in various types of beta decay, by further efforts to detect double beta decay and by the refinement of present neutrino-detection experiments.

The experimental determination of nuclear properties will increase in accuracy and scope as the equipment and methods grow in diversity and ingenuity.

For example, the steady improvement in resolution and efficiency of spectrometers will facilitate the determination of level energies.

For character assignments to nuclear energy levels there are now a number of methods at our disposal, which we have just begun to exploit. For short-lived states preceded by some previous radiation there is the delayed-coincidence method, which Sunyar has recently developed into a form that permits one to measure lifetimes as low as a few times 10⁻¹¹ sec. For electric dipole or quadrupole transitions leading to the ground state the Coulomb-excitation method is able to cover a wide lifetime range. As a final example, the use of molecular-beam and paramagnetic-resonance methods for the spin determination of radioactive nuclei will doubtless increase and serve to check assignments made on the basis of decay-scheme studies. Further, it may be expected that the refinement of the theoretical interpretation of experimental results, e.g., a better theory of stripping, of internal conversion and of the angular distribution of inelastically scattered particles, will make level-character assignments of excited states more definite.

Radioactive nuclei will be found further away from the stability region, owing to the use of heavier bombarding particles and faster detection methods. The study of the binding energies of nuclei of this type may throw further light on the nature of nucleon-nucleon forces. Also, the number of known elements will certainly be increased.

Relatively long-lived isotopes may well be found among the far-transuranic nuclides because of magic-number stability

There may be, for instance, another region of relative stability at the doubly magic nucleus ¹²⁶X³¹⁰ (the closing of the *j* neutron shell).

New accelerators like variable-energy cyclotrons and tandem Van de Graaffs capable of producing beams of well defined and sufficiently high energies will make it possible to study radiation and particle widths of excited states of light nuclei with accuracy, thus testing the shell-model wave functions, and to explore the level schemes of medium-weight nuclei (50 < *A* < 150).

Atomic-beam methods and possibly the study of μ -mesic X-rays will yield new data on electric and magnetic moments of higher order. The boldness of some thinkers in this field is indicated by the title of a recent theoretical paper: "Nuclear Hexadecapole Moments." It may even be possible to get a better idea of the charge and current distribution within the proton by means of the 6-Bev electron synchrotron now being constructed at Cambridge, Massachusetts, which will permit an extension of the very successful work carried out at Stanford.

One important goal of nuclear-physics research is the deduction of nuclear forces from meson fields. Among these fields the role of the π -meson field will probably be first understood, but it is clear that any ultimate theory will not be able to ignore the role of the "strange particles" (*K* mesons and hyperons). In the meantime, the new field of "hypernuclear" physics is likely to develop considerably and to help indirectly in the understanding of nuclear phenomena.

In Nucl. Phys. **81** (1966) 1, Myers and Swiatecki present a mass table based on a postulated shell-correction expression with $Z = 126$ as the next magic proton number beyond $Z = 82$. However they state that Meldner suggested to them that $Z = 114$ was another possibility. They also observed that in previously published Nilsson modified-oscillator level diagrams there were large spherical gaps at this proton number (but since they were not labeled their significance was not always recognized).

ARKIV FÖR FYSIK

UTGIVET AV
KUNGL. SVENSKA VETENSKAPSAKADEMIEN

Band 36 nr 69

PROCEEDINGS OF THE LYSEKIL SYMPOSIUM 1966

C. GUSTAFSON, I. L. LAMM, B. NILSSON and S. G. NILSSON

Nuclear deformabilities in the
rare-earth and actinide regions with excursions
off the stability line and into the
super-heavy region



ALMQVIST & WIKSELL

STOCKHOLM

1967




Fig. 3. Single-proton levels for $Z > 82$. $\kappa = 0.0577$, $\mu = 0.65$. Note added in proof: Level denoted [510 $1/2$] in middle of figure should be [530 $1/2$].

On the other hand, for a more complicated r -dependence than the one assumed in eq. (1), one may not be able to conserve the volumes of all equipotential surfaces simultaneously. One may then have to be content with conserving a weighted average of these surfaces.

In the actual calculations it is convenient to make a small change of eq. (1) by first transforming to the variables $\xi = x\sqrt{M\omega_1}/\hbar$, $\eta = \dots$, etc., according to ref. [7]. The parameters ϵ_{20} and ϵ_{40} are replaced by ϵ and ϵ_4 . We then write the total single-particle Hamiltonian:

1.D.4:
1.E.2

Nuclear Physics A95 (1967) 420—442; © North-Holland Publishing Co., Amsterdam
Not to be reproduced by photoprint or microfilm without written permission from the publisher

SHELL EFFECTS IN NUCLEAR MASSES AND DEFORMATION ENERGIES

V. M. STRUTINSKY

*I. V. Kurchatov Institute of Atomic Energy
Moscow, USSR*

Received 20 October 1966

Abstract: Nuclear shells correspond to an inhomogeneous distribution of nucleons in phase space, whereas the distribution in quasi-classical phenomenological models (the liquid drop model) is supposed to be homogeneous. Starting from this point, Nilsson's level scheme is used to calculate the shell-model correction to the "liquid drop energy" of the nucleus as a function of the occupation number and deformation. A strong correlation between the shell correction and nucleon level density at the Fermi energy was observed. In magic and mid-shell nuclei the calculated deformation energy oscillates around the LDM value. Discussion of problems related to the deformation energy such as nuclear deformations, shell effects in nuclear masses, in deformed nuclei and in nuclear fission, etc. is presented. The role of nucleon pairing is discussed.

1. Introduction

The problem of nuclear masses and their dependence on deformation is one of the oldest in nuclear theory. There are several approaches to the problem, one of which is the so-called liquid drop model (LDM), which is commonly used in the description of nuclear masses and in nuclear fission theories. The LDM is based on the assumption of a classically uniform distribution of nucleons in phase space and thus ignores completely nucleon shell effects. The importance of such (quantum) effects is stressed however in the Bohr-Mottelson unified model. In more microscopic models, an attempt is made to reduce the problem to residual nucleon interactions, e.g. quadrupole-quadrupole and pairing interactions.

A completely quantitative microscopic description of nuclear deformations is hardly possible at present. Modern theories of nuclear excitations use a renormalized Hamiltonian and assume that the properties of the average field are known from empirical data. The problem of nuclear masses and deformations is complicated by the necessity of calculating the average field and the energy related to it and some other quantities such as the surface tension constant. In the surface region the nucleon density decreases to zero in a very short distance of the order of $A^{-\frac{1}{3}}R_0$, i.e. a large density gradient is present. The problem of surface tension arises for which features of nucleon interactions such as saturation (velocity dependence) are important.

It was assumed that relatively stable nuclei near the next magic number would be separated from the last stable nuclei at the end of the currently known ones by a sea of instability. Some large-scale calculations happened to include some such nuclei (although they were focused on actinides and the next assumed magic numbers). Some nuclei in the sea of instability were actually calculated to have large negative shell corrections below -5 MeV. But the significance was not noted at the time. (next 4 frames).

1.D.2:
2.J

Nuclear Physics A131 (1969) 1–66; © North-Holland Publishing Co., Amsterdam
Not to be reproduced by photoprint or microfilm without written permission from the publisher

ON THE NUCLEAR STRUCTURE AND STABILITY OF HEAVY AND SUPERHEAVY ELEMENTS †

SVEN GÖSTA NILSSON †† and CHIN FU TSANG

Lawrence Radiation Laboratory, University of California Berkeley, California

and

ADAM SOBICZEWSKI, ZDZISLAW SZYMAŃSKI and SLAWOMIR WYCECH

Institute for Nuclear Research and University of Warsaw, Warsaw, Poland

and

CHRISTER GUSTAFSON, INGER-LENA LAMM, PETER MÖLLER and BJÖRN NILSSON

Department of Mathematical Physics, Lund Institute of Technology Lund, Sweden

Received 14 February 1969

Abstract: Nuclear potential energy surfaces as a function of deformations are calculated on the basis of a modified oscillator model. In particular, quadrupole (P_2) and hexadecapole (P_4) deformations are considered. The average behavior of the surface is normalized to that of a liquid drop through the employment of a generalized Strutinsky prescription. In this way a synthesis of the single-particle model and the liquid-drop model is obtained.

Lowest minima in the potential energy surfaces give the ground state masses and distortions. These results compare extremely well with experimental data. Spontaneous fission half-lives are also obtained. The inertial parameters associated with fission barrier penetration are derived empirically as well as by a microscopic model. Shape (fission) isomeric states are also found. Their N and Z dependence in the present model are discussed and results tabulated.

The calculations are extended to the predicted superheavy region around $Z = 114$ and $N = 184$. The total overall stability with respect to alpha and beta decay, and spontaneous fission is found to be most favorable in the vicinity of $Z = 110$ and $N = 184$. Detailed diagrams and tables are exhibited.

Introduction

It was found a long time ago ^{1,2,3}) that simple equilibrium calculations based on the deformable shell model ⁴) were able to reproduce the experimental quadrupole moments in the “rare earth” and “actinide” regions. In the Mottelson-Nilsson calculations ¹) single-particle energies are simply added as a function of the quadrupole distortions, and the shape corresponding to the minimum energy is found. The calculations neglected the effects of Coulomb and pairing interactions, which are, however, considered by Bès and Szymański ²), Szymański ⁵), and also by Sobiczewski ⁶). On the whole, the results of ref. ¹) are reproduced, indicating that Coulomb and pairing forces at the equilibrium point counteract each other. In these latter

† Work supported by the U.S. Atomic Energy Commission, the Swedish Council of Atomic Research and the Polish Atomic Energy Commission.

†† Present address: The Department of Mathematical Physics, Lund Institute of Technology, Lund, Sweden.




Fig. 16. Experimental and theoretical mass values for $150 < A < 340$ plotted relative to the spherical liquid drop value as of ref. ¹¹).

CALCULATED GROUND-STATE PROPERTIES OF HEAVY NUCLEI

P. MÖLLER[†], S. G. NILSSON[†] and J. R. NIX

Los Alamos Scientific Laboratory, University of California^{††}
Los Alamos, New Mexico 87544

Received 10 April 1974

Abstract: Ground-state distortions and single-particle corrections are calculated for nuclei with $Z \geq 68$ and $N \geq 106$ by use of the macroscopic-microscopic method as developed by Strutinsky. The microscopic part is calculated primarily by use of the folded Yukawa single-particle potential. Its parameters are redetermined to fit actinide data. The modified oscillator potential is also used in some of the studies. Two methods for calculating the macroscopic energy are investigated. One is the droplet model of Myers and Swiatecki, and the other is a modified liquid-drop model in which the surface-energy term is modified to take into account the finite range of the nuclear force. Single-particle level diagrams for the folded Yukawa potential are also presented. They are plotted as functions of the distortion parameters ϵ , ϵ_4 and ϵ_6 . Theoretical and experimental single-particle levels at the ground state for actinide nuclei are also compared.

1. Introduction

Today many calculations of nuclear properties are carried out by use of the macroscopic-microscopic method, which was developed in its present form by Strutinsky¹⁾. In this method the total potential energy of the nucleus is expressed as the sum of two terms, a macroscopic term and a microscopic term. A recent review of the calculation of fission barriers²⁾ contains some results obtained by use of the folded Yukawa single-particle potential to calculate the microscopic corrections and the droplet model of Myers and Swiatecki³⁻⁵⁾ to calculate the macroscopic part of the energy. A more detailed study of the results obtained for fission barriers is presented in ref. 6). However, compared to the large distortions involved in fission, much more experimental information is available at ground-state distortions. In this paper we use primarily the folded Yukawa single-particle potential to study single-particle levels, nuclear shapes, and single-particle corrections at ground-state distortions. These quantities depend not only on the single-particle potential but also, to a lesser extent, on the method that is used to calculate the macroscopic energy. We present results obtained by use of the droplet model of Myers and Swiatecki³⁻⁵⁾ and alternatively by use of a modified liquid-drop model in which the surface-energy term is modified to take into account the finite range of the nuclear force⁷⁾. Details on various aspects of the calculations can be found in refs. 2, 7-9) and in references quoted

[†] On leave from the University of Lund, Lund, Sweden.

^{††} This work was supported by the US Atomic Energy Commission and the Swedish Atomic Research Council.

TABLE 2 (continued)

Z	N	A	ε	ε_4	β_2	β_4	Single-particle correction (MeV)
104	162	264	0.208	0.064	0.230	-0.058	-5.11
	142	246	0.207	-0.015	0.224	0.036	-1.74
	144	248	0.212	-0.005	0.229	0.025	-2.21
	146	250	0.215	0.002	0.233	0.018	-2.75
	148	252	0.220	0.011	0.239	0.007	-3.41
	150	254	0.225	0.021	0.245	-0.004	-4.14
	152	256	0.228	0.031	0.250	-0.015	-4.76
	154	258	0.225	0.038	0.246	-0.024	-4.77
	156	260	0.220	0.044	0.241	-0.033	-4.83
	158	262	0.217	0.054	0.238	-0.045	-5.06
106	160	264	0.213	0.060	0.235	-0.053	-5.50
	162	266	0.208	0.070	0.230	-0.066	-6.08
	142	248	0.210	-0.002	0.228	0.021	-1.60
	144	250	0.213	0.005	0.232	0.014	-2.15
	146	252	0.222	0.018	0.242	-0.001	-2.80
	148	254	0.225	0.028	0.246	-0.012	-3.58
	150	256	0.228	0.034	0.250	-0.019	-4.42
	152	258	0.230	0.041	0.252	-0.027	-5.17
	154	260	0.227	0.047	0.249	-0.035	-5.31
	156	262	0.223	0.054	0.246	-0.044	-5.49
108	158	264	0.218	0.060	0.241	-0.052	-5.83
	160	266	0.215	0.067	0.238	-0.061	-6.37
	162	268	0.212	0.077	0.235	-0.073	-7.06
	142	250	0.213	0.015	0.232	0.002	-1.53
	144	252	0.220	0.024	0.240	-0.009	-2.16
	146	254	0.222	0.031	0.242	-0.017	-2.91
	148	256	0.225	0.038	0.246	-0.024	-3.77
	150	258	0.227	0.044	0.249	-0.031	-4.68
	152	260	0.230	0.051	0.253	-0.038	-5.51
	154	262	0.225	0.057	0.248	-0.047	-5.80
108	156	264	0.222	0.064	0.245	-0.056	-6.11
	158	266	0.218	0.070	0.242	-0.064	-6.58
	160	268	0.213	0.073	0.236	-0.069	-7.24
	162	270	0.212	0.080	0.235	-0.077	-8.05

The ground-state single-particle correction is the nuclear ground-state mass relative to the spherical macroscopic energy, which is calculated here by use of the modified-surface-energy model. In this table the ground-state zero-point energy (which is frequently taken equal to 0.5 MeV) is not included. Single-particle levels for $^{228}_{88}\text{Ra}$ and $^{250}_{98}\text{Cf}$ are used to calculate the potential energy for nearby nuclei. In both regions the values $a = 0.8$ fm, $\lambda_n = 36$ and $\lambda_p = 34$ are used.

One then arrives at the following expression for the macroscopic energy in the modified-surface-energy model:

$$E_{\text{macro}} = -a_v(1 - \kappa_v I^2)A + E(\xi_v) + \frac{4}{3}\pi V_0 R_0^3 + \frac{3}{5} \frac{e^2 Z^2}{r_0 A^{\frac{1}{3}}} \left[B_c(\xi_v) - \frac{5}{6}\pi^2 \frac{d^2}{r_0^2 A^{\frac{1}{3}}} - \frac{0.7636}{Z^{\frac{1}{3}}} \right]. \quad (3)$$

In 1975 Seeger and Howard published the first global mass table with deformations and level structure based on *calculated* microscopic correction based on a general nuclear-structure model and the Strutinsky method. No other mass table in this issue is based on a general theoretical nuclear structure model able to model many correlated nuclear properties.

These Numbers Form Volume 17
Volume 17 (1975) Nos. 1-4 (1975) June 1976

Atomic Data AND Nuclear Data Tables

1975 MASS PREDICTIONS

S. Maripuu, Special Editor

EDITOR:

Katharine Way

MANUSCRIPT EDITOR:

Lu W. Rose

ASSOCIATE EDITORS:

E. U. Baranger
B. Bederschi
F. Boehm
J. W. Cooper
A. Dalgarno
S. Datz
B. S. Dzhelepov
R. Geballe
J. A. Harvey
W. C. Lineberger
J. B. Marion
E. W. McDaniel
F. K. McGowan
E. Merzbacher
M. Mladjenovic
G. R. Satchler
S. Shafroth

Introduction	1
Development of the Semiempirical Droplet Model. <i>W. D. Myers</i>	411
A New Semiempirical Shell Correction to the Droplet Model (Gross Theory of Nuclear Magics). <i>H. v. Groot, E. R. Hilf,</i> <i>and K. Takahashi</i>	418
A Semiempirical Liquid-Drop Plus Shell-Correction Formula. <i>P. A. Seeger and W. M. Howard</i>	428
A Semiempirical Shell-Model Formula. <i>S. Liran and N. Zeldes</i>	431
An <i>Ab-Initio</i> Shell-Model Formula. <i>M. Bauer</i>	442
Masses of Spherical Nuclei from the Energy-Density Method. <i>M. Beiner, R. J. Lombard, and D. Mas</i>	450
Updated Mass Predictions from the Garvey-Kelson Mass Relations. <i>J. Jänecke</i>	455
Ensemble Averaging of Mass Values. <i>E. Comay and I. Kelson</i>	463
Masses from Inhomogeneous Partial Difference Equations. <i>J. Jänecke and B. P. Eynon</i>	467
A 1975 Midstream Atomic Mass Evaluation. <i>A. H. Wapstra</i> <i>and K. Bos</i>	474
EXPLANATION OF TABLE	476
TABLE. 1975 Mass-Excess Predictions	477

ACADEMIC PRESS
New York and London

A Subsidiary of Harcourt Brace Jovanovich, Publishers



In 1981 a global mass table based on the folded-Yukawa single-particle model was published. In the summer of 1982 in the LBL cafeteria Peter Armbruster asked Peter Moller: “Do you think the large (negative) shell corrections you obtain in the vicinity of $Z = 108$ and $N = 162$ are related to the just discovered new elements at the GSI”. The calculated shell corrections were subsequently plotted in color where the large shell corrections for DEFORMED nuclei in the previously assumed “sea of instability” clearly stand out. (next 3 frames)

**ATOMIC MASSES AND NUCLEAR GROUND-STATE DEFORMATIONS CALCULATED
WITH A NEW MACROSCOPIC-MICROSCOPIC MODEL**

PETER MÖLLER

Department of Mathematical Physics, University of Lund
S-220 07 Lund 7, Sweden

Theoretical Division, Los Alamos National Laboratory
Los Alamos, New Mexico 87545

and

J. RAYFORD NIX*

Sektion Physik, Universität München
D-8046 Garching, Federal Republic of Germany

Theoretical Division, Los Alamos National Laboratory
Los Alamos, New Mexico 87545

We tabulate the atomic masses and nuclear ground-state deformations of 4023 nuclides ranging from ^{16}O to $^{279}112$, calculated on the basis of a Yukawa-plus-exponential macroscopic model and a folded-Yukawa microscopic model, with new terms included to account for several previously neglected physical effects. With the values of only five constants determined from a least-squares adjustment to ground-state masses, the resulting root-mean-square error in the calculated ground-state masses of 1323 nuclides ranging from ^{16}O to ^{258}No for which experimental values are known with experimental errors less than 1 MeV is 0.835 MeV.

* Alexander von Humboldt Senior U. S. Scientist Awardee

TABLE. Calculated Ground-State Electric Multipole Moments and Masses,
Compared to Experimental Masses Where Available
See p. 171 for Explanation of Table

Z	A	Q ₂ (b)	Q ₄ (b ²)	Calc.mic. (MeV)	Calc.mass (MeV)	Discr. (MeV)	Z	A	Q ₂ (b)	Q ₄ (b ²)	Calc.mic. (MeV)	Calc.mass (MeV)	Discr. (MeV)
108	258	11.6	0.23	-3.11	120.91								
	259	11.6	0.23	-3.56	121.04								
	260	11.7	0.23	-3.69	120.12								
	261	11.3	0.01	-3.87	120.80								
	262	11.3	-0.14	-3.86	120.29								
	263	11.2	-0.28	-4.08	121.18								
	266	10.9	-0.64	-4.60	121.77								
	267	10.9	-0.78	-5.05	122.93								
	268	10.9	-0.93	-5.25	122.98								
	269	10.9	-1.07	-5.85	124.23								
	270	10.9	-1.22	-6.04	124.55								
	271	10.5	-1.28	-6.32	126.36								
	272	10.2	-1.35	-5.75	127.67								
	273	10.2	-1.35	-5.38	130.37								
	274	9.8	-1.28	-4.73	132.00								
	275	9.1	-1.25	-4.36	134.92								
	276	7.7	-0.90	-3.90	136.60								
	277	7.1	-0.72	-3.97	139.31								
	278	5.4	-0.24	-3.78	140.94								
	279	5.0	-0.28	-4.21	143.51								
109	260	11.4	0.15	-3.61	129.13								
	261	11.4	0.01	-3.76	128.07								
	262	11.0	-0.07	-3.96	128.45								
	263	11.0	-0.21	-4.04	127.73								
	269	10.6	-1.01	-5.69	128.99								
	270	10.6	-1.15	-6.32	129.94								
	271	11.0	-1.37	-6.56	130.09								
	272	10.6	-1.44	-6.88	131.59								
	273	10.3	-1.37	-6.30	132.81								
	274	10.3	-1.51	-5.96	135.21								
	275	10.3	-1.38	-5.21	136.83								
	276	9.6	-1.36	-4.91	139.42								
	277	8.5	-1.23	-4.46	140.97								
	278	6.8	-0.77	-4.44	143.51								
	279	5.4	-0.38	-4.30	144.98								
110	263	11.1	-0.07	-3.51	135.72								
	272	10.7	-1.46	-6.32	135.35								
	273	10.3	-1.52	-6.66	136.71								
	274	10.4	-1.53	-6.07	137.67								
	275	10.4	-1.53	-5.71	139.98								
	276	10.1	-1.46	-5.02	141.28								
	277	9.3	-1.43	-4.80	143.68								
	278	7.2	-0.88	-4.49	144.83								
	279	5.8	-0.49	-4.74	146.99								
111	275	10.5	-1.55	-5.64	144.60								
	276	9.8	-1.52	-5.61	146.31								
	277	9.0	-1.35	-5.10	147.32								
	278	8.0	-1.07	-5.06	149.28								
	279	7.3	-0.89	-4.91	150.15								
112	278	8.1	-1.08	-4.93	152.57								
	279	7.7	-0.85	-5.09	154.22								






Fig. 2c. Plot of the ground-state microscopic correction, as calculated in [6] for 4023 nuclei. The fluctuations in the shell correction are larger in the heavier region than in the lighter region.

In the 1985–1992 time frame an improved mass table FRDM1992 encompassing 8979 nuclides was developed (but not in print until 1995, but labeled FRDM1992 because it was finalized and available in the LANL T2 at that time). Next two frames compares odd-even ground-state spins to exp. Third frame compares spins obtained with the first choice of spin-orbit and diffuseness parameters, the new set clearly better. Forth frame shows that remarkably better masses are obtained with the new, current single- particle parameters. The comparison between the impact on masses of two parameter sets is for an interim version of the mass models, the final mass model FRDM2012 agrees even better with experimental masses. For details see AD-NDT 109–110 (2016) and ADNDT 125 (2019).


Calculated ground-state spin compared to experiment




Calculated ground-state spin compared to experiment





Calculated ground-state spin compared to experiment







The next 5 frames compare calculated Q_α with experimental data. The FRDM data were all published/submitted before the experiments so they represent actual predictions. Many features of the data are reproduced by the FRDM, notice in particular the kink in the vicinity of $Z = 108$ and $N = 162$.










Beginning in 1999 computer tech became sufficiently powerful to make practical to calculate fission potential-energy surfaces versus 5 independent shape coordinates for a total of millions of different shapes. The calculations showed results in agreement with the old assumptions that many actinide systems divide into one large spherical fragment and one smaller deformed. In 2011 Randrup et al (PRL 106, 132503 (2011)) showed that remarkably accurate fission yields could be calculate based on random walks on these potential energy surfaces without introducing any adjustable parameters. Next 4 frames. The random walk method was then used to calculate, or predict, where asymmetric fission would occur, 5th frame.

Five Essential Fission Shape Coordinates





45	$Q_2 \sim$ Elongation (fission direction)
⊗	
35	$\alpha_g \sim (M1-M2)/(M1+M2)$ Mass asymmetry
⊗	
15	$\epsilon_{f1} \sim$ Left fragment deformation
⊗	
15	$\epsilon_{f2} \sim$ Right fragment deformation
⊗	
15	$d \sim$ Neck

⇒ 5 315 625 grid points – 306 300 unphysical points

⇒ **5 009 325 physical grid points**

Fission Barrier and Associated Shapes for ^{232}Th





Brownian shape motion

Nuclear deformation energy: $E_{\text{def}}(i,j,k,l,m)$

Bias potential: $V_{\text{bias}}(i) = V_0 (Q_0/Q_2)^2$

Level density parameter: $a_A = A/(8 \text{ MeV})$

Temperature T : $E^* - E_{\text{def}} = a_A T^2$


$$\Rightarrow V(\chi) = E_{\text{def}} + V_{\text{bias}}$$

Metropolis walk:

Change shape: $\chi \rightarrow \chi'$?


$$\begin{cases} V(\chi') < V(\chi): \text{ move with } P = 1 \\ V(\chi') > V(\chi): \text{ move with } P = \exp(-\Delta V/T) \end{cases}$$

Scission: Critical neck radius $c_0 \approx 2.5 \text{ fm}$




P. Möller *et al*, Nature 409 (2001) 785



N. Metropolis *et al*, J Chem Phys 26 (1953) 1087





Fission-Fragment Symmetric-Yield to Peak-Yield Ratio





In 1960 Johansson (Johansson was later professor in Lund and dean of Lund University) suggested (Nuclear Physics 22 (1961) 529) that the asymmetry of actinide fission was due to octupole coupling (splitting) between levels of type $[51\Lambda]$ to levels of type $[40\Lambda]$ at the saddle. He also remarkably predicted that the same couplings could occur at lower levels namely between levels in the rare earth region of type $[41\Lambda]$ to levels of type $[30\Lambda]$ in the rare earth region and might lead to asymmetric fission there. This coupling does occur as predicted. Next two frames see Physics Letters B 789 (2019) 679–684. However, although the saddle shapes are both asymmetric the potential-energy surface for actinides has a persistent asymmetric valley from saddle to scission whereas for Hg the saddle is also asymmetric but beyond the saddle the shape evolves toward symmetry (and splits before reaching symmetry is the interpretation and yield code results)







Contrasting Fission Potential-Energy Surfaces $\text{Hg} \leftrightarrow \text{U}$





The yield models have been developed to describe isotopic yields $Y(Z,N)$ and odd-even staggering. The next 5 frames illustrate the steps towards the current model; the last panel in the last frame illustrating the current results of the yield models, see *Eur. Phys. J. A* (2015) 51: 173 for the whole story.






Next Track-Point Candidates








The next 4 frames illustrate barrier heights and fission half-lives. The first shows that fission half-lives (shorter than 30d) have mainly been observed for barrier heights above 5 MeV. The next two refer to a Polish Woods-Saxon calculation of barrier heights. The 4th frame are barriers calculated in the folded-Yukawa. Frame 1 and 4 are from PRC 91 (2015) 024310. Frame 4 shows that barriers in the r-process path or decay back from the r-process are so low that fission would occur. The two different models in frame 3 and 4 both indicate that barriers are calculated to be too low to allow nuclei with approximately $Z > 120$ and $N > 190$ to exist. It would be desirable to study isotopes with neutron numbers closer to $N = 184$ than currently available to test these theoretical predictions.

Calculated Fission-Barrier Height



Adiabatic fission barriers in superheavy nuclei

P. Jachimowicz

Institute of Physics, University of Zielona Góra, Szafrańska 4a, 65516 Zielona Góra, Poland

M. Kowal* and J. Skalski

National Centre for Nuclear Research, Hoża 69, PL-00-681 Warsaw, Poland

(Received 15 June 2016; published 4 January 2017)

Using the microscopic-macroscopic model based on the deformed Woods–Saxon single-particle potential and the Yukawa-plus-exponential macroscopic energy, we calculated static fission barriers B_f for 1305 heavy and superheavy nuclei $98 \leq Z \leq 126$, including even-even, odd-even, even-odd and odd-odd systems. For odd and odd-odd nuclei, adiabatic potential-energy surfaces were calculated by a minimization over configurations with one blocked neutron or/and proton on a level from the 10th below to the 10th above the Fermi level. The parameters of the model that have been fixed previously by a fit to masses of even-even heavy nuclei were kept unchanged. A search for saddle points has been performed by the “imaginary water flow” method on a basic five-dimensional deformation grid, including triaxiality. Two auxiliary grids were used for checking the effects of the mass asymmetry and hexadecapole nonaxiality. The ground states (g.s.) were found by energy minimization over configurations and deformations. We find that the nonaxiality significantly changes first and second fission saddle in many nuclei. The effect of the mass asymmetry, known to lower the second, very deformed saddles in actinides, in the heaviest nuclei appears at the less deformed saddles in more than 100 nuclei. It happens for those saddles in which the triaxiality does not play any role, which suggests a decoupling between effects of the mass asymmetry and triaxiality. We studied also the influence of the pairing interaction strength on the staggering of B_f for odd- and even-particle numbers. Finally, we provide a comparison of our results with other theoretical fission barrier evaluations and with available experimental estimates.

DOI: [10.1103/PhysRevC.95.014303](https://doi.org/10.1103/PhysRevC.95.014303)**I. INTRODUCTION**

Although fission barrier heights B_f are not directly measurable quantities, i.e., are not quantum observables, they are very useful in estimating nuclear fission rates. As the activation energy E_a (per mole) in chemistry gives a rate k of a chemical reaction at temperature T via the Arrhenius law $k = Ae^{-E_a/RT}$ (where R is the gas constant and A is the frequency factor) [1,2], the fission barrier gives the fission rate Γ_f of an excited (as they usually are in nuclear reactions) nucleus via: $\Gamma_f \sim e^{-B_f/kT_{\text{eff}}}$, where T_{eff} is an effective temperature derived from the excitation energy, and k is the Boltzmann constant. For example, knowing fission barriers of possible fission products helps to predict a cross section for a production of a given evaporation residue in a heavy-ion reaction: one can figure out whether neutron or alpha emission wins a competition with fission at each stage of the deexcitation of a compound nucleus. Moreover, one can try to understand the experimentally established intriguing growth of the total cross sections around $Z = 118$; for its correlation with B_f ; see, e.g., Fig. 6 and the related discussion in Ref. [3]. On the other hand, the prediction of the spontaneous or low-energy (i.e., from a weakly excited state) fission rates, governed by the regime of the collective quantum tunneling, requires an additional knowledge of the barrier shape and mass parameters.

A nonobservable status of the fission barrier, again in analogy to that of the activation energy in chemistry, is

reflected in its possible dependence on a reaction type and/or the excitation energy (effective temperature) range. This leads to some uncertainty in calculations of fission barriers. In particular, it is not clear whether intrinsic configurations should be conserved along the level crossings, which increases B_f , or the adiabatic state should be followed. This is especially relevant for odd- A and odd-odd nuclei, in which sharp crossings of levels occupied by the odd particle exclude the strictly adiabatic scenario. It is known that, if the projection of the single-particle angular momentum Ω on the symmetry axis of a nucleus is conserved, the diabatic effect on the fission barrier can be huge; see, e.g., Ref. [4]. As there is no accepted formula for a barrier correction due to the nonadiabaticity, it is usually ignored, even in odd- N and/or odd- Z nuclei.

A general idea is that, at the excitation energies close to and higher than the barrier but still not inducing sizable dissipative corrections, the adiabatic barrier could be used for calculating fission rates.

Since calculations of potential-energy surfaces (PESs) for odd- A and odd-odd nuclei involve a repetition of calculations for many low-lying quasiparticle states which multiplies the effort (especially in odd-odd systems), systematic studies of their fission barriers are rather scarce. Up to now, they were provided mainly by the Los Alamos microscopic-macroscopic (MM) model and recently by some self-consistent models [5]. The current state of theoretical predictions in fission of even-even nuclei (with $Z \geq 100$) has been discussed recently in Ref. [6].

In the present paper we extend our MM model based on the deformed Woods–Saxon potential, which up to now was applied mainly to even-even nuclei [7], to odd- A and odd-odd

*michal.kowal@ncbj.gov.pl







FIG. 19. Calculated fission-barrier heights B_f for superheavy nuclei.


Calculated Fission-Barrier Heights




The potential-energy model has been developed to allow two asymmetry variables namely fragment Z_1 and N_1 and in the remaining fragment $Z_2=Z-Z_1$ and $N_2=N-N_1$ where Z and N are the fissioning compound system proton and neutron numbers, see Eur. Phys. J. A (2015) 51: 173 for full details. By random walks on the resulting 6D potential surface isotopic yields are obtained, next 3 frames, from Physics Letters B 812 (2021) 136017 show some results. The models have also been extended to calculate fragment neutron emission versus fragment mass, frame 4, details in Phys. Rev. C 103, (2021) 014609.









C O N C L U S I O N S

- All global nuclear-structure models are based on “effective” forces. To expect infinite accuracy with global models is unrealistic.
- Both Wood-Saxon and folded-Yukawa based models give properties of SHE elements to useful accuracy. Both models predicted subsequently observed Q_α values well, in particular the kink near $Z = 108$ and $N = 162$, but show no hope of stability for new elements with $Z > 120$ and no isotopes with $N > 190$.
- Remaining differences between these models and between the models and experiment to a large extent reflect somewhat unavoidable model uncertainties.
- Obviously smaller deviations can be achieved by local adjustments of parameters, but for those of us who strive to improve global model accuracy, this would be a null results.

- Most HFB models have poor (mass) results for known nuclei, therefore their stability predictions in the SHE region are irrelevant. It would be desirable to reduce the number of parameter sets and not use different parameter sets for different purposes, and work to understand how to bring the model results into useful agreement with known data such as nuclear masses.
- For heavy systems it is not the lowest minimum that is the most stable, it is the minimum with the highest fission barrier, a fact overlooked in some calculations.
- Although much used, 2D FISSION potential-energy surfaces obtained in constrained HFB calculations are flawed and do not, and cannot, reflect the properties of the “full” higher-dimensional potential energy function.
- The folded-Yukawa model has been extended to describe many fission properties such as fission (charge, isotopic) yields and neutron-emission versus fragment mass, with encouraging accuracy.

List of publications relevant to Moller presentation 2022-06-15
Three sections: Ground-state properties, beta-decay and astrophysics, fission

GROUND-STATE PROPERTIES

1. CALCULATED GROUND-STATE PROPERTIES OF HEAVY NUCLEI,
P. Möller, S. G. Nilsson and J. R. Nix
Nucl. Phys. A229 (1974) 292
2. NUCLEAR MASS FORMULA WITH A YUKAWA-PLUS-EXPONENTIAL MACROSCOPIC MODEL
AND A FOLDED-YUKAWA SINGLE-PARTICLE POTENTIAL,
P. Möller and J. R. Nix
Nucl. Phys. A361 (1981) 117
3. ATOMIC MASSES AND NUCLEAR GROUND-STATE DEFORMATIONS CALCULATED WITH A
NEW MACROSCOPIC-MICROSCOPIC MODEL,
P. Möller and J. R. Nix
ATOMIC DATA AND NUCLEAR DATA TABLES, 26 (1981) 165-196
4. NUCLEAR SHAPES AND SHAPE TRANSITIONS,
R. Bengtsson, P. Möller, J. R. Nix and J.-y. Zhang
Phys. Scripta 29 (1984) 402
5. ON THE STABILITY OF THE TRANSEINSTEINIUM ELEMENTS
P. Möller, G. A. Leander and J. R. Nix
Z. Phys. A – Atomic Nuclei **323** (1986) 41
6. NUCLEAR MASSES FROM A UNIFIED MACROSCOPIC-MICROSCOPIC MODEL,
P. Möller and J. R. Nix
Los Alamos Preprint LA-UR-3983
ATOMIC DATA AND NUCLEAR DATA TABLES, **39** (1988) 213
(The preprint has more extensive tables than the publication)
7. NUCLEAR MASS FORMULA WITH A FINITE-RANGE DROPLET MODEL AND A FOLDED-
YUKAWA SINGLE-PARTICLE POTENTIAL,
P. Möller, W. D. Myers, W. J. Swiatecki and J. Treiner
Los Alamos Preprint LA-UR-86-4149
Berkeley Preprint LBL-22686
ATOMIC DATA AND NUCLEAR DATA TABLES **39** (1988) 225
(The preprint has more extensive tables than the publication)
8. NUCLEAR PAIRING MODELS,
P. Möller and J. R. Nix
Nucl. Phys. **A536** (1992) 20
9. NUCLEAR GROUND-STATE MASSES AND DEFORMATIONS,
P. Möller, J. R. Nix, W. D. Myers and W. J. Swiatecki
ATOMIC DATA AND NUCLEAR DATA TABLES, **59** (1995) 185–381
10. GLOBAL CALCULATIONS OF GROUND-STATE AXIAL SHAPE-ASYMMETRY OF NUCLEI,
Peter Möller, Ragnar Bengtsson, Peter Olivius, B. Gillis Carlsson, and Takatoshi Ichikawa,
Phys. Rev. Lett. **97** (2006) 162502

11. GLOBAL MICROSCOPIC CALCULATIONS OF GROUND-STATE SPINS AND PARITIES FOR ODD-MASS NUCLEI
L. Bonneau, P. Quentin, and P. Möller,
Phys. Rev. C **76** 024320 (2007)
12. A NON-DISAPPEARING MAGIC TRICK,
Ragnar Bengtsson and Peter Möller,
Nature, **449** (2007) 411–413
13. AXIAL AND REFLECTION ASYMMETRY OF THE NUCLEAR GROUND-STATE Peter Möller,
Ragnar Bengtsson, B. G. Carlsson, P. Olivius, T. Ichikawa, H. Sagawa, A. Iwamoto,
ATOMIC DATA AND NUCLEAR DATA TABLES, **94** (2008) 758–780
14. GLOBAL CALCULATIONS OF NUCLEAR SHAPE COEXISTENCE Peter Möller, Arnold J.
Sierk, Ragnar Bengtsson, Hiroyuki Sagawa, and Takatoshi Ichikawa,
Phys. Rev. Lett. **103** (2009) 212501
15. NEW FINITE-RANGE DROPLET MASS MODEL AND EQUATION-OF-STATE PARAMETERS
P. Möller, W. D. Myers, H. Sagawa, and S. Yoshida,
Phys. Rev. Lett. **108** (2012) 052501 (LA-UR-11-11461)
16. NUCLEAR SHAPE ISOMERS
P. Möller, A. J. Sierk, R. Bengtsson, H. Sagawa, T. Ichikawa,
ATOMIC DATA AND NUCLEAR DATA TABLES, **98** (2012) 149–300
17. NUCLEAR GROUND-STATE MASSES AND DEFORMATIONS: FRDM(2012)
P. Möller, A. J. Sierk, T. Ichikawa, and H. Sagawa,
ATOMIC DATA AND NUCLEAR DATA TABLES 109–110 (2016) 1-204

BETA-DECAY AND ASTROPHYSICS-RELATED PUBLICATIONS

18. CALCULATION OF GAMOW-TELLER β -STRENGTH FUNCTIONS IN THE RUBIDIUM REGION IN THE RPA APPROXIMATION WITH NILSSON MODEL WAVE FUNCTIONS,
J. Krumlinde and P. Möller
Nucl. Phys. A417 (1984) 419
19. ISOTOPIC R-PROCESS ABUNDANCES AND NUCLEAR STRUCTURE FAR FROM STABILITY: IMPLICATIONS FOR THE R-PROCESS MECHANISM,
K.-L. Kratz, J.-P. Bitouzet, F.-K. Thielemann, P. Möller and B. Pfeiffer
ApJ **403** (1993) 216–238
20. NUCLEAR PROPERTIES FOR ASTROPHYSICAL AND RADIOACTIVE ION BEAM APPLICATIONS,
P. Möller, J. R. Nix, and K.-L. Kratz
ATOMIC DATA AND NUCLEAR DATA TABLES, **66** (1997) 131-343
21. RP-PROCESS NUCLEOSYNTHESIS AT EXTREME TEMPERATURE AND DENSITY CONDITIONS, H. Schatz, A. Aprahamian, J. Görres, M. Wiescher, T. Rauscher, J. F. Rembges, F.-K. Thielemann, B. Pfeiffer, P. Möller, K.-L. Kratz, H. Herndl, B. A. Brown and H. Rebel,
Phys. Rep. **294** (1998) 167-263
22. HEATING IN THE ACCRETING NEUTRON STAR OCEAN: IMPLICATIONS FOR SUPERBURST IGNITION
S. Gupta, E. Brown H. Schatz, P. Möller, and K.-L. Kratz,
Astrophysical Journal, **662** (2007) 1188–1197
23. NEUTRON REACTIONS IN ACCRETING NEUTRON STARS: A NEW PATHWAY TO EFFICIENT CRUST HEATING Sanjib S. Gupta, Toshihiko Kawano, and Peter Möller,
Phys. Rev. Lett. **101** (2008) 231101
24. STRONG NEUTRINO COOLING BY CYCLES OF ELECTRON CAPTURE AND β^- DECAY IN NEUTRON STAR CRUSTS,
H. Schatz, S. Gupta, P. Möller, M. Beard, E. F. Brown, A. T. Deibel, L. R. Gasques, W. R. Hix, L. Keek, R. Lau, A. W. Steiner, and M. Wiescher,
Nature **505** (2014) 62
25. A HIGH-ENTROPY-WIND R-PROCESS STUDY BASED ON NUCLEAR-STRUCTURE QUANTITIES FROM THE NEW FINITE-RANGE DROPLET MODEL FRDM(2012)
Karl-Ludwig Kratz, Khalil Farouqi, and Peter Möller,
ApJ **792** (2014) 6
26. THE IMPACT OF INDIVIDUAL NUCLEAR MASSES ON R-PROCESS ABUNDANCES
M.R. Mumpower, R. Surman, D.-L. Fang, M. Beard, P. Möller, T. Kawano, and A. Aprahamian.
Phys. Rev. C **92**, (2015) 035807

27. 94β -DECAY HALF-LIVES OF NEUTRON-RICH $_{55}\text{Cs}$ TO $_{67}\text{Ho}$: EXPERIMENTAL FEEDBACK AND EVALUATION OF THE R-PROCESS RARE-EARTH PEAK FORMATION
J. Wu, S. Nishimura, G. Lorusso, P. Möller, E. Ideguchi, P.-H. Regan, H. Sakurai, 7 G.S. Simpson, P.-A. Söderström, P.M. Walker, H. Watanabe, Z.Y. Xu, Y.L. Ye, H. Baba, F. Browne., 12 R. Daido, P. Doornenbal, Y.F. Fang, G. Gey, T. Isobe, J.J. Liu, C.S. Lee, P.S. Lee, Z. Li, Z. Korkulu, Z. Patel, S. Rice, L. Sinclair, T. Sumikama, M. Tanaka, V. Phong, A. Yagi, R. Yokoyama, G.X Zhang, N. Aoi, T. Alharbi, F.L. Bello Garrote, G. Benzoni, A.M. Bruce, R.J. Carroll, K.Y. Chae, Z. Dombradi, A. Estrade, A. Gottardo, C.J. Griffin, H. Kanaoka, I. Kojouharov, F.G. Kondev, S. Kubono, I. Kuti, N. Kurz, S. Lalkovski, G.J. Lane, E.J. Lee, G. Lotay, C.-B. Moon, H. Nishibata, I. Nishizuka, C.R. Nita, A. Odahara, Zs. Podolyák, O.J. Roberts, C. Shand, H. Schaffner, S. Terashima, J. Taprogge, Z. Vajta, and S. Yoshida,
Phys. Rev. Lett. **118** (2017) 072701
28. NEUTRON- γ COMPETITION FOR β -DELAYED NEUTRON EMISSION
M. R. Mumpower, T. Kawano, and P. Möller
Phys. Rev. C **94** (2016) 064317
29. NUCLEAR PROPERTIES FOR ASTROPHYSICAL AND RADIOACTIVE-ION-BEAM APPLICATIONS (II)
P. Möller, M. R. Mumpower, T. Kawano, W.D. Myers
ATOMIC DATA AND NUCLEAR DATA TABLES **135** (2019) 1–192

FISSION-RELATED PUBLICATIONS

30. CALCULATED FISSION PROPERTIES OF THE HEAVIEST ELEMENTS,
P. Möller, J. R. Nix and W. J. Swiatecki
Nucl. Phys. **A469** (1987) 1
31. NEW DEVELOPMENTS IN THE CALCULATION OF HEAVY-ELEMENT FISSION BARRIERS,
P. Möller, J. R. Nix and W. J. Swiatecki
Nucl. Phys. **A492** (1989) 349-387
32. **TOPICAL REVIEW: STABILITY OF HEAVY AND SUPERHEAVY ELEMENTS**,
P. Möller and J. R. Nix
Journal of Physics G: Nuclear and Particle Physics, TOPICAL REVIEW, **20** (1994) 1681–1747
33. REALISTIC FISSION SADDLE-POINT SHAPES.
P. Möller and A. Iwamoto,
Phys. Rev. **C 61** (2000) 47602
34. TOPOLOGY OF FIVE-DIMENSIONAL, MILLION-GRID-POINT FISSION POTENTIAL-ENERGY SURFACES IN THE 3QS PARAMETERIZATION,
Peter Möller and Akira Iwamoto,
Proc. Conf on Nuclear Shapes and Motions, Santa Fe, New Mexico, Oct. 25–28 (1998)
ACTA PHYSICA HUNGARICA NEW SERIES-HEAVY ION PHYSICS, 10 pp. 241-251 1999
35. NUCLEAR FISSION MODES AND FRAGMENT MASS ASYMMETRIES IN A FIVE-DIMENSIONAL DEFORMATION SPACE, Peter Möller, David G. Madland, Arnold J. Sierk, and Akira Iwamoto,
Nature **409** (2001) 785–790
36. INTO THE FISSION VALLEY,
Peter Möller and Arnold J. Sierk,
Nature, **422** (2003) 485–486
37. FIVE-DIMENSIONAL FISSION-BARRIER CALCULATIONS FROM ^{70}Se TO ^{252}Cf ,
Peter Möller, Arnold J. Sierk, and Akira Iwamoto,
Phys. Rev. Lett. **92** (2004) 072501
38. HEAVY-ELEMENT FISSION-BARRIERS Peter Möller, Arnold J. Sierk, T. Ichikawa, A. Iwamoto, Ragnar Bengtsson, Henrik Uhrenholt, and Sven Åberg,
Phys. Rev. C **79** (2009) 064304 38 pages.
39. ORIGIN OF THE NARROW, SINGLE PEAK IN THE FISSION-FRAGMENT MASS DISTRIBUTION FOR ^{258}Fm , T. Ichikawa, A. Iwamoto, and P. Möller,
Phys. Rev. C **79** (2009) 014305
40. A NEW TYPE OF ASYMMETRIC FISSION IN PROTON-RICH NUCLEI
A.N. Andreyev, J. Elseviers, M. Huyse, P. Van Duppen, S. Antalic, A. Barzakh, N. Bree, T.E. Cocolios, V. F. Comas, J. Diriken, D. Fedorov, V. Fedosseev, S. Franchoo, J.A. Heredia, O. Ivanov, U. Köster, B. A. Marsh, K. Nishio, R.D. Page, N. Patronis, M. Seliverstov, I. Tsekhanovich, P. Van den Bergh, J. Van De Walle, M. Venhart, S. Vermote, M. Veselsky, C. Wagemans, T. Ichikawa, A. Iwamoto, P. Möller, and A.J. Sierk,
Phys. Rev. Lett. **105** (2010) 252502

41. BROWNIAN SHAPE MOTION ON FIVE-DIMENSIONAL POTENTIAL-ENERGY SURFACES:NUCLEAR FISSION-FRAGMENT MASS DISTRIBUTIONS J. Randrup and P. Möller, Phys. Rev. Lett. **106** (2011) 132503106
42. FISSION-FRAGMENT MASS DISTRIBUTIONS FROM STRONGLY DAMPED SHAPE EVOLUTION J. Randrup, P. Möller, and A. J. Sierk Phys. Rev. C **84** (2011) 034613
43. CALCULATED FISSION YIELDS OF NEUTRON-DEFICIENT MERCURY ISOTOPES, P. Möller, J. Randrup, and A. J. Sierk, Phys. Rev. C, **85** (2012) 024306
44. THE CONTRASTING FISSION POTENTIAL-ENERGY STRUCTURE OF ACTINIDES AND MERCURY ISOTOPES, Takatoshi Ichikawa, Akira Iwamoto, Peter Möller, and Arnold J. Sierk, Phys. Rev. C **86** (2012) 024610
45. CHARACTER AND PREVALENCE OF THIRD MINIMA IN ACTINIDE FISSION BARRIERS, Takatoshi Ichikawa, Peter Möller, and A. J. Sierk Phys. Rev. C **87** (2013) 054326,
46. ENERGY DEPENDENCE OF FISSION-FRAGMENT MASS DISTRIBUTIONS FROM STRONGLY DAMPED SHAPE EVOLUTION, Jørgen Randrup and Peter Möller, Phys. Rev. C **88** (2013) 064606,
47. FISSION-FRAGMENT CHARGE YIELDS: VARIATION OF ODD-EVEN STAGGERING WITH ELEMENT NUMBER, ENERGY, AND CHARGE ASYMMETRY, Peter Möller, Jørgen Randrup, Akira Iwamoto, and Takatoshi Ichikawa, Phys. Rev. C **90** (2014) 014601
48. CALCULATED FISSION-FRAGMENT YIELD SYSTEMATICS IN THE REGION $74 \leq Z \leq 94$ AND $90 \leq N \leq 150$, Peter Möller, Jørgen Randrup, Phys. Rev. C **91** (2015) 044316
49. FISSION BARRIERS AT THE END OF THE CHART OF THE NUCLIDES, Peter Möller, Arnold J. Sierk, Takatoshi Ichikawa, Akira Iwamoto, and Matthew Mumpower, Phys. Rev. C **91** (2015) 024310,
50. A METHOD TO CALCULATE FISSION-FRAGMENT YIELDS $Y(Z, N)$ VERSUS PROTON AND NEUTRON NUMBER IN THE BROWNIAN SHAPE-MOTION MODEL; APPLICATION TO CALCULATIONS OF U AND PU CHARGE YIELDS Peter Möller and Takatoshi Ichikawa, European Physics Journal A. **51** (2015) 173

51. EVOLUTION OF URANIUM FISSION-FRAGMENT CHARGE YIELDS WITH NEUTRON NUMBER
Peter Möller and Christelle Schmitt,
European Physics Journal A **53** (2017) 7
52. NUCLEAR SHAPE EVOLUTION BASED ON MICROSCOPIC LEVEL DENSITIES
D. E. Ward, B.G. Carlsson, T. Døssing, and P. Möller, J. Randrup, and S. Åberg
Physical Review C, **95** (2017) 024618
53. THE MICROSCOPIC MECHANISM BEHIND THE FISSION-BARRIER ASYMMETRY (II): THE RARE-EARTH REGION $50 < Z < 82$ AND $82 < N < 126$ T. Ichikawa, P Möller
Phys. Lett. B **789** (2019) 679–684
54. EXCITATION ENERGY PARTITION IN FISSION M. Albertsson, B.G. Carlsson, T. Døssing, P. Möller, J. Randrup, S. Åberg
Phys. Lett. B **803** (2020) 135276
55. CALCULATED FISSION-FRAGMENT MASS YIELDS AND AVERAGE TOTAL KINETIC ENERGIES OF HEAVY AND SUPERHEAVY NUCLEI
M. Albertsson, B.G. Carlsson, T. Døssing, P. Möller, J. Randrup, S. Åberg
Eur. Phys. J. A **56**, (2020) 46
56. ON THE ISOTOPIC COMPOSITION OF FISSION FRAGMENTS
C. Schmitt and P. Möller,
Phys. Lett. B **812** (2021) 136017
57. CORRELATION STUDIES OF FISSION FRAGMENT NEUTRON MULTIPLICITIES
M. Albertsson, B. G. Carlsson, T. Døssing, P. Möller, J. Randrup, S. Åberg
Phys. Rev. C **103**, (2021) 014609