



ESA/Hubble, NASA

# Effects of Neutrino Flavor Conversions in Core-Collapse Supernovae


Multi-Messenger Astrophysics in the Dynamic Universe

4<sup>th</sup> February, 2026

Jakob Ehring (Academia Sinica, Institute of Physics)




# Accretion phase of core-collapse supernovae




(not to scale)

# Neutrino luminosity and mean energy at 400 km




# Neutrino luminosity and mean energy at 400 km





**Production (cooling):**


**Thermal (flavor blind):**



**Charged current (e only):**




## Neutrino luminosity and mean energy at 400 km





**Production (cooling):**

**Thermal (flavor blind):**




**Charged current (e only):**




**Interaction (heating):**

**All flavors: scattering**



**Electron flavor: absorption**



**TL;DR**

**What are neutrino flavor conversions?**

**Propagation**

**Interaction**

$$(\partial_t + \vec{v} \cdot \vec{\nabla}_x) \rho = C$$

# Neutrinos are quantum particles!



Johan Jarnestad/The Royal Swedish Academy of Sciences, 2015

Propagation

Interaction

$$(\partial_t + \vec{v} \cdot \vec{\nabla}_x) \rho = C$$

# Neutrinos are quantum particles!



Johan Jarnestad/The Royal Swedish Academy of Sciences, 2015

Propagation

Density Matrix

Interaction

$$(\partial_t + \vec{v} \cdot \vec{\nabla}_x) \rho = -i[H, \rho] + C$$

Quantum propagation phenomena

The diagram shows the equation  $(\partial_t + \vec{v} \cdot \vec{\nabla}_x) \rho = -i[H, \rho] + C$  enclosed in a blue dashed box. Three arrows point from labels above to parts of the equation: 'Propagation' points to the left-hand side, 'Density Matrix' points to the  $\rho$  term, and 'Interaction' points to the right-hand side. A fourth arrow points from the text 'Quantum propagation phenomena' below to the entire equation.

# Neutrinos are quantum particles!



Johan Jarnestad/The Royal Swedish Academy of Sciences, 2015

Propagation Interaction

Density Matrix

$$(\partial_t + \vec{v} \cdot \vec{\nabla}_x) \rho = -i[H, \rho] + C$$



Quantum propagation phenomena

$$H = H_{\text{vac}} + H_{\text{MSW}} + H_{\nu\nu}$$



For details see Meng-Ru's talk

# A parametric description of flavor conversions in CCSN simulations


# Parametrized recipe:



# Parametrized recipe:




# Parametrized recipe:




$$\nu_e, \bar{\nu}_e \rightarrow \nu_x, \bar{\nu}_x$$

See Ehring, et al. 2023a



# Parametrized recipe:




$$\nu_e, \bar{\nu}_e \rightarrow \nu_x, \bar{\nu}_x$$

At higher energies:

$$\nu_x, \bar{\nu}_x \rightarrow \nu_e, \bar{\nu}_e$$


See Ehring, et al. 2023a




# Probe effect in different regions:

Use density to restrict flavor conversions to different regions:

- Inside Heating Region ( $\rho_{\text{crit}} = 10^{10} \text{ g/cm}^3$ ) (A)
- Inside Cooling Region ( $\rho_{\text{crit}} = 10^{12} \text{ g/cm}^3$ ) (B)




# Results from simulations in axial symmetry (2D)




**(A): Inside heating region**

# Local heating




From Ehring, et al. 2023a


# Local heating



From Ehring, et al. 2023a


# Local heating







**Low mass progenitor system  
Conversions in heating region**

- **Earlier onset of explosion**
- **Increased specific heating**




**(B): Inside heating and cooling region**





**Heavy progenitor system**  
**Conversions in heating and cooling**  
**region**



- **Tendency to black hole formation is increased in all models with flavour conversion**
- **Additionally increased cooling leads to faster PNS contraction**





**From Ehring, et al. 2023b**





$9.0M_{\odot}$     $11.2M_{\odot}$     $12.00M_{\odot}$     $13.00M_{\odot}$     $14.00M_{\odot}$     $15.00M_{\odot}$     $16.00M_{\odot}$   
 $17.00M_{\odot}$     $18.00M_{\odot}$     $19.00M_{\odot}$     $20M_{\odot}$     $21.00M_{\odot}$     $23.00M_{\odot}$




$12.00 M_{\odot}$   $13.00 M_{\odot}$   $14.00 M_{\odot}$   $15.00 M_{\odot}$   $16.00 M_{\odot}$   
 $17.00 M_{\odot}$   $18.00 M_{\odot}$   $19.00 M_{\odot}$   $20 M_{\odot}$   $21.00 M_{\odot}$   $23.00 M_{\odot}$




$18.00 M_{\odot}$   $19.00 M_{\odot}$   $21.00 M_{\odot}$   $16.00 M_{\odot}$



$21.00 M_{\odot}$





- low mass ( $m \lesssim 12 M_{\odot}$ ): more vigorous explosions
- intermediate mass ( $12 M_{\odot} \lesssim m \lesssim 15 M_{\odot}$ ): faster (?) BH formation
- intermediate mass ( $16 M_{\odot} \lesssim m \lesssim 19 M_{\odot}$ ): increased explodability
- higher masses ( $m \gtrsim 20 M_{\odot}$ ): prevents (?) explosions

# Summary

## ➔ Flavor conversions influence the dynamics

➔ They can help AND hinder AND enable AND prevent explosions

➔ Flavor conversions enhance

- Heating in gain region
- Cooling in cooling region and PNS mantle

➔ Temporal dependence on the effects


- Soon after bounce: largest effect on heating
- At later times: cooling effect remains

Visit us in Taiwan for a workshop on Neutrino Flavor Conversions  
in CCSN and BNSM

23.03.2026 - 27.03.2026





# Backup




**From Ehring, et al. 2023b**


# 11.2 M<sub>⊙</sub> - 3D



From Mori and Mori 2025



From Wang and Burrows 2025a



From Akako et al 2026

