



Challenge to GW170817 with Numerical Relativity



Max-Planck-Institut
für Gravitationsphysik
(Albert-Einstein-Institut)

Kenta Kiuchi (CRA/YITP)



Center for Gravitational Physics and
Quantum Information

Yukawa Institute for Theoretical Physics, Kyoto University

Happy 60th birthday, Shibata san!



@ banquet last week

~~Japanese quates: 名選手、名監督にあらず~~

~~⇒ An excellent player is not necessarily an excellent director (supervisor)~~


Shibata san's contribution to relativistic astrophysics

Timetable of the 2nd week (February 2 - February 6)

February 2 (Mon.)	February 3 (Tue.)	February 4 (Wed.)	February 5 (Thu.)	February 6 (Fri.)
10:00-10:55 Registration	10:00-11:00 A. Tanikawa	9:30-10:20 D. Radice	9:30-10:20 M. Aloy	9:30-10:20 O. Just
		10:20-11:10 E. Most	10:20-11:10 P. Mösta	10:20-11:10 S. Fujibayashi
10:55-11:00 Opening	11:00-11:30 Coffee	11:10-11:30 Coffee	11:10-11:30 Coffee	11:10-11:30 Coffee
11:00-12:30 T. Piran	11:30-12:30 Discussion (Tanikawa, Morisaki)	11:30-12:30 M. Miravet-Tenes , E. Gutierrez, A. Reboul-Salze	11:30-12:30 T. Kuroda , C. Musolino , A. Mizuta	11:30-12:30 A. Janiuk, D. Desai, A.T.L. Lam
12:30-14:00 Lunch & Poster	12:30-14:00 Lunch & Poster	12:30-14:00 Photos Lunch & Poster	12:30-14:00 Lunch & Poster	12:30-14:00 Lunch & Poster
14:00-15:00 A. Perego	14:00-15:00 M.-R. Wu	14:00-14:50 K. Kawaguchi	14:00-14:50 T. Dietrich	14:00-15:40 A. Agarwal , A. Shankar, P. Hammond, Y. Gao , M. Jacobi
15:00-15:30 Coffee	15:00-15:30 Coffee	14:50-15:50 J. Ehring , L. Longo Micchi , P. Cheong	14:50-15:50 Y. Qiu , H.-J. Kuan, K. Hayashi	15:40-16:10 Coffee
15:30-17:00 Discussion (Perego, Wanajo)	15:30-17:00 Discussion (Wu, Nagakura)	15:50-16:20 Coffee	15:50-16:20 Coffee	16:10- M. Shibata
		16:20-17:50 Poster & Free Discussion	16:20-17:50 Discussion (Radice, Dietrich)	


Please upload the talk slide, Shibata san!

Shibata san's contribution to relativistic astrophysics



These lists are the tip of the iceberg of Shibata san's pioneering works.


Shibata san's contribution to relativistic astrophysics



GW170817 + GRB170817A + AT2017gfo

I appreciate Shibata-san's tremendous contributions to relativistic astrophysics, particularly in binary neutron star mergers.


Reality



Introduction



Imagination



There is no self-consistent model to explain it.

We as numerical relativists must explain

► SGRB (+Afterglow observations)

► Kilonova emission: $M_{\text{eje}} \approx 0.05 M_{\odot}$ (Hotokezaka & Nakar20), Red+Blue? (Kitamura et al. 25)

at least qualitatively, and quantitatively at the best.

A “package” for EM counterpart modeling

NR simulation

(GR+EOS+ ν -Rad.+MHD)






R-process nucleosynthesis calculation



Kilonova/GRB light curve modeling



Observation



Sys. Err.

- ▶ Resolution
- ▶ Approx. GR
- ▶ Approx. ν -Rad
- ▶ MHD approx.


...

- ▶ Reaction rate
- ▶ Mass model

...


- ▶ Photon rad. transfer

...



Neutron rich matter ejection mechanism

S. Fujibayashi, KK, et al. 18



Many Refs.

Hotokezaka et al. 13,
Fujibayashi et al. 18,20,21,22

Bauswein et al. 13, 17

Just et al. 14, 21

Siegel & Metzger, 18

Combi & Siegel, 23

Radice et al. 18

+ more



► One missing mass ejection channel = **Lorentz force-driven ejecta**

Recent progress indicates its relevance. (Mösta et al. 20, Combi & Siegel 23, KK et al. 24, Most 23)

To B or not to B in binary neutron star merger (by Victoria M. Kaspi)

$P - \dot{P}$ Diagram

Image of the binary pulsar



- Assumption : Rotational energy is dissipated by the magnetic dipole radiation $\Rightarrow B \propto (P\dot{P})^{1/2}$

To B or not to B in binary neutron star merger (by Victoria M. Kaspi)

► B-field in observed binary NSs : $10^{9.7} - 10^{12.2}$ G

Kinetic energy before the merger $\sim 10^{53}$ g cm² s⁻² $(M/2.7M_{\text{sun}})(v/0.3c)^2$

B-field energy $\sim 10^{41}$ g cm² s⁻² $(B/10^{12}\text{G})^2(R/10^6\text{cm})^3$

B-field is irrelevant in BNS mergers ?

No ! \Rightarrow Several amplification mechanisms (Magneto Hydro Dynamical instabilities) could amplify the B-field up to the dynamically important level

Lorentz force (MHD)-driven ejecta as a “new” channel
 Question: How can be a **strong and large-scale field** established?

Mean field dynamo theory

$$\partial_t \bar{\mathbf{B}} = \nabla \times (\bar{\mathbf{U}} \times \bar{\mathbf{B}} + \bar{\mathcal{E}}), \quad \mathbf{Q} = \bar{\mathbf{Q}} + \mathbf{q}, \quad \bar{\mathbf{Q}} = \text{Axisym. Ave.}$$

$$\bar{\mathcal{E}} = \overline{\mathbf{u} \times \mathbf{b}} \quad \mathbf{u} \ \& \ \mathbf{b} : \text{turbulence of the velocity and b-field.}$$


α Ω dynamo

$$\bar{\mathcal{E}}_i = \alpha_{ij} \bar{B}_j + \beta_{ij} \overline{(\nabla \times \mathbf{B})_j} \approx \alpha_{ii} \bar{B}_i$$

$$\partial_t \bar{B}_\varphi = R \bar{B}^A \nabla_A \Omega \quad (A = R, z, \ \Omega - \text{effect})$$

$$\partial_t \bar{B}_R = \partial_z \mathcal{E}_\varphi \approx \partial_z (\alpha_{\varphi\varphi} \bar{B}_\varphi) \quad (\alpha - \text{effect})$$

$$\partial_t \bar{B}_z = -\partial_R \mathcal{E}_\varphi \approx \partial_R (\alpha_{\varphi\varphi} \bar{B}_\varphi)$$



Generation of a large-scale field via α Ω dynamo

α Ω dynamo theory prediction (Check list)




1. \bar{B}_φ should be anticorrelated with $\bar{B}_{R/z}$.
2. \bar{E}_φ should be correlated or anti-correlated with \bar{B}_φ .
3. Dynamo cycle period $P_{\text{theory}} = 2\pi (\alpha_{\phi\phi} d\Omega/d\ln R k_z/2)^{-1/2}$
4. Dynamo wave propagation direction according to the Yoshimura-Parker rule $\alpha_{\phi\phi} \nabla \Omega \times \mathbf{e}_\phi$

Question: What generate electromotive force (EMF), i.e., fluctuation component of velocity and magnetic field?

$$\bar{\mathcal{E}} = \overline{\mathbf{u} \times \mathbf{b}}$$

Generation of a large-scale field via $\alpha \Omega$ dynamo

Kelvin Helmholtz instability (Rasio and Shapiro 99, Price & Rosswog 05)



$$\sigma \propto k$$

High grid resolution is key.


(KK et al. 14,15,18,24, Palenzuela et al. 22, Aguilera-Miret et al. 20, 22, 23)

Generation of a large-scale field via $\alpha \Omega$ dynamo

Magneto Rotational Instability (MRI) (Balbus & Hawley 91)

► Differential rotation: $\nabla\Omega < 0$, $\lambda_{MRI}^{RNS} \approx 80m \left(\frac{B_p}{10^{15}G}\right) \left(\frac{\rho}{10^{15}g\text{ cm}^{-3}}\right)^{-\frac{1}{2}} \left(\frac{\Omega}{8000\text{ rads}^{-1}}\right)^{-1}$

$$\lambda_{MRI}^{BH-Disk} \approx 1,000m \left(\frac{B_p}{10^{15}G}\right) \left(\frac{\rho}{10^{13}g\text{ cm}^{-3}}\right)^{-\frac{1}{2}} \left(\frac{\Omega}{6000\text{ rads}^{-1}}\right)^{-1}$$




Again, high resolution is key. (Shibata et al. 05, Duez et al. 05, Siegel et al. 13, KK et al. 18,24)

Methodology

- ▶ Einstein's solver (Shibata & Nakamura 95, Baumgarte & Shapiro 98, Barker et al, 06, Campanelli et al. 06, Hilditch et al. 13)
- ▶ Nuclear theory-based equation of state for the NS matter (SFHo/BHBLp/DD2) (Steiner et al. 13, Banik et al. 14)
- ▶ Relativistic magnetohydrodynamics solver (KK et al. 22, Migone et al. 09, Gardiner & Stone 08)
- ▶ Neutrino-radiation transfer solver (Sekiguchi et al. 12)

+ for more technical issues (see KK et al. 22)



All the works, we quantify the ability of our simulation set up to resolve the KHI and MRI:

$$Q_{MRI} \equiv \frac{\lambda_{MRI}}{\Delta x} \geq 10$$

Code Name

- ▶ I was asked more than 100 times about which NR code we use.
- ▶ The code was originally developed by Shibata san, and my colleagues implemented new features, e.g., neutrino, MHD, etc.
- ▶ It doesn't have a specific name.

Numerical relativity code for **A**strophysical and **N**on-**A**strophys**I**cal objects = No name in Japanese

Computational facilities



Fugaku@Riken (Japan)



Raven@MPCDF (Germany)

Supercomputer = Experimental labo.

Inferring the EOS from ab initio simulations





Lifetime of the remnant massive neutron star

Prompt
~ 0s

Short-lived
~ O(0.01) s

Intermediate-lived
~ O(0.1)s

Long-lived
~ O(1) s



$L_{iso} \sim 10^{49}$ erg/s
Hayashi et al. PRL 25

No jet until 1s at least.
KK PRL 23

$L_{iso} \sim 10^{52}$ erg/s
KK 26 in prep.

$L_{iso} \sim 10^{52}$ erg/s
KK Nature Astro. 24

EOS stiffness: Soft (SFHo)
Binary mass: Large



Stiff (DD2)
Small

Long-lived remnant formation (KK et al. Nature Astron. 24)

0.2s simulation, DD2-1.35-1.35M_⊙, $\Delta x_{\text{finiest}}=12.5$ m, $B_{\text{max}}=10^{15.5}$ G, 36M CH


Y_e

ρ ($g\ cm^{-3}$) + B - field line



B (G)

©K. Hayashi

2,000 km




Time: 0.02 ms




Kelvin-Helmholtz dynamo at the merger

KH amplification at the merger




Growth rate vs initial B-field



► In reality, the KH dynamo would produce a **strong**, but **small-scale** magneto turbulence (see also Palenzuela et al. 22, Aguilera-Miret et al. 22, 23).

Neutrino viscosity/drag on MRI



Neutrino viscosity/drag could be irrelevant for MRI.

Dispersion relation for MRI with the neutrino viscosity:

$$\left[\left(\tilde{\sigma} + \tilde{k}^2 \tilde{\nu} \right) \tilde{\sigma} + \tilde{k}^2 \right]^2 + \tilde{\kappa}^2 \left[\tilde{\sigma}^2 + \tilde{k}^2 \right] - 4\tilde{k}^2 = 0,$$

$$\tilde{\sigma} \equiv \frac{\sigma}{\Omega}, \quad \tilde{k} \equiv \frac{k v_A}{\Omega}, \quad \tilde{\kappa}^2 \equiv \frac{\kappa^2}{\Omega^2}, \quad \tilde{\nu} \equiv \frac{\nu \Omega}{v_A^2}$$

Dispersion relation for the neutrino drag:

$$\left[\left(\tilde{\sigma} + \tilde{\Gamma} \right) \tilde{\sigma} + \tilde{k}^2 \right]^2 + \tilde{\kappa}^2 \left[\tilde{\sigma}^2 + \tilde{k}^2 \right] - 4\tilde{k}^2 = 0,$$

$$\tilde{\Gamma} \equiv \frac{\Gamma}{\Omega}$$


$$\nu = 1.2 \times 10^{10} (\rho/10^{13} \text{gcm}^{-3})^{-2} (T/10 \text{MeV})^2 \text{cm}^2 \text{s}^{-1}$$

$$\Gamma = 6 \times 10^3 (T/10 \text{MeV})^6 \text{s}^{-1} \quad \text{Guilet et al. 16}$$

Check list to pin down an $\alpha \Omega$ dynamo


Prerequisite

$$\text{MRI quality factor: } \frac{\lambda_{MRI}}{\Delta x} > 10$$



MRI is well resolved in $\Delta x=12.5\text{m}$ run \Rightarrow Turbulence is developed
MRI is not resolved in $\Delta x=200\text{m}$ run \Rightarrow No turbulence

Generation of a large-scale field via $\alpha \Omega$ dynamo




- ▶ Waves generated in the MRI-active region propagates towards the polar
- ▶ The B-field deep inside the core in the polar region stays buried throughout the simulation

Check list to pin down an $\alpha \Omega$ dynamo


Ω effect

\bar{B}_ϕ (G)




α effect

\bar{B}_ϕ (G)




$t - t_{\text{merger}}$ (ms)

\bar{B}_R (G)



$t - t_{\text{merger}}$ (ms)

\mathcal{E}_ϕ (G cm s^{-1})



$t - t_{\text{merger}}$ (ms)

$t - t_{\text{merger}}$ (ms)

Check list to pin down an $\alpha \Omega$ dynamo

Pearson correlation between \bar{E}_ϕ and \bar{B}_ϕ





Table 1 The $\alpha\Omega$ dynamo period prediction and simulation data at several radii



R (km)	$\alpha_{\phi\phi}$ (cm/s)	Ω (rad/s)	Shear rate	k_z (/cm)	P_{theory} (s)	P_{sim} (s)
20	-8.1×10^6	4025	$q = -1.0$	6.3×10^{-6}	<u>0.020</u>	<u>0.018</u>
30	-1.0×10^7	2515	$q = -1.34$	4.2×10^{-6}	<u>0.021</u>	<u>0.018–0.024</u>
40	-1.0×10^7	1688	$q = -1.44$	3.3×10^{-6}	<u>0.037</u>	<u>0.018–0.030</u>
50	-4.4×10^6	1200	$q = -1.50$	2.6×10^{-6}	<u>0.062</u>	<u>0.030–0.040</u>

Check list to pin down an $\alpha \Omega$ dynamo



Dynamo wave propagates to the z direction according to the Yoshimura-Parker rule $\alpha_\phi \nabla \Omega \times e_\phi$


Jet from long-lived remnant formation




- ▶ Poynting flux dominated luminosity outflow is $L_{\text{jet}}^{\text{iso}} \approx 10^{52} \text{ erg/s}$
- ▶ Relativistic outflow is confined in a region with $\theta \sim 12^\circ$.
- ▶ Terminal Lorentz factor ≈ 10 -20.
- ▶ The standard resolution (200m) underestimates the luminosity by a factor of 10-100.

Mass ejection from long-lived remnant formation


Ejecta mass evolution



Mass histogram vs Y_e



Mass histogram vs v_{∞}



- ▶ $M_{\text{eje, peak, dyn}} \approx 10^{-3} M_{\odot}$, $M_{\text{eje, peak, post}}$ (Lorentz force-driven) $\approx 0.1 M_{\odot}$,
- ▶ $Y_{e, \text{peak, dyn}} \approx 0.24$, $Y_{e, \text{peak, post}} \approx 0.22$,
- ▶ $v_{\infty, \text{peak, dyn}} \approx 0.1-0.3 c$, $v_{\infty, \text{peak, post}} \approx 0.1c$
- ▶ The standard resolution (200m) underestimates the ejecta mass by a factor of 10 (see also Mösta et al. 20).

Inferring the EOS from ab initio simulations





Lifetime of the remnant massive neutron star

Prompt
~ 0s

Short-lived
~O(0.01) s

Intermediate-lived
~O(0.1)s

Long-lived
~O(1) s



$L_{iso} \sim 10^{49}$ erg/s
Hayashi et al. PRL 25

No jet until 1s at least.
KK PRL 23

$L_{iso} \sim 10^{52}$ erg/s
KK 26 in prep.

$L_{iso} \sim 10^{52}$ erg/s
KK Nature Astro. 24

EOS stiffness: Soft (SFHo)
Binary mass: Large




Stiff (DD2)
Small

Prompt BH formation motivated by GW190425 (Hayashi, KK et al. 24)

1.5s simulation, SFHo-1.25-1.65M_⊙, $\Delta x_{\text{finiest}}=150\text{m}$, $B_{\text{max}}=10^{15}\text{G}$, 130M CH
 ρ (g cm⁻³) + B - field line B (G) Y_e



2,000 km

100 km



©K. Hayashi

Prompt BH formation




► Butterfly diagram suggests MRI-driven $\alpha \Omega$ dynamo: $P_{\alpha \Omega} = 0.03s$ vs $P_{BF} = 0.03-0.04s$

► Generation of large-scale B-field \Rightarrow Blandford-Znajek mechanism.


$$L_{jet} \sim 10^{49} \text{ erg/s}$$

Mass ejection from prompt BH formation


Ejecta mass evolution



Mass histogram vs Y_e



Mass histogram vs v_∞



► Dynamical ejecta $\approx 1.6 \times 10^{-3} M_\odot$, Post-merger ejecta (MRI-driven turbulent viscosity & Lorentz force) $\approx 4.7 \times 10^{-3} M_\odot$

► $Y_{e, \text{peak, dyn}} \approx 0.08$, $Y_{e, \text{peak, post}} \approx 0.28$

► $v_{\infty, \text{peak, dyn}} \approx 0.2c$, $v_{\infty, \text{peak, post}} \approx 0.08c$

Inferring the EOS from ab initio simulations





Lifetime of the remnant massive neutron star

Prompt
~ 0s

Short-lived
~0(0.01) s

Intermediate-lived
~0(0.1)s

Long-lived
~0(1) s



$L_{iso} \sim 10^{49}$ erg/s
Hayashi et al. PRL 25

No jet until 1s at least.
KK PRL 23

$L_{iso} \sim 10^{52}$ erg/s
KK 26 in prep.

$L_{iso} \sim 10^{52}$ erg/s
KK Nature Astro. 24

EOS stiffness: Soft (SFHo)
Binary mass: Large



Stiff (DD2)
Small

Short-lived remnant formation (KK et al. PRL 23)


1.2s simulation, SFHo-1.2-1.5M_⊙, $\Delta x_{\text{finiest}}=150\text{m}\&200\text{m}$, 85M CH

ρ (g cm^{-3})

Y_e

$B(G)$


©K. Hayashi




(see also, Just et al. 14, 21)

No “jet” from the short-lived remnant formation


No jet until 1s at least.



Butter-fly diagram




Ram-pressure/Mag-pressure




- ▶ BF diagram indicates the $\alpha \Omega$ dynamo generates the large-scale B-field.
- ▶ Resultant large-scale is not strong enough to overcome the ram-pressure. Why? Disk rotational energy is $\approx 10^{51}$ erg, c.f. Remnant NS rotational energy is $\approx 10^{53}$ erg.

Mass ejection from the short-lived remnant formation


Ejecta mass evolution



Mass histogram vs Y_e



Mass histogram vs v_∞



▶ $M_{\text{eje, peak, dyn}} \approx 6 \times 10^{-3} M_\odot$, $M_{\text{eje, peak, post}} \approx 8 \times 10^{-3} M_\odot$ (MRI-driven turbulent viscosity)

▶ $Y_{\text{e, peak, dyn}} \approx 0.03$, $Y_{\text{e, peak, post}} \approx 0.26-0.27$.

▶ $v_{\infty, \text{ peak, dyn}} \approx 0.2-0.3c$, $v_{\infty, \text{ peak, post}} \approx 0.08-0.10c$.

Inferring the EOS from ab initio simulations





Lifetime of the remnant massive neutron star

Prompt
~ 0s

Short-lived
~ 0(0.01) s

Intermediate-lived
~ 0(0.1) s

Long-lived
~ 0(1) s



$L_{iso} \sim 10^{49}$ erg/s
Hayashi et al. PRL 25

No jet until 1s at least.
KK PRL 23

$L_{iso} \sim 10^{52}$ erg/s
KK 26 in prep.

$L_{iso} \sim 10^{52}$ erg/s
KK Nature Astro. 24

EOS stiffness: Soft (SFHo)
Binary mass: Large



Stiff (DD2)
Small

Intermediate lived remnant formation case (KK in prep)


$M_{\text{chirp}} = 1.186 M_{\odot}$, BHB Λ_{ϕ} , $\Delta x_{\text{finest}} = 12.5 \text{ m}$, 0.7 second, $B_{\text{max}} = 10^{15} \text{ G}$, 150M CH

$\rho \text{ (g cm}^{-3}\text{)}$

$B \text{ (G)}$


Y_e

©K. Hayashi



Pinning down the $\alpha\Omega$ dynamo in BNS

Butterfly diagram



Theory vs simulation

R [km]	$\alpha_{\phi\phi}$ [cm/s]	k_z [cm $^{-1}$]	$d\Omega/d \ln R$ [s $^{-1}$]	$P_{\alpha\Omega}$ [ms]	P_{BF} [ms]
10	8.0×10^6	2.4×10^{-5}	5359	<u>8.8</u>	<u>12</u>
20	$0.5 - 1.0 \times 10^7$	7.0×10^{-6}	4955	<u>15 - 25</u>	<u>17</u>
30	1.5×10^7	3.3×10^{-6}	3505	<u>21</u>	<u>17</u>

- ▶ A couple of predictions based on the mean-field large-scale dynamo theory (not shown)
- ▶ The most relevant prediction is


$$P_{\alpha\Omega} = 2 \pi (\alpha_{\phi\phi} d\Omega/d \ln R k_z)^{-1/2}$$

$$\partial_t \bar{B}_\varphi = R \bar{B}^R \nabla_R \Omega,$$


$$\partial_t \bar{B}_R = \partial_z \bar{\mathcal{E}}_\varphi \approx \partial_z (\alpha_{\phi\phi} \bar{B}_\varphi)$$

“Jet” from the intermediate-lived remnant formation

Poynting flux



Required jet kinetic energy (GW170817)




► $L_{\text{jet}}^{\text{iso}} \sim 10^{52} \text{ erg/s} \Rightarrow$ If 1 second duration and 10% convergence efficiency are assumed, it is consistent with the required jet kinetic energy in GW170817 (see also Sadeh & Waxman 24).


► $\Gamma_{\infty} \approx 30$

Intermediate lived remnant formation case

Ejecta mass evolution



Mass histogram vs Y_e




► $M_{\text{eje, dyn}} \approx 2 \times 10^{-3} M_{\odot}$, $M_{\text{eje, post}}$ (Lorentz-force-driven) $\approx 8 \times 10^{-2} M_{\odot}$.

► $Y_{\text{e, peak, dyn}} \approx 0.03$, $Y_{\text{e, peak, post}} \approx 0.2$

Intermediate lived remnant formation case

GW signals



The energy emitted in the post-merger phase is below the observed upper limit (LVK18).

Inferring the EOS from “jet” launching





Lifetime of the remnant massive neutron star

Prompt
~ 0s

Short-lived
~0(0.01) s

Intermediate-lived
~0(0.1)s

Long-lived
~0(1) s



$L_{iso} \sim 10^{49}$ erg/s

No jet until 1s at least.

$L_{iso} \sim 10^{52}$ erg/s

$L_{iso} \sim 10^{52}$ erg/s

SFHo

BHB Λ_ϕ

DD2

Inferring the EOS from R-process nucleosynthesis





Lifetime of the remnant massive neutron star

Prompt
~ 0s

Short-lived
~0(0.01) s

Intermediate-lived
~0(0.1)s

Long-lived
~0(1) s



$$M_{eje} \approx 0.006 M_{\odot}$$

$$M_{eje} \approx 0.014 M_{\odot}$$

$$M_{eje} \approx 0.08 M_{\odot}$$

$$M_{eje} \approx 0.1 M_{\odot}$$

$$M_{eje}^{GW170817} \approx 0.05 M_{\odot}$$

R-process universality:

SFHo

BHB \wedge ϕ

DD2



Kilonova in GW170817:

SFHo

BHB \wedge ϕ

DD2


End-to-End modeling of GW170817



Nuclear EOS predicts a “blind” spot on this diagram
Mass asymmetry? Jet-ejecta interaction? Different BNS merger types?

A prelude to magnetar formation in BNS

1.35-1.35 M_{\odot} , DD2, $\Delta x=6.25\text{m}$, $B_{0,\text{max}} = 3 \times 10^{12}\text{ G}$, 480M CH



At least, the EM energy reaches an expected saturation level (see also Gutiérrez et al. 26).

Another potential large-scale dynamo



$$\partial_t \bar{\mathbf{B}} = \nabla \times (\bar{\mathbf{U}} \times \bar{\mathbf{B}} + \bar{\mathcal{E}}),$$

$$\bar{\mathcal{E}} = \overline{\mathbf{u} \times \mathbf{b}}$$



Taylor-Spruit dynamo loop

MRI-inactive

MRI-active



Spruit 98





Saturated turbulent magnetic field

Barrère et al. 25

- ▶ Deep inside (Outside) core is MRI-inactive (active) region
- ▶ Bulk EM energy is contained in the MRI-inactive region.

Taylor-Spruit dynamo could be the case in the core.

Taylor-Spruit dynamo in BNS (Reboul-Salze et al. 25)



► Solving the linear perturbation equation on top of the simulation data.
(gray: stable, white unstable)


► Numerical viscosity alters the critical strength of the instability
Taylor-Spruit dynamo is the next challenge.

Conclusion

- ▶ A self-consistent direct modelling of BNS merger is feasible.
- ▶ For the long-lived case, $L_{jet}^{iso} \sim 10^{52} \text{ erg/s}$, $M_{eje} \approx 0.1 M_{\odot} \gg M_{eje}^{GW170817}$, and the solar R-process can not be reproduced.
- ▶ The intermediate case, $L_{jet}^{iso} \sim 10^{52} \text{ erg/s}$, $M_{eje} \approx 0.08 M_{\odot} \sim M_{eje}^{GW170817}$.
- ▶ For the short-lived case, no strong jet, $M_{eje} \approx 0.014 M_{\odot} \ll M_{eje}^{GW170817}$, and the solar R-process is reproduced.
- ▶ For the prompt collapse case, $L_{eje}^{iso} \sim 10^{49} \text{ erg/s}$, $M_{eje} \approx 0.006 M_{\odot}$.

Caveat: A large systematics in hydro. simulation, in particular, MHD-turbulent case.

Mean B-field in MRI-active region




- ▶ Deep inside (Outside) core is MRI-inactive (active) region
- ▶ Bulk EM energy is contained in the MRI-inactive region.

Prompt BH formation

Rest-mass density

MRI-quality factor

Mean poloidal B-field



Once the MRI starts to be resolved, the mean poloidal B-field is generated.

$P_{\text{theory}} = 0.03$ s vs $P_{\text{BF}} = 0.03-0.04$ s

Electromagnetic emission in compact binary mergers

R(paid)-process nucleosynthesis and EM

(Lattimer & Schramm 74, Metzger et al. 10, Li & Paczynski 98)

Role of the r-process elements

- ▶ Heating source via radio-active decay (Kasen et al. 17)

$$\dot{\epsilon} \approx 10^{10} \text{ erg s}^{-1} \text{ g}^{-1} \left(\frac{t}{\text{day}} \right)^{-1.3}$$

- ▶ Opacity source (Lanthanide elements)

(Barnes & Kasen 13, Tanaka & Hotokezaka 13)

$$\kappa \approx 10 \text{ cm}^2 \text{ g}^{-1}$$


Properties of electromagnetic emission (Optical-IR)

- ▶ Peak time (diffusion time = dynamical time)

$$t_{\text{peak}} \approx 5.7 \text{ day} \left(\frac{\kappa}{10 \text{ cm}^{-2} \text{ g}^{-1}} \right)^{1/2} \left(\frac{M_{\text{eje}}}{0.03 M_{\odot}} \right)^{1/2} \left(\frac{v_{\text{ej}}}{0.2c} \right)^{-1/2}$$

- ▶ Peak Luminosity



$$L \approx \dot{\epsilon} M_{\text{ej}} \approx 6 \times 10^{41} \text{ erg s}^{-1} \left(\frac{M_{\text{eje}}}{0.03 M_{\odot}} \right) \left(\frac{t}{\text{day}} \right)^{-1.3}$$



Slide courtesy of M. Tanaka

R-process nucleosynthesis and its opacity



Tanaka et al. 17



- ▶ Electron fraction Y_e (# of electron/# of baryon) is a key quantity
- ▶ $Y_e \gtrsim 0.25$ produces negligible / small amount of lanthanide \Rightarrow low opacity in optical
- ▶ $Y_e \lesssim 0.25$ produces lanthanide \Rightarrow high opacity in IR
- ▶ Neutrino reaction determines Y_e of the ejecta

Super long-term BNS inspiral simulation (KK 25)

Residual phase error



- ▶ Fourth-order accurate finite volume Riemann solver (KK 25)
- ▶ $\delta\phi_{\text{error}} = 0.27 \pm 0.07 \text{ rad}$ (*new solver*) vs $\delta\phi_{\text{error}} = 0.58 \pm 0.22 \text{ rad}$ (*old solver*)
- ▶ ≈ 100 GW cycles

Black Hole - Neutron Star mergers

Key ingredients for tidal disruption in BH-NS

Tidal force $>$ NS self gravity $\Rightarrow r \lesssim (M_{\text{BH}}/M_{\text{NS}})^{-2/3} (M_{\text{NS}}/R_{\text{NS}})^{-1} M_{\text{BH}} \equiv r_{\text{tidal}}$

If $r_{\text{tidal}} > r_{\text{isco}} \Rightarrow$ Tidal disruption

$r_{\text{tidal}} < r_{\text{isco}} \Rightarrow$ No tidal disruption



*ISCO = Inner Stable Circular Orbit

Key ingredients of the mass ejection in BH-NS are

- ▶ Spin of BH
- ▶ Mass ratio ($M_{\text{BH}}/M_{\text{NS}}$)
- ▶ Compactness of NS ($M_{\text{NS}}/R_{\text{NS}}$)




Stiff EOS = large compactness

Compactness




Ab initio simulation of BH-NS (K. Hayashi, KK et al. 22,23)

Time: 0.01 ms




Numerical modeling of BH-NS merger

Magnetically tower “jet”



Isotropic Poynting Luminosity



- Magnetically tower “jet” builds up magnetosphere
⇒ L_{iso} and θ_{jet} are roughly consistent with the observed values.