

Red and Slow:

Nebular Phase Kilonovae in the Infra-Red

Quentin Pognan 18.02.2026
quentin.pognan@aei.mpg.de




MAX PLANCK INSTITUTE
FOR GRAVITATIONAL PHYSICS
(Albert Einstein Institute)




With K. Kawaguchi, S. Wanajo,
S. Fujibayashi,
A. Jerkstrand, J. Grumer

Artist Impression, NASA




Coalescence,
r-process in
ejecta




Watson+ 2019




Hotokezaka+ 2022



Evolution of a Kilonova



Hotokezaka+ 2023




Gillanders & Smartt+ 2025

Nebular phase:


- Optically thin
- NLTE physics
- Slow evolution of T and X_e
- Low energy $\rightarrow$ NIR-MIR lines

Method and Goals


- Use time-dependent NLTE radiative transfer simulations to explore a parameter space of nebular phase KN spectra in the IR: 1 – 30 μm
 - Input data from end-to-end simulation of symmetric binary NS merger with long-lived remnant, ejecta split into post-merger and dynamical components
- Parameter space of total ejecta mass and fraction of dynamical ejecta
 - $M_{\text{ej}} = 0.05, 0.01, 0.005 M_{\text{sol}}$, $f_{\text{dyn}} = 0, 0.01, 0.05, 0.1, 0.5$
- Produce self-consistent nebular phase spectra of KNe
 - Particular focus on the impact of lanthanides in the IR



Numerical Relativity
Fujibayashi+ 2020




GR Hydrodynamics
Kawaguchi+ 2021



NLTE Radiative Transfer
Jerkstrand+ 2011,2012
QP+ 2022


Merger Model

- DD2 EoS, $1.35 M_{\text{sol}}$ symmetric BNS system
 - Long lived remnant scenario
- No lanthanides, 3rd peak or actinides in post-merger ejecta
 - In reality small trace abundances
 - Not impactful in terms of opacity or emission lines
- Post-merger vs dynamical ejecta cutoff at 0.15 s
 - Consistency between 3D NR and 2D HD simulation
- Compositions limited to 30 elements, chosen at 25 days
 - Representative of average composition in 10 – 75 day timespan






Spectra: Common Features

- “Standard” model with $M_{\text{ej}} = 0.05 M_{\text{sol}}$ and $f_{\text{dyn}} = 0.05 \rightarrow X_{\text{La}} = 0.0027$
- Key features at 10 days:
 - Very blended emission at ≤ 2 micron
 - Dominant lanthanides are Nd II and Ce III
 - Te III at 2 micron is blended with these
 - Strong Br I/II emission at 3 micron
 - Blend of Se I/III between 4 – 6 micron
 - Some Ni III emission at 7.3 micron (from stable Ni isotopes)





Time Evolution

- The spectra evolve in time, rapidly in the 10 – 30 day range and more slowly after
 - This is linked to the slow evolution of T and X_e
- Initially we have increasing T and X_e
 - Change of dominant ionisation state leads to observable changes
 - Se I $\rightarrow$ Se III leads to a transition from a single to double peak feature
- At late time, time-dependent effects lead to freeze-out of X_e




Impact of Ejecta Mass

- Smaller ejecta masses yield lower densities at a given time
 - Line cooling is suppressed due to inefficient collisions → typically hotter
 - Both ionisation and recombination are inhibited → similar $X_e \sim 0.7-1.5$
- Spectral evolution is “compressed” in time
 - E.g. $0.01M_{\text{sol}}$ model at 20 days resembles $0.05M_{\text{sol}}$ model at ~ 40 days
- Freeze-out of X_e occurs earlier
 - SED shape does not evolve much from this point onwards




Lanthanide-poor Case

- Consider lanthanide-poor ejecta
 - E.g. $f_{\text{dyn}} = 0.01$ model with $X_{\text{La}} = 0.001$
c.f. $f_{\text{dyn}} = 0.05$ with $X_{\text{La}} = 0.0027$
- Some emission from Ce III and Nd II at ≤ 3 micron
- Compare to fiducial lanthanide-free model $f_{\text{dyn}} = 0$, and $f_{\text{dyn}} = 0.05$
 - Relative contributions are different, e.g. 2 micron blended feature but...
- Overall spectral shape is very similar
 - Unlikely to observationally distinguish cases with $X_{\text{La}} \leq 0.0027$ in nebular phase at IR wavelengths




Lanthanide-rich Case

- Consider lanthanide-rich ejecta
 - E.g. $f_{\text{dyn}} = 0.5$ model with $X_{\text{La}} = 0.0260$
c.f. $f_{\text{dyn}} = 0.05$ with $X_{\text{La}} = 0.0027$
- At 10 days, lanthanides completely dominate
 - Te III at 2 micron is largely subdominant
 - No characteristic Se I emission peak at 5 micron
- Due to large f_{dyn} , ejecta has high heating rate $\rightarrow$ noticeably brighter
 - Combined with an accurate determination of M_{ej} , case should be distinguishable from cases with $X_{\text{La}} < 0.026$





Lanthanide-rich Case

- At high enough X_{La} , lanthanide emission remains significant
 - At late times
 - At smaller masses
- Note that the Se III double peak feature does appear with time
 - Heavier composition $\rightarrow$ W III contributing up to $\sim 20\%$ of emission at 4.5 micron
- Spectral shape past 4 micron is similar to more mildly lanthanide-bearing models
 - Slightly brighter due to higher power, but difficult to distinguish observationally




Further into the MIR



- Further into the MIR: mainly 1st, 2nd peak and lighter elements, little to no lanthanides
- Important Ni III line at 11 micron, could be combined with line at 7.3 micron for reliable species ID
 - But luminosity is significantly lower!

Generalising Lanthanide Emission


- Where and when does lanthanide emission dominate in the IR in the nebular phase?
 - Typically at wavelengths < 4 micron
 - Earlier times/heavier masses $\rightarrow$ higher density
- If looking for lanthanides:
 - Earlier is better
 - “Blue” NIR wavelengths ≤ 4 micron
 - But heavy blending is expected...
- If avoiding lanthanides:
 - Wavelengths ≥ 4 micron
 - Later is better
 - But remember luminosity is dropping



Flux fraction coming from lanthanide species in the 1.2 – 3.6 micron range


What about Te?

- Te III has been suggested as responsible for emission at 2 micron in AT2017gfo and AT2023vfi
- These models suggest that Te III does indeed play a non-negligible role...
- But it is never the sole emitting species in any model at any epoch
 - In lanthanide-bearing models, blending with Ce III and Nd II occurs
 - In lanthanide-poor models, blending with Zr II, Kr II and Kr III occurs
- Estimating Te mass assuming the emission is fully Te III overestimates (potentially drastically) the value
 - Highlights need for two separate lines from same species for reliable identification and mass calculations




1st or 3rd: Se vs W

- *Spitzer* photometry of AT2017gfo at 43 and 74 days shows emission at 4.5 micron and non-detection at 3.6 micron
- Suggested species for 4.5 micron emission include Se III and W III
 - Our models favour Se III since underlying merger model is long-lived remnant case
 - Dominance of W III requires significantly more neutron rich conditions
- Our $f_{\text{dyn}} = 0.01$ model ($X_{\text{La}} = 0.001$) most closely matches the observed photometry
 - Exploration of short-lived remnant cases and possibly BHNS mergers should be explored




Cool and Mysterious: AT2023vfi

- AT2023vfi spectra at 29 and 61d showed cool, Blackbody like continua
 - E.g. 29d: $T \sim 660\text{K}$, $R \sim 6 \times 10^{15} \text{ cm}$ ($v_{\text{phot}} = 0.08c$)
- Origin of this continuum is unclear
 - Photosphere? Dust? Something else?
- Can expansion line opacity make an optically thick BB continuum?
 - Our models are optically thin in NIR at these times
 - Take most lanthanide-rich $f_{\text{dyn}} = 0.5$ model and make a “dense” variation with $v_{\text{in}} = 0.02c \rightarrow$ no continuum formed
- Expansion line opacity cannot form an optically thick NIR continuum at these times in reasonable ejecta conditions




AT2023vfi: Lines

- AT2023vfi also had GRB afterglow and emission line components
 - For comparison to models, subtract best-fit afterglow and BB components
- Case A: BB component is a real “photosphere” with $v_{\text{phot}} = 0.08c$
 - Scale down model spectra corresponding to mass between $v_{\text{in}} = 0.05$ - $v_{\text{phot}} = 0.08c$
- Closest match is $f_{\text{dyn}} = 0.5$ model
 - 4.5 micron emission is mix of Se III and W III
 - 2 micron emission does not reproduce observed line shape
 - Excess flux between 2.5 – 4.0 micron




AT2023vfi: Lines

- Case B: Assume BB continuum comes from something other than a photosphere
 - In this case don't scale down model spectra, simply compare to continuum-subtracted spectrum
- Closest match is $f_{\text{dyn}} = 0.01$ model
 - 4.5 micron emission is mainly Se III
 - Te III at 2.1 micron, but not luminous enough
 - Again flux excess between 2.5 – 3.6 micron
- No model reproduces the 29d spectrum of AT2023vfi...



AT2023vfi: Remnant?

- In all models, we find excess Br I/II flux at ~ 3 micron
 - Is Br absent from line-forming ejecta of AT2023vfi at 29d?
- Suppose it is... let's follow the thread
 - Br is a first peak element $\rightarrow$ other first peak elements also low abundance?
 - Se may also be absent $\rightarrow$ 4.5 micron emission is dominated by W III?
- Requisite W III mass is $\sim 9 \times 10^{-4} M_{\text{sol}}$ (McCann+2025)
 - Only $1.4 \times 10^{-4} M_{\text{sol}}$ in $f_{\text{dyn}} = 0.5$ model at 30d
- Our merger model is for long-lived remnant case, could AT2023vfi instead be a short-lived case?
 - Further work needed... but AT2023vfi likely not a long-lived remnant case



McCann+ 2025

Conclusions

- Want to explore IR spectra of KNe in the nebular phase
- Use time-dependent NLTE radiative transfer simulations to explore a parameter space of ejecta mass and lanthanide fraction
- Find several key results:
 - Lanthanide impact is typically limited to wavelengths ≤ 4 micron
 - Te III at 2.1 micron is always blended: Ce III, Nd II, Zr II, Kr II, Kr III
 - Se III dominates emission at 4.5 micron and 5.7 micron
 - Ni III has important features at 7.3 and 11 micron
 - Observations combining NIR and MIR spectroscopy could provide 2 line IDs for 2 different species!
- Comparing our model to AT2023vfi at 29d:
 - Line opacity alone cannot produce an optically thick thermal continuum
 - No model reproduces the observed line emission → some hints that a long-lived remnant case is in tension with spectra of AT2023vfi

**Thank you for your
attention!**

