

MINOS anomaly and non-standard interactions etc.

--- Critical views ---

Osamu Yasuda
Tokyo Metropolitan University

Sept. 23 @6th IDS-NF Mtg

1. Introduction

2. High energy behavior of ν_{atm} data & NSI

3. MINOS anomaly & NSI etc.

4. Conclusions

1. Introduction

Candidates for new physics to be tested at future LBL

Scenarios	Phenomenological bound on deviation from standard case
NSI at production / detection	$O(1\%)$
NSI in propagation $\varepsilon_{\mu\alpha}$	$O(1\%)$
NSI in propagation $\varepsilon_{[e\tau][e\tau]}$	$O(100\%)$
Violation of unitarity due to heavy particles	$O(0.1\%)$
Light sterile neutrinos	$O(10\%)$

● NP in propagation (NP matter effect)

$$i \frac{d}{dt} \begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \left[U \text{diag}(E_1, E_2, E_3) U^{-1} + A \begin{pmatrix} 1 + \epsilon_{ee} & \epsilon_{e\mu} & \epsilon_{e\tau} \\ \epsilon_{\mu e} & \epsilon_{\mu\mu} & \epsilon_{\mu\tau} \\ \epsilon_{\tau e} & \epsilon_{\tau\mu} & \epsilon_{\tau\tau} \end{pmatrix} \right] \begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix}$$

$$A \equiv \sqrt{2} G_F N_e \quad N_e \equiv \text{electron density}$$

NP

● Constraints on $\epsilon_{\alpha\beta}$

Davidson et al., JHEP 0303:011,2003; Berezhiani, Rossi, PLB535 ('02) 207; Barranco et al., PRD73 ('06) 113001; Barranco et al., arXiv:0711.0698

Biggio et al., JHEP 0908, 090 (2009) w/o 1-loop arguments

related to each other by ν_{atm}

can be improved by ν_{atm}


$$\left(\begin{array}{l} |\epsilon_{ee}| \lesssim 4 \times 10^0 \\ |\epsilon_{e\mu}| \lesssim 3 \times 10^{-1} \\ |\epsilon_{\mu\mu}| \lesssim 7 \times 10^{-2} \\ |\epsilon_{e\tau}| \lesssim 3 \times 10^0 \\ |\epsilon_{\mu\tau}| \lesssim 3 \times 10^{-1} \\ |\epsilon_{\tau\tau}| \lesssim 2 \times 10^1 \end{array} \right)$$

● Sterile neutrinos: (3+1)-scheme

If we forget about **LSND/MB anti- ν** , then (3+1)-scheme is a possible scenario, provided that the mixing angles satisfy all the constraints of the negative results (w/ less motivation).

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \\ \nu_s \end{pmatrix} = U \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \\ \nu_4 \end{pmatrix}$$

$$U = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} & U_{e4} \\ U_{\mu1} & U_{\mu2} & U_{\mu3} & U_{\mu4} \\ U_{\tau1} & U_{\tau2} & U_{\tau3} & U_{\tau4} \\ U_{s1} & U_{s2} & U_{s3} & U_{s4} \end{pmatrix}$$



$$U = R_{34}(\theta_{34}, 0) R_{24}(\theta_{24}, 0) R_{23}(\theta_{23}, \delta_3) R_{14}(\theta_{14}, 0) R_{13}(\theta_{13}, \delta_2) R_{12}(\theta_{12}, \delta_1)$$

θ_{34} : ratio of $\nu_\mu \leftrightarrow \nu_\tau$ and $\nu_\mu \leftrightarrow \nu_s$ in ν_{atm}

θ_{24} : ratio of $\sin^2\left(\frac{\Delta m_{\text{atm}}^2 L}{4E}\right)$ and $\sin^2\left(\frac{\Delta m_{\text{SBL}}^2 L}{4E}\right)$ in ν_{atm}

θ_{14} : mixing angle in ν_{reactor} at $L=O(10\text{m})$



Constraints from ν_{atm} and SBL

Donini-Maltoni-Meloni-Migliozzi-Terranova, JHEP 0712:013,'07

$$U = R_{34}(\theta_{34}) R_{24}(\theta_{24}) R_{23}(\theta_{23}, \delta_3) R_{14}(\theta_{14}) R_{13}(\theta_{13}, \delta_2) R_{12}(\theta_{12}, \delta_1)$$

Assumption on rapid oscillations in ν_{atm} :
 $\Delta m^2_{41} > 0.1 \text{ eV}^2$

θ_{34} : could be relatively large




(3+1)-scheme

$$P(\bar{\nu}_e \rightarrow \bar{\nu}_e) = 1 - 4|U_{e4}|^2(1 - |U_{e4}|^2) \sin^2(\Delta m_{41}^2 L/4E)$$

$$P(\nu_\mu \rightarrow \nu_\mu) = 1 - 4|U_{\mu 4}|^2(1 - |U_{\mu 4}|^2) \sin^2(\Delta m_{41}^2 L/4E)$$

$$\sin^2 2\theta_{\text{Bugey}} > 4|U_{e4}|^2(1 - |U_{e4}|^2) = \sin^2 2\theta_{14}$$

$$\sin^2 2\theta_{\text{CDHSW}} > 4|U_{\mu 4}|^2(1 - |U_{\mu 4}|^2) \cong \sin^2 2\theta_{24}$$



2. High energy behavior of ν_{atm} data & NP

- Standard case with $N_\nu=2$

$$1 - P(\nu_\mu \rightarrow \nu_\mu) = \sin^2 2\theta_{\text{atm}} \sin^2 \left(\frac{\Delta m_{\text{atm}}^2 L}{4E} \right) \propto \frac{1}{E^2}$$

- Standard case with $N_\nu=3$

$$1 - P(\nu_\mu \rightarrow \nu_\mu) \sim \left(\frac{\Delta m_{31}^2}{2AE} \right)^2 \left[\sin^2 2\theta_{23} \left(\frac{c_{13}^2 AL}{2} \right)^2 + s_{23}^2 \sin^2 2\theta_{13} \sin^2 \left(\frac{AL}{2} \right) \right] \propto \frac{1}{E^2}$$

- Deviation of $1-P(\nu_\mu \rightarrow \nu_\mu)$ due to NP contradicts with data

$$1 - P(\nu_\mu \rightarrow \nu_\mu) \simeq \mathbf{c}_0 + \frac{\mathbf{c}_1}{E} + \frac{c_{20}L^2 + c_{21} \sin^2(c_{22}L)}{E^2}$$

→ High ν_{atm} data gives constraints on NP:

$$|\mathbf{c}_0| \ll 1, |\mathbf{c}_1| \ll 1$$

● with NSI

$$1 - P(\nu_\mu \rightarrow \nu_\mu) \simeq \mathbf{c}_0 + \frac{\mathbf{c}_1}{E} + \frac{c_{20}L^2 + c_{21} \sin^2(c_{22}L)}{E^2}$$

$$|\mathbf{c}_0| \ll 1 \rightarrow |\varepsilon_{e\mu}| \ll 1, |\varepsilon_{\mu\mu}| \ll 1, |\varepsilon_{\mu\tau}| \ll 1$$

$|\varepsilon_{\mu\tau}| \ll 1$: Already shown by Fornengo et al. PRD65, 013010, '02;
Gonzalez-Garcia&Maltoni, PRD70, 033010, '04; Mitsuka@nufact08

$|\varepsilon_{\mu\mu}| \ll 1$: Already shown from other expts. by Davidson et al.
JHEP 0303:011, '03

$|\varepsilon_{e\mu}| \ll 1$: New observation (analytical consideration only)

$$|\mathbf{c}_1| \ll 1 \rightarrow |\varepsilon_{\tau\tau} - |\varepsilon_{e\tau}|^2 / (1 + \varepsilon_{ee})| \ll 1$$

Already shown by
Friedland-Lunardini,
PRD72:053009, '05

- Summary of the constraints on $\epsilon_{\alpha\beta}$

To a good approximation, we are left with 3 independent variables ϵ_{ee} , $|\epsilon_{e\tau}|$, $\arg(\epsilon_{e\tau})$:

$$A \begin{pmatrix} 1 + \epsilon_{ee} & \epsilon_{e\mu} & \epsilon_{e\tau} \\ \epsilon_{\mu e} & \epsilon_{\mu\mu} & \epsilon_{\mu\tau} \\ \epsilon_{\tau e} & \epsilon_{\tau\mu} & \epsilon_{\tau\tau} \end{pmatrix}$$




$$A \begin{pmatrix} 1 + \epsilon_{ee} & 0 & \epsilon_{e\tau} \\ 0 & 0 & 0 \\ \epsilon_{e\tau}^* & 0 & |\epsilon_{e\tau}|^2 / (1 + \epsilon_{ee}) \end{pmatrix}$$

Furthermore, v_{atm} data implies

$$\tan\beta = |\epsilon_{e\tau}| / (1 + \epsilon_{ee}) < 1.3$$

Friedland-Lunardini,
PRD72:053009,'05

Allowed region in $(\epsilon_{ee}, |\epsilon_{e\tau}|)$




● with ν_s

$$1 - P(\nu_\mu \rightarrow \nu_\mu) \simeq \mathbf{C_0} + \frac{\mathbf{C_1}}{E} + \frac{c_{20}L^2 + c_{21} \sin^2(c_{22}L)}{E^2}$$

$$|\mathbf{C_0}| \propto s_{24}^2 \ll 1 \rightarrow s_{24}^2 \ll 1$$




$$|\mathbf{C_1}| \propto s_{34}^2 \ll 1 \rightarrow s_{34}^2 \ll 1$$



Donini-Maltoni-Meloni-Migliozzi-
Terranova, JHEP 0712:013,'07

3. MINOS anomaly


Vahle@nu2010



Effort to explain with $\epsilon_{\mu\tau}$

Mann-Cherdack-Musial-Kafka,
arXiv:1006.5720 [hep-ph]

$$A \begin{pmatrix} 1 & 0 & 0 \\ 0 & \epsilon_{\mu\mu} & \epsilon_{\mu\tau} \\ 0 & \epsilon_{\tau\mu} & \epsilon_{\tau\tau} \end{pmatrix}$$



ν_{atm} constraint


Contradicts with ν_{atm} constraint

Effort to explain with $\epsilon_{\mu\tau}$

Kopp-Machado-Parke, arXiv:1009.0014 [hep-ph]

ν_{atm} constraint

$$A \begin{pmatrix} 1 & 0 & 0 \\ 0 & \epsilon_{\mu\mu} & \epsilon_{\mu\tau} \\ 0 & \epsilon_{\tau\mu} & \epsilon_{\tau\tau} \end{pmatrix}$$



Contradicts with ν_{atm} constraint

Effort to explain with gauging $L_\alpha-L_\beta$

Heeck-Rodejohann, arXiv:1007.2655 [hep-ph]

$$A \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} V & 0 & 0 \\ 0 & -V & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} V & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -V \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 & 0 \\ 0 & V & 0 \\ 0 & 0 & -V \end{pmatrix}$$

L_e-L_μ L_e-L_τ $L_\mu-L_\tau$

ν_{sol} constraint

$$\left. \frac{\Delta m_{31}^2}{E} \right|_{\text{MINOS}} \sim A \sim V_{\text{MINOS}} = \frac{\text{const}}{d_{\text{Sun-Earth}}}$$

To keep the success of ν_{sol} ,
we have to avoid L_e-L_μ , L_e-L_τ

$$100 \times V_{\text{MINOS}} \sim 100 \times \left. \frac{\Delta m_{31}^2}{E} \right|_{\text{MINOS}}$$

$$\sim \left. \frac{\Delta m_{21}^2}{E} \right|_{\text{solar}} \ll V_{\text{solar}} = \frac{\text{const}}{R_{\text{Sun}}}$$

ν_{atm} constraint

$L_\mu-L_\tau$ contradicts with $|\varepsilon_{\mu\mu} - \varepsilon_{\tau\tau}| \ll 1$

Effort to explain with $\epsilon_{e\tau}$

$$A \begin{pmatrix} 1 + \epsilon_{ee} & 0 & \epsilon_{e\tau} \\ 0 & 0 & 0 \\ \epsilon_{e\tau}^* & 0 & |\epsilon_{e\tau}|^2 / (1 + \epsilon_{ee}) \end{pmatrix}$$

Preliminary work by OY,
unpublished (2010; to be published
or not to be, that is a question)

● Best fit point lies in
the excluded region of


ν_{atm}

● $\chi^2(\text{SM}) - \chi^2(\text{min}) = 0.4$

(2dof): 0.25σ (not
significant at all)

→ Probably not worth



introducing $\epsilon_{e\tau}$



Effort to explain with ν_s or ν_s +gauged B-L

Engelhardt-Nelson-Walsh, Phys.Rev.D81:113001,2010


- old data is used
- $\theta_{24}=0 \rightarrow$ no conflict with CDHSW
- $\theta_{34} \neq 0 \rightarrow 0 \leq \theta_{34} < 30^\circ$
- $\theta_{23} = \pi/4$




$$\begin{pmatrix} \pm(V_{CC} - V_{NC} - V_{B-L}) & 0 & 0 & 0 \\ 0 & \pm V_{NC} - V_{B-L} & 0 & 0 \\ 0 & 0 & \pm V_{NC} - V_{B-L} & 0 \\ 0 & 0 & 0 & V_{B-L} \end{pmatrix}$$

Best fit


- with B-L $\Delta m^2_{32} = 2.5 \times 10^{-3} \text{eV}^2$, $\Delta m^2_{42} = 2.5 \times 10^{-2} \text{eV}^2$, $V_{NC}/2 + V_{B-L} = 5 \times 10^{-14} \text{eV}$, $\chi^2 = 24.8$ (20 dof)
- w/o B-L $\Delta m^2_{32} = 1.8 \times 10^{-3} \text{eV}^2$, $\Delta m^2_{42} = 9.5 \times 10^{-2} \text{eV}^2$, $\chi^2 = 28.1$ (21 dof)



(a) M vs. m



(b) V vs. m



(c) V vs. M

$$M = (\Delta m_{42}^2)^{1/2}$$

$$m = (\Delta m_{32}^2)^{1/2}$$

$$V = V_{\text{NC}}/2 + V_{\text{B-L}}$$

Engelhardt-Nelson-Walsh,
Phys.Rev.D81:113001,2010

Effort to explain with ν_s

Preliminary work by OY,
unpublished (2010; to be published
or not to be, that is a question)

- new data is used

- Δm^2_{42} is fixed as 1eV^2

- $\theta_{24}=0 \rightarrow$ no conflict with CDHSW

- potential enhancement for ν / suppression for $\bar{\nu}$ occurs if $\theta_{34} < \pi/4$ & NH or $\theta_{34} > \pi/4$ & IH

$$\tan 2\tilde{\theta}_{23} = \frac{\Delta E_{32} \sin 2\theta_{23}}{\Delta E_{32} \cos 2\theta_{23} \pm V s_{34}^2}$$

$$\sin^2 2\tilde{\theta}_{23} = \frac{(\Delta E_{32} \sin 2\theta_{23})^2}{(\Delta E_{32} \cos 2\theta_{23} \pm V s_{34}^2)^2 + (\Delta E_{32} \sin 2\theta_{23})^2}$$

$$1 - P(\nu_\mu \rightarrow \nu_\mu) = \sin^2 2\tilde{\theta}_{23} \sin^2 \left(\frac{\Delta \tilde{E}_{32} L}{2} \right)$$

$$\Delta \tilde{E}_{32} \equiv \sqrt{(\Delta E_{32} \cos 2\theta_{23} \pm V s_{34}^2)^2 + (\Delta E_{32} \sin 2\theta_{23})^2}$$

$\sin^2 2\tilde{\theta}_{23}$ is almost 1 anyway \rightarrow difficult to distinguish ν & $\bar{\nu}$

(Best fit point with ν_s) \simeq (Best fit point for $N_\nu=3$ case)

\rightarrow Probably not worth introducing ν_s

4. Conclusions

- People made efforts to account for MINOS anomaly, but they all seem either to give little contribution to distinguish ν & $\bar{\nu}$ or to have conflict with atmospheric neutrinos and/or solar neutrinos.
- After all, MINOS anomaly is only a 2σ effect, so we should wait until we have more statistics.